

MAT415 Lecture Notes

Arky!! :3c

'26 Fall Semester

Contents

Day 1: Gaussian Integers (Sep. 8, 2026)	2
Day 2: Algebra Preliminaries; Algebraic Numbers and Integers (Sep. 10, 2026)	4
Day 3: Minimal Polynomials (Sep. 15, 2026)	8
Day 4: Number Fields (Sep. 17, 2026)	10
Day 5: Norm and Trace (Sep. 22, 2026)	14
Day 6: Integral Basis (Sep. 24, 2026)	16
Day 7: Dedekind Domains (Sep. 29, 2026)	20
Day 8: Properties of Dedekind Domains, Ideal Class Group (Oct. 1, 2026)	22
Day 9: Ramification (Oct. 6, 2026)	25

Day 1: Gaussian Integers (Sep. 8, 2026)

The Quercus site for this class will be up tonight or early tomorrow. This class will have (approximately) weekly homework. There will be regular oral quizzes with the TA, about 5–10 minutes, where you will discuss one of the problems on the homework.

Consider the following example,

$$1 + 1 = 2, \quad 1 + 4 = 5, \quad 4 + 9 = 13, \quad 1 + 16 = 17.$$

Theorem 1.1 (Fermat). Let $p > 2$; p can be expressed as a sum of two squares if and only if $p \equiv 1 \pmod{4}$.

Proof. We start with the easy direction. Observe that the squares of the numbers $0, 1, 2, 3, \dots$ are all equal to 0 or 1 modulo 4, i.e., $0^2 = 0$, $1^2 = 1$, $2^2 = 4 \equiv 0 \pmod{4}$, and $3^2 = 9 \equiv 1 \pmod{4}$, and so on (you can always express a number as either $2n$ or $2n + 1$, then squaring, to check that this pattern continues). It follows that if $p = n^2 + m^2$, then $p \equiv 1 \pmod{4}$; if it were the sum of the squares of two odd numbers, then p would be even, which contradicts it being prime.

For the other direction, consider the ring of Gaussian integers,

$$\mathbb{Z}[i] = \{a + bi \mid a, b \in \mathbb{Z}\} \subset \mathbb{C}.$$

Consider $p = n^2 + m^2$. Notice that the right hand side cannot be factored as *integers*, but it could be factored in $\mathbb{Z}[i]$ as $(n + im)(n - im)$; in this manner, we may instead ask the question of how to factor $p \in \mathbb{Z}[i]$. The continuation of this proof will use the fact that $\mathbb{Z}[i]$ is a UFD, but we will not prove that today. In particular, $\mathbb{Z}[i]$ is a Euclidean domain with $\alpha \mapsto |\alpha|^2$ is a Euclidean norm. $\square$

Given $z = x + iy \in \mathbb{Z}[i]$, define the norm of z as $N(z) = z \cdot \bar{z} = x^2 + y^2 \in \mathbb{Z}_{\geq 0}$.¹ Observe that we have $N(z \cdot w) = N(z) \cdot N(w)$.

Claim 1.2. $z \in \mathbb{Z}[i]$ is a unit if and only if $N(z) = 1$, i.e., the units are $\{\pm 1, \pm i\}$.

Proof. Suppose $N(z) = 1$; by definition, we see $z \cdot \bar{z} = 1$, and so $\bar{z}$ constitutes the inverse of z . Conversely, suppose $z \in \mathbb{Z}[i]$ is a unit; if there exists another number w such that $z \cdot w = 1$, then, since norm is multiplicative, we may take the norm of both sides to get that $N(z) \cdot N(w) = 1$. Since $N(z) \in \mathbb{Z}_{\geq 0}$, it must be that $N(z) = 1$. $\square$

Claim 1.3. If p is a prime such that $p \equiv 1 \pmod{4}$, then p is not prime in $\mathbb{Z}[i]$.

We will prove the claim later; for now, observe that 1.1 and 1.3 imply each other; one is obvious, and for the direction that the claim implies the theorem, suppose p is not prime, so we may write it as the product of two non-units, $p = \alpha \cdot \beta$, where $\alpha, \beta \in \mathbb{Z}[i]$. We have that $p^2 = N(p) = N(\alpha) \cdot N(\beta)$; but $N(\alpha), N(\beta) \neq 1$, so it must be that $N(\alpha) = N(\beta) = p$. By writing $\alpha = a + bi$, we get that $N(\alpha) = a^2 + b^2 = p$, which is the desired expression as seen in 1.1.

We now prove 1.3 in two ways. For the first way, consider Wilson's theorem:

Theorem 1.4 (Wilson). Let p be prime; then $(p - 1)! \equiv -1 \pmod{p}$.

Proof. The proof of this is simple; since $\mathbb{Z}/p$ is a field, every nonzero residue modulo p has an inverse, where $\alpha \equiv \alpha^{-1}$ if and only if $\alpha \equiv 1$ or -1 , i.e., $\alpha^2 = 1$. It follows that the product of $1, \dots, p - 1$ without 1 and $p - 1$ is 1, so we are done. $\square$

¹at this point, Rozenblyum made a comment about complex number norms versus algebra norms. idk. sure.

Corollary 1.5. If $p = 4n + 1$, then $((2n)!)^2 \equiv -1 \pmod{p}$.

Proof. By Wilson's theorem, we have that $-1 \equiv (p-1)!$. Let us write the factorial out;

$$-1 \equiv (p-1)! = (1 \cdot 2 \cdot \dots \cdot 2n)((p-2n)(p-2n+1) \dots (p-1)).$$

Since we're working modulo p , we may realize the right hand side expression as

$$\dots \equiv (1 \cdot 2 \cdot \dots \cdot 2n)((-2n)(-2n+1) \dots (-1)) = (2n)!(-1)^{2n}(2n)! = ((2n)!)^2. \quad \square$$

The above corollary allows us to write²

$$p \mid ((2n)!)^2 + 1,$$

which we may factor as $((2n)! + i)((2n)! - i)$; if p was prime, then it divides one of them, but we know that $p \mid (a+bi) = (2n)! \pm i$ can't happen.

We now proceed to the second proof. Consider the cyclic group

$$(\mathbb{Z}/p)^\times = \mathbb{Z}/(p-1);$$

if $p = 4n + 1$, then $\mathbb{Z}/(p-1) \cong \mathbb{Z}/(4n)$, whence the existence of some element of order 4 in $(\mathbb{Z}/p)^\times$. Let such an element be called m ; it follows that $m^4 \equiv 1 \pmod{p}$ and $m^2 \equiv -1 \pmod{p}$, but then we would have $p \mid m^2 + 1$, which is a contradiction. While this proof was fast, it didn't really tell us much about the arithmetic that went on in the first proof.

The morals as to the two above proofs are that

- (i) It is important to understand arithmetic in more general number rings (such as prime factorization). As an example,

$$\mathbb{Z}[\sqrt{-5}] = \{a + b\sqrt{-5} \mid a, b \in \mathbb{Z}\}$$

is not a UFD (see the MAT347 notes for a reference on this).

- (ii) Questions about number rings are closely related to questions in modular arithmetic.

²Rozenblyum proceeded to stand stunned in front of the blackboard for 3 minutes. When pointed out on how to continue the discussion, he said that this was proof your "IQ drops precipitously when in front of a classroom".

Day 2: Algebra Preliminaries; Algebraic Numbers and Integers (Sep. 10, 2026)

We set some conventions for this class. Unless specified otherwise, a ring will always refer to a commutative ring with unity.

Definition 2.1 (Preliminaries). Let R be a ring.

- (i) R is an *integral domain* if it admits no zero divisors, i.e., whenever $ab = 0$ for $a, b \in R$, then $a = 0$ or $b = 0$.
- (ii) We will write $a \mid b$ if “ a divides b ” for $a, b \in R$ if there exists some $c \in R$ such that $b = ac$.
- (iii) An integral domain R is a *principal ideal domain* (PID) if every ideal is principal, i.e., generated by a single element. Specifically, if $I \subset R$ is an ideal, then there exists $a \in R$ such that $I = \langle a \rangle$.
- (iv) An element $u \in R$ is a *unit* if it has a multiplicative inverse. An element $p \in R$ is *irreducible* if it is not a unit, and whenever $p = ab$, either a or b is a unit. An element $p \in R$ is *prime* if, whenever $p \mid ab$, then $p \mid a$ or $p \mid b$.
- (v) An integral domain R is a *unique factorization domain* (UFD) if every non-unit $x \in R$ is uniquely (up to permutation and multiplication by units) a product of irreducible elements.

Proposition 2.2. If R is a UFD, then irreducibles are primes, and primes are irreducible.

Note that primes are always irreducible. You can reference my MAT347 notes for a proof on this.

Proposition 2.3. Let R be a ring. Then $p \in R$ is prime if and only if $R/\langle p \rangle$ is an integral domain.

Proof. Pick $\bar{a}, \bar{b} \in R/\langle p \rangle$ such that $\bar{a}\bar{b} = 0$ in $R/\langle p \rangle$. Then $ab \in \langle p \rangle$, so $p \mid ab$, where $p \mid a$ or $p \mid b$ as p is prime, so $\bar{a} = 0$ or $\bar{b} = 0$. The converse direction is obtained by reversing this proof. $\square$

We now continue writing out the definitions.

Definition 2.4 (Preliminaries, cont.). Let R be a ring.

- (vi) An integral domain R is a *Euclidean domain* (ED) if there exists a map $v: R \setminus \{0\} \rightarrow \mathbb{Z}_{\geq 0}$ such that, for all $a, b \in R$ with $b \neq 0$, there exists $q, r \in R$ such that $a = bq + r$ with $r = 0$ or $v(r) < v(b)$. In addition, for all nonzero $a, b \in R$, v must satisfy $v(a) \leq v(ab)$.

Theorem 2.5. Let R be an integral domain. If R is a Euclidean domain, then it is a PID. If R is a PID, then it is a UFD.

Once again, the proof is in my MAT347 notes. Let us give some examples of this.

- $R = \mathbb{Z}$ is a Euclidean domain with $v(n) = |n|$.
- If F is a field, then $F[x]$ is a Euclidean domain with $v(f(x)) = \deg f$.
- Let n be an integer, and let $R = \mathbb{Z}/n$ be the integers mod n . R is an integral domain if and only if n is prime; in fact, if n is prime, then $\mathbb{Z}/n$ is a field. For a quick proof of this fact, let $x \in \mathbb{Z}$; then $\bar{x} \in \mathbb{Z}/n$ is a unit if and only if x is coprime to n . If p is prime, then $(\mathbb{Z}/p)^\times$ is a group of order $p - 1$.

Theorem 2.6. Let $p \in \mathbb{N}$ be a prime number.

- (i) (Fermat's little theorem) If $a \in \mathbb{Z}$ is coprime to p , then $a^{p-1} \equiv 1 \pmod{p}$.
- (ii) $(\mathbb{Z}/p)^\times$ is a cyclic group.

Corollary 2.7. Let $p \in \mathbb{N}$ be a prime number. Then there exists $g \in \mathbb{Z}$, called the *primitive root modulo* p , such that (i) if a is coprime to p , then $a \equiv g^r \pmod{p}$ for some $r \geq 0$, and (ii) if $r \in \mathbb{Z}$ such that $g^r \equiv 1 \pmod{p}$, then $p - 1 \mid r$.

Definition 2.8. An *algebraic number* α is a complex number $\alpha \in \mathbb{C}$ for which there exists $a_0, \dots, a_{n-1} \in \mathbb{Q}$ (with $n \geq 1$) such that

$$\alpha^n + a_{n-1}\alpha^{n-1} + \dots + a_1\alpha + a_0 = 0,$$

i.e., $p(\alpha) = 0$ for some monic $p(x) \in \mathbb{Q}[x]$. An *algebraic integer* is a complex number $\alpha \in \mathbb{C}$ such that there exists $a_0, \dots, a_{n-1} \in \mathbb{Z}$ (with $n \geq 1$) such that

$$\alpha^n + a_{n-1}\alpha^{n-1} + \dots + a_1\alpha + a_0 = 0,$$

i.e., $p(\alpha) = 0$ for some monic $p(x) \in \mathbb{Z}[x]$. Algebraic numbers are denoted by $\mathbb{Q}^{\text{alg}}$ or $\overline{\mathbb{Q}}$, and algebraic integers are denoted by $\mathbb{Z}^{\text{alg}}$ or $\overline{\mathbb{Z}}$. The bar denotes algebraic closure.

In particular, algebraic numbers form a field, and algebraic integers form a ring. This is not obvious from the definition, but we will take it as true for now. Let us give some examples.

- Any $\alpha \in \mathbb{Q}$ is an algebraic number, since we may pick $p(x) = x - \alpha = 0$. Similarly, any $n \in \mathbb{Z}$ is an algebraic integer per the exact same construction.
- $\mathbb{Z}^{\text{alg}} \subset \mathbb{Q}^{\text{alg}}$.
- $\sqrt{-5} \in \mathbb{Z}^{\text{alg}}$, since $(\sqrt{-5})^2 + 5 = 0$.
- $\omega = \frac{1}{2}(-1 + \sqrt{-3})$ an algebraic integer, since $\omega^2 + \omega + 1 = 0$.
- For $n \geq 1$, we have that $\zeta = \exp(2\pi i/n) \in \mathbb{Z}^{\text{alg}}$, since $\zeta^n - 1 = 0$. ζ is the *primitive n^{th} root of unity*.
- Let $\alpha \in \mathbb{Q}^{\text{alg}}$ and $r \in \mathbb{Q}$. Then $r\alpha \in \mathbb{Q}^{\text{alg}}$. We will see later that, for all $\alpha \in \mathbb{Q}^{\text{alg}}$, there exists some $r \in \mathbb{Z}$ such that $r\alpha \in \mathbb{Z}^{\text{alg}}$.
- $\mathbb{Q}^{\text{alg}}$ is countable, since it is the union of the (finitely many) roots of polynomials in $\mathbb{Q}[x]$, of which are countable. $\mathbb{C}$ is not countable, however, so $\mathbb{Q}^{\text{alg}} \subsetneq \mathbb{C}$.
- $e, \pi \notin \mathbb{Q}^{\text{alg}}$.

Another way to characterize $\mathbb{Q}^{\text{alg}}$ and $\mathbb{Z}^{\text{alg}}$ is to consider subsets $V \subset \mathbb{C}$ such that V is a vector space (respectively, a finitely generated abelian group) over $\mathbb{Q}$. As an example, $V = \mathbb{Q}$ is trivially a vector space over $\mathbb{Q}$, and $V = \{r + s\sqrt{-5} \mid r, s \in \mathbb{Q}\}$ is one as well. In particular, it contains the finitely generated abelian group $\{r + s\sqrt{-5} \mid r, s \in \mathbb{Z}\}$.

Lemma 2.9. Let $\alpha \in \mathbb{C}$. Then

- $\alpha \in \mathbb{Q}^{\text{alg}}$ if and only if there exists a nonzero finite dimensional vector space $V \subset \mathbb{C}$ over $\mathbb{Q}$ such that $\alpha V \subset V$.
- $\alpha \in \mathbb{Z}^{\text{alg}}$ if and only if there exists a finitely generated abelian group $M \subset \mathbb{C}$ such that $\alpha M \subset M$.

Proof. We start by proving (i). In the forwards direction, suppose $\alpha \in \mathbb{Q}^{\text{alg}}$; then $\alpha^n + a_{n-1}\alpha^{n-1} + \dots + a_0 = 0$. Let

$$V = \{r_0 + r_1\alpha + \dots + r_{n-1}\alpha^{n-1} \mid r_0, \dots, r_{n-1} \in \mathbb{Q}\},$$

which we readily see as finite-dimensional as a vector space over $\mathbb{Q}$. It follows that $\alpha V \subset V$, since

$$\begin{aligned}\alpha(r_0 + r_1\alpha + \cdots + r_{n-1}\alpha^{n-1}) &= r_0\alpha + r_1\alpha^2 + \cdots + r_{n-1}\alpha^n \\ &= r_0\alpha + r_1\alpha^2 + \cdots + r_{n-1}(-a_{n-1}\alpha^{n-1} - \cdots - a_0) \in V.\end{aligned}$$

Conversely, suppose $V \subset \mathbb{C}$ is finite-dimensional with $\alpha V \subset V$. Let $v_1, \dots, v_d$ be a basis of V over $\mathbb{Q}$. For each i , we see that

$$\alpha v_i = \sum_{j=1}^d r_{ij} v_j, \quad r_{ij} \in \mathbb{Q};$$

i.e., $v \mapsto \alpha v$ is represented by a matrix $R := (r_{ij})$ in this basis. Specifically, R is a $d \times d$ matrix with entries in $\mathbb{Q}$, which we may think of as a complex matrix:

$$R \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^d r_{1j} v_j \\ \vdots \\ \sum_{j=1}^d r_{dj} v_j \end{pmatrix} = \alpha \begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix}.$$

It follows that $\alpha \in \mathbb{C}$ is an eigenvalue of R , whence α is a root of the characteristic polynomial $\det(R - Ix) \in \mathbb{Q}[x]$, so it has to be an algebraic number.

The proof for (ii), i.e., the algebraic integers case, follows analogously. Note that we need to know that the finitely generated abelian group is free; per the classification theorem (of finitely generated abelian groups), every such V can be written in terms of its free part and torsion; but there is no torsion, so we may reiterate our earlier argument as desired. $\square$

Proposition 2.10. $\mathbb{Q}^{\text{alg}} \subset \mathbb{C}$ is a subfield, and $\mathbb{Z}^{\text{alg}} \subset \mathbb{C}$ is a subring.

Proof. Let $\alpha, \beta \in \mathbb{Q}^{\text{alg}}$. Then, per 2.9, there exist nonzero finite-dimensional subspaces $V, W \subset \mathbb{C}$ such that $\alpha V \subset V$ and $\beta W \subset W$. Consider

$$V \cdot W = \{v_1 w_1 + \dots + v_t w_t \mid v_j \in V, w_j \in W\};$$

observing that $v_1, \dots, v_n$ span V and $w_1, \dots, w_m$ span W , we simply have that $v_i w_j$ for $1 \leq i \leq n$ and $1 \leq j \leq m$ span $V \cdot W$. Thus, $V \cdot W$ is finite-dimensional and nonempty.

Now, for any $v \in V$ and $w \in W$, we have that $\alpha \cdot (vw) = (\alpha v) \cdot w \in V \cdot W$, and $\beta \cdot (vw) = v \cdot (\beta w) \in V \cdot W$. By linearity for $z \in V \cdot W$, we have $\alpha z \in V \cdot W$ and $\beta z \in V \cdot W$, whence $(\alpha + \beta)(V \cdot W) \subset V \cdot W$ and $(\alpha \cdot \beta)(V \cdot W) \subset V \cdot W$, so $\alpha + \beta$ and $\alpha\beta$ are both algebraic numbers per 2.9. An analogous argument holds for $\alpha, \beta \in \mathbb{Z}^{\text{alg}}$.

It remains to show that nonzero $\alpha \in \mathbb{Q}^{\text{alg}}$ have inverses, i.e., $\alpha^{-1} \in \mathbb{Q}^{\text{alg}}$. Indeed, the “multiplying by α ” map, $\alpha: V \rightarrow V$, is injective, so it is isomorphic, and so there exists α^{-1} as the inverse map to α . In particular, it preserves V . Alternatively, we may write

$$\alpha^n + a_{n-1}\alpha^{n-1} + \cdots + a_0 = 0 \implies \alpha \cdot \left(-\frac{1}{\alpha_0}\right) (\alpha^{n-1} + a_{n-1}\alpha^{n-2} + \cdots + a_1) = 1,$$

which is a more explicit construction for α^{-1} . $\square$

We now look at more examples.

- Every element of $\mathbb{Z}[\sqrt{-5}]$ is an algebraic integer. Indeed, every $\alpha \in \mathbb{Z}[\sqrt{-5}]$ is of the form $a + b\sqrt{-5}$ with $a, b \in \mathbb{Z}$, and there is clearly an easy monic quadratic $p(x) \in \mathbb{Z}[x]$ such that $p(\alpha) = 0$.
- $\alpha = \sqrt{2} + \sqrt{3} \in \mathbb{Z}^{\text{alg}}$, since it satisfies $\alpha^4 - 10\alpha^2 + 1 = 0$.

Proposition 2.11. Let $\alpha \in \mathbb{Q}^{\text{alg}}$. Then there exists a unique monic irreducible polynomial $f(x) \in \mathbb{Q}[x]$ such that $f(\alpha) = 0$. Moreover, if $g(x) \in \mathbb{Q}[x]$ also satisfies $g(\alpha) = 0$, then $f \mid g$.

Proof. Consider $I = \{h(x) \in \mathbb{Q}[x] \mid h(\alpha) = 0\} \neq \emptyset$ (nonempty, as α is algebraic). I is clearly an ideal, and $\mathbb{Q}[x]$ is a PID, so we may write $I = \langle f \rangle$ for some f . Since f is uniquely determined up to units, we may divide any choice of f by its leading coefficient to let f be monic. It remains to check that f is irreducible. Let $f = gh$; then $0 = f(\alpha) = g(\alpha)h(\alpha)$. It must be that one of $g(\alpha)$ or $h(\alpha)$ is zero, with the other being in I . Without loss of generality, let $g(x) \in I$; but then $f \mid g$, so $\deg g \geq \deg f$, whence h is a unit. $\square$

Day 3: Minimal Polynomials (Sep. 15, 2026)

Office hours will be held from 3–4pm on Wednesdays in BA6110. Last time, we defined the algebraic numbers, $\mathbb{Q}^{\text{alg}}$, and the algebraic integers, $\mathbb{Z}^{\text{alg}}$. We also saw that $\mathbb{Q}^{\text{alg}}$ is a field, and $\mathbb{Z}^{\text{alg}}$ is a ring.

Proposition 3.1. Let $\alpha \in \mathbb{Q}^{\text{alg}}$ be nonzero; there is a unique monic irreducible polynomial $f(x) \in \mathbb{Q}[x]$ such that $f(\alpha) = 0$. Moreover, if $g(x) \in \mathbb{Q}[x]$ such that $g(\alpha) = 0$, then $f \mid g$.

Definition 3.2. Let $\alpha \in \mathbb{Q}^{\text{alg}}$. Then the unique monic irreducible $f(x) \in \mathbb{Q}[x]$ with $f(\alpha) = 0$ is called the *minimal polynomial* of α . The *degree* of α is the degree of $f(x)$ itself.

If $\alpha \in \mathbb{Q}^{\text{alg}}$, write $\mathbb{Q}(\alpha) = \{h(\alpha) \mid h \in \mathbb{Q}[x]\}$. If $\alpha \in \mathbb{Z}^{\text{alg}}$, write $\mathbb{Z}[\alpha] = \{h(\alpha) \mid h \in \mathbb{Z}[x]\}$. Clearly, $\mathbb{Q}(\alpha) \subset \mathbb{C}$, and we call it the subring generated by $\mathbb{Q}$ and α in $\mathbb{C}$. Similarly, $\mathbb{Z}(\alpha)$ is the smallest subring of $\mathbb{C}$ containing α .

As an example, take $\mathbb{Q}(\sqrt{-5}) = \{a + b\sqrt{-5} \mid a, b \in \mathbb{Q}\}$, and $\mathbb{Z}[\sqrt{-5}] = \{a + b\sqrt{-5} \mid a, b \in \mathbb{Z}\}$.

Proposition 3.3. Let $\alpha \in \mathbb{Q}^{\text{alg}}$.

- (i) $\mathbb{Q}(\alpha) = \mathbb{Q}[x]/p(x)$, where p is the minimal polynomial of α .
- (ii) As a vector space over $\mathbb{Q}$, $\mathbb{Q}(\alpha)$ has basis $1, \alpha, \dots, \alpha^{n-1}$, where $n = \deg(\alpha)$.
- (iii) $\mathbb{Q}(\alpha)$ is a field.

Proof. Consider the map $\text{ev}_\alpha: \mathbb{Q}[x] \rightarrow \mathbb{Q}(\alpha)$, given by $h \mapsto h(\alpha)$. ev_α is a surjective ring homomorphism by definition of $\mathbb{Q}(\alpha)$ (cf. earlier), whence

$$\mathbb{Q}(\alpha) \cong \frac{\mathbb{Q}[x]}{\ker(\text{ev}_\alpha)}.$$

In particular, we saw that $\ker(\text{ev}_\alpha) = \{h \in \mathbb{Q}[x] \mid h(\alpha) = 0\} = \langle f(x) \rangle$, where $f(x) = a_0 + \dots + a_{n-1}x^{n-1} + x^n$, so we are done with (i). (ii) immediately follows from earlier. For (iii), pick $\beta \in \mathbb{Q}(\alpha)$ nonzero, and let $\beta = h(\alpha)$ with $h \in \mathbb{Q}[x]$, for which $f \nmid h$, since $h(\alpha) \neq 0$. By the division algorithm (or Bezout's lemma), there must exist polynomials $A, B \in \mathbb{Q}[x]$ such that $Af + Bh = 1$. Evaluating at α , we see that $A(\alpha)f(\alpha) = 0$, so $B(\alpha)h(\alpha) = 1$, so we have found an inverse $\beta^{-1} = B(\alpha) \in \mathbb{Q}(\alpha)$ as desired. $\square$

Example 3.4. We give some examples of irreducible polynomials.

- (i) The minimal polynomial of $\sqrt{-5}$ is $x^2 + 5$.³
- (ii) Let $\omega = \frac{1}{2}(-1 + \sqrt{-3}) = \exp(2\pi i/3)$. Clearly, $\omega^3 = 1$. Since $x^3 - 1$ is not irreducible, we see that we may factor and obtain the actual minimal polynomial for ω to be $x^2 + x + 1$.

Is $\frac{1}{2} \in \mathbb{Z}^{\text{alg}}$? Recall Gauss' lemma; suppose R is a UFD. A polynomial $f(x) \in R[x]$ is *primitive* if the GCD of its coefficients is 1; Gauss' lemma states that if $f, g \in R[x]$ is primitive, then fg is primitive as well. As a quick proof, suppose not; then there exists $p \in R$ prime such that $p \mid fg$; since R/p is an integral domain, let $\bar{f}, \bar{g} \in (R/p)[x]$ be the corresponding reductions. We have that $\bar{f} \cdot \bar{g} = 0$, but then it must be that $\bar{f} = 0$ or $\bar{g} = 0$, which is a contradiction.

³so... any idea why this is irreducible? "the eisenstein criterion?" *grimaces* wayyyy overkill...

Lemma 3.5. Let R be a UFD, with F its field of fractions. Suppose $g, h \in F[x]$ are monic such that $f := gh \in R[x]$. Then $g, h \in R[x]$.

Proof. Let c, d be the LCM of the denominators of the coefficients of g, h . Then cg and dh are primitive polynomials in $R[x]$. Gauss' lemma would then imply that $cg \cdot dh = (cd) \cdot gh$ is a primitive polynomial; but $gh \in R[x]$, so $cd \mid (cd) \cdot gh$ in $R[x]$, i.e., cd must be a unit in R . It follows that c and d are both units in R , whence $g, h \in R[x]$. $\square$

Corollary 3.6. If $\alpha \in \mathbb{Z}^{\text{alg}}$, then its minimal polynomial lies in $\mathbb{Z}[x]$.

Proof. Let $p(x) \in \mathbb{Q}[x]$ be the minimal polynomial of α . Since $\alpha \in \mathbb{Z}^{\text{alg}}$, there exists a monic $h(x) \in \mathbb{Z}[x]$ such that $h(\alpha) = 0$. $p(x) \mid h(x)$, i.e., $h(x) = p(x)q(x)$, for some q with rational coefficients. From here, the lemma tells us that $p, q \in \mathbb{Z}[x]$. $\square$

Corollary 3.7. $\mathbb{Q} \cap \mathbb{Z}^{\text{alg}} = \mathbb{Z}$.

Proof. Let $\alpha \in \mathbb{Q} \cap \mathbb{Z}^{\text{alg}}$. The minimal polynomial of α is $x - \alpha$, so $\alpha \in \mathbb{Z}$. $\square$

Definition 3.8. Algebraic numbers $\alpha, \beta \in \mathbb{Q}^{\text{alg}}$ are *conjugate* if they have the same minimal polynomial.

Lemma 3.9. We have some more properties of minimal polynomials.

- (i) If $f \in \mathbb{Q}[x]$ is monic and irreducible of degree d , then f has d distinct roots in $\mathbb{C}$.
- (ii) If α is an algebraic number of degree d , then there exist exactly d algebraic numbers conjugate to α .

Clearly, (i) implies (ii). We will prove this next lecture.

Day 4: Number Fields (Sep. 17, 2026)

Office hours have been moved to 2–3pm on Thursdays. Last time, given an algebraic number $\alpha \in \mathbb{Q}^{\text{alg}}$, we defined the conjugates of α to be the algebraic numbers with the same minimal polynomial, and we introduced the following lemma,

Lemma 4.1. (i) Let $f \in \mathbb{Q}[x]$ be monic and irreducible of degree d ; then f has d distinct roots in $\mathbb{C}$. (ii) If α is an algebraic number of degree d , then there exist exactly d algebraic numbers conjugate to α .

Proof. It is clear that (i) implies (ii), so it suffices to prove (i). First, $\mathbb{C}$ is algebraically closed, so $f = (x - \alpha_1) \dots (x - \alpha_d)$ with each $\alpha_i \in \mathbb{C}$. We need to show that $\alpha_i \neq \alpha_j$ for all i, j ; suppose $\alpha \in \mathbb{C}$ is a root of f with multiplicity greater than 1. Then f being monic and irreducible implies that f is the minimal polynomial of α . By taking the derivative of f , we see that $f'(\alpha) = 0$; but then $\deg f' = d - 1$, which is a contradiction. $\square$

Let's do some examples.

- (i) The conjugates of $\sqrt{5}$ are $\sqrt{5}$ and $-\sqrt{5}$, since $x^2 - 5 = 0$ is the minimal polynomial of $\sqrt{5}$. Recall that it is easy to check if a quadratic polynomial is irreducible (check the rational roots).
- (ii) Let $f(x) = x^4 - 10x^2 + 1$, which we know (from previous classes) may be factored as

$$f(x) = (x - \sqrt{2} - \sqrt{3})(x - \sqrt{2} + \sqrt{3})(x + \sqrt{2} - \sqrt{3})(x + \sqrt{2} + \sqrt{3}).$$

We claim that f is irreducible; there are many ways to check this, see the MAT347 notes. [In my opinion, the easiest would be to check the extensions of \$\mathbb{Q}\$.](#)

Corollary 4.2 (Viète). If $\alpha \in \mathbb{Q}^{\text{alg}}$ has minimal polynomial $f(x) = x^d + a_{d-1}x^{d-1} + \dots + a_0$ with conjugates $\alpha_1, \dots, \alpha_d$, then

$$\alpha_1 \dots \alpha_d = (-1)^d a_0, \quad \alpha_1 + \dots + \alpha_d = -a_{d-1}.$$

The proof is immediate; just write $f(x) = (x - \alpha_1) \dots (x - \alpha_d)$.

Definition 4.3. An *algebraic number field* is a subfield of F of $\mathbb{C}$ which is finite-dimensional over $\mathbb{Q}$.

Equivalently, an algebraic number field is a finite extension of $\mathbb{Q}$. (recall from MAT347 that any finite extension is generated by a single element, i.e., the primitive element theorem. cf. D&F §14.4.25). If F is a number field, then the *ring of integers* of F is denoted $\mathcal{O}_F = F \cap \mathbb{Z}^{\text{alg}}$.

Remark 4.4. We have the following properties of number fields.

- (i) Let $V = F \subset \mathbb{C}$ be a number field; then $\alpha \cdot V \subset V$ for $\alpha \in F$, and so $\alpha \in \mathbb{Q}^{\text{alg}}$. Specifically, any element of an algebraic number field is an algebraic number, $F \subset \mathbb{Q}^{\text{alg}}$.
- (ii) $\mathcal{O}_F$ is a subring of $\mathbb{C}$ containing $\mathbb{Z}$.
- (iii) $\mathbb{Q} \cap \mathbb{Z}^{\text{alg}} = \mathbb{Z}$, i.e., $\mathcal{O}_{\mathbb{Q}} = \mathbb{Z}$.
- (iv) Let $\alpha \in \mathbb{Q}^{\text{alg}}$; we have that $\mathbb{Q}(\alpha) \subset \mathbb{C}$ is an algebraic number field. The primitive element theorem states that *every* number field is of this form.

Example 4.5. Let $d \in \mathbb{Z} \setminus \{0, 1\}$ be squarefree. Let $F = \mathbb{Q}(\sqrt{d}) = \{a + b\sqrt{d} \mid a, b \in \mathbb{Q}\}$. What is $\mathcal{O}_F$? Observe that

$$(a + b\sqrt{d})(a - b\sqrt{d}) = a^2 - db^2, \quad (a + b\sqrt{d}) + (a - b\sqrt{d}) = 2a,$$

so the minimal polynomial is $x^2 - 2ax + (a^2 - db^2)$. We want that $2a \in \mathbb{Z}$ and $a^2 - db^2 \in \mathbb{Z}$. If $d \equiv 2, 3 \pmod{4}$, then a, b are actually integers (checked by bashing). If $d \equiv 1 \pmod{4}$, then a, b are both integers, or half-integers. Thus,

$$\mathcal{O}_F = \begin{cases} \mathbb{Z} + \mathbb{Z}\sqrt{d}, & d \equiv 2, 3 \pmod{4}, \\ \mathbb{Z} + \mathbb{Z}\left(\frac{-1+\sqrt{d}}{2}\right), & d \equiv 1 \pmod{4}. \end{cases}$$

Note that, for $\alpha \in \mathbb{Z}^{\text{alg}}$ and $F = \mathbb{Q}(\alpha)$, $\mathcal{O}_F$ can be bigger than $\mathbb{Z}[\alpha]$.

Lemma 4.6. (i) If $\alpha \in \mathbb{Q}^{\text{alg}}$, there exists a nonzero integer $m \in \mathbb{N}$ such that $m\alpha \in \mathbb{Z}^{\text{alg}}$.

(ii) If F is a number field, then $F = \text{Frac}(\mathcal{O}_F)$.

Proof. We start by proving that (i) implies (ii). Since $\mathbb{Z} \subset \mathcal{O}_F$, we have that $\alpha = \frac{m\alpha}{m} \in \text{Frac}(\mathcal{O}_F)$. *Note that he skips a lot of intuition here, but the point is that $\mathcal{O}_F$ contains the integers, so from (i), we may write α as a fraction in terms of elements of $\mathcal{O}_F$; since $\mathcal{O}_F = F \cap \mathbb{Z}^{\text{alg}}$, it is enough to have $\alpha \in \mathcal{O}_F$ to get (ii).*

To prove (i), let $f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0$ (with $a_i \in \mathbb{Q}$) be the minimal polynomial of α . Then there exists m such that $ma_i \in \mathbb{Z}$ for all i ; it follows that we may multiply $f(x)$ throughout by m^n to see that $m\alpha$ is a root of

$$x^n + ma_{n-1}x^{n-1} + m^2a_{n-2}x^{n-2} + \dots + m^na_0 \in \mathbb{Z}[x]. \quad \square$$

We do more examples now.

- (i) Consider the Gaussian integers $F = \mathbb{Q}(i)$, where $i = \sqrt{-1}$ and $-1 \equiv 3 \pmod{4}$. Then $\mathcal{O}_F = \mathbb{Z}[i]$, for which $\mathcal{O}_F$ is a Euclidean domain with Euclidean norm $N(\alpha) = \alpha\bar{\alpha}$.
- (ii) Let $\omega = \exp(2\pi i/3) \in \mathbb{C}$. We have that $F = \mathbb{Q}(\omega) = \mathbb{Q}(\sqrt{-3})$. Since $-3 \equiv 1 \pmod{4}$, we have that $\mathcal{O}_F = \mathbb{Z} + \mathbb{Z}\omega = \mathbb{Z}[\omega] \neq \mathbb{Z}[\sqrt{-3}]$. We claim that this is a Euclidean domain, with $N(\alpha) = \alpha\bar{\alpha}$ the Euclidean norm.
- (iii) Let $F = \mathbb{Q}\sqrt{-5}$. $-5 \equiv 3 \pmod{4}$, so $\mathcal{O}_F = \mathbb{Z}[\sqrt{-5}]$. Consider the Euclidean norm $N(a + b\sqrt{-5}) = a^2 + 5b^2 \in \mathbb{Z}_{\geq 0}$.

Lemma 4.7. The only units in $\mathbb{Z}[\sqrt{-5}]$ are -1 and 1 . The elements $3, 7, 1 + 2\sqrt{-5}, 1 - 2\sqrt{-5}$ are all irreducible.

Proof. We claim that $\alpha \in \mathbb{Z}[\sqrt{-5}]$ is a unit if and only if $N(\alpha) = 1$. Suppose $\alpha\beta = 1$; then $N(\alpha\beta) = N(\alpha)N(\beta) = 1$. Since $N(\alpha) \in \mathbb{Z}_{\geq 0}$, if $N(\alpha) = 1$, then $\alpha\bar{\alpha} = 1$, meaning $\bar{\alpha} = \alpha^{-1}$. In this manner, it is enough to solve $a^2 + 5b^2 = 1$, from which we get $b = 0$ and $a = \pm 1$, whence the units are ± 1 .

We now check that our claimed elements are irreducible. Suppose 3 is not irreducible. Let us write $3 = \alpha\beta$, where α, β are non-units. Per above, we see that this must mean that their norms are not 1 ; so

$$9 = N(3) = N(\alpha)N(\beta) \implies N(\alpha) = N(\beta) = 3;$$

since $\alpha = a + b\sqrt{-5}$ with $N(\alpha) = a^2 + 5b^2 \neq 3$, we get a contradiction, and we are done. The process is analogous to check that 7 is irreducible, i.e., we arrive at a contradiction when we find that $a^2 + 5b^2 = 7$ has no integer solutions.

To check that $1 \pm 2\sqrt{-5}$ is irreducible (we'll only check the $1 + 2\sqrt{-5}$ case because the other follows similarly), suppose $1 + 2\sqrt{-5} = \alpha\beta$, where α, β are non-units. Since

$$21 = N(1 + 2\sqrt{-5}) = N(\alpha)N(\beta),$$

we must have $N(\alpha)$ is 3 or 7, which is not possible as we saw earlier. Thus, we are done. $\square$

The moral of this situation is that, to analyze Diophantine equations, we really just want to bash things out with factorization. In addition, norms are really useful for testing these properties as well!

We recall some important properties of field extensions (perhaps, from MAT347). Let L/F be a finite extension (i.e., $F \subset L$). We denote the degree of the extension as

$$\deg L/F = |L : F| := \dim_F L.$$

As an example, $|\mathbb{Q}(\alpha) : \mathbb{Q}| = \deg \alpha$. The tower law states that, if L/M and M/F are finite extensions, and $\omega_1, \dots, \omega_n$ is a basis for L/M , then

$$L = M\omega_1 \oplus \dots \oplus M\omega_n.$$

We also have

$$|L : F| = |L : M| |M : F|.$$

We have the following key example. Let F be a number field, and take $\alpha \in F$; then

$$|F : \mathbb{Q}| = |F : \mathbb{Q}(\alpha)| |\mathbb{Q}(\alpha) : \mathbb{Q}|,$$

for which the latter is simply just $\deg \alpha$. In particular, $\deg \alpha \mid |F : \mathbb{Q}|$. What can we do with this? Suppose $\alpha = \sqrt{2} + \sqrt{3}$. $\mathbb{Q}(\alpha)$ contains $\sqrt{2}$ and $\sqrt{3}$. Since

$$\sqrt{2} = \frac{\alpha^3 - 9\alpha}{-4},$$

we see that $\mathbb{Q}(\sqrt{2}) \subset \mathbb{Q}(\alpha)$. By writing

$$\deg \alpha = |F : \mathbb{Q}| = |F : \mathbb{Q}(\sqrt{2})| |\mathbb{Q}(\sqrt{2}) : \mathbb{Q}| = 4,$$

where $|\mathbb{Q}(\sqrt{2}) : \mathbb{Q}| = 2$, we see that $|F : \mathbb{Q}(\sqrt{2})| = 2$.

We now discuss norm and trace.

Definition 4.8. Let L/F be a finite extension. The *norm* of α , denoted $N_{L/F}(\alpha)$, is the determinant of the F -linear map $\phi_\alpha = \bullet\alpha : L \rightarrow L$. Similarly, the trace of α is $\text{Tr}_{F/L}(\alpha) := \text{tr}(\phi_\alpha)$.

Let us look at some examples.

- (i) $\mathbb{C}/\mathbb{R}$ has basis $\{1, i\}$. The matrix for multiplication by $x + iy$ is

$$\begin{pmatrix} x & -y \\ y & x \end{pmatrix},$$

(which you may perhaps recognize as the Cauchy–Riemann equations). Then $N_{\mathbb{C}/\mathbb{R}}(x + iy) = x^2 + y^2$, and $\text{Tr}_{\mathbb{C}/\mathbb{R}}(x + iy) = 2x$.

- (ii) $\mathbb{Q}(i)/\mathbb{Q}$ follows the exact same process. The norm and trace of $a + bi$ is $a^2 + b^2$, $2a$ respectively, since the basis and matrix are the same.
- (iii) $\mathbb{Q}(\omega)/\mathbb{Q}$, where $\omega^2 + \omega + 1 = 0$ (and $\omega = \exp(2\pi i/3)$), has basis $1, \omega$. Multiplication by $a + b\omega$ in this basis is

$$\begin{pmatrix} a & -b \\ b & a - b \end{pmatrix},$$

so $N_{\mathbb{Q}(\omega)/\mathbb{Q}}(a + b\omega) = a^2 + b^2 - ab$, and $\text{Tr}_{\mathbb{Q}(\omega)/\mathbb{Q}}(a + b\omega) = 2a - b$. The norm is the product of the conjugates, and conjugate in this situation is the complex conjugate (really, this has to do with the fact that the degree is 2; not every degree 2 extension is like this, though).

Proposition 4.9. We give the main properties of the norm here. $N_{L/F}: L \rightarrow F$ has the following properties,

- (i) $N_{L/F}(\alpha_1 \alpha_2) = N_{L/F}(\alpha_1) N_{L/F}(\alpha_2)$;
- (ii) $N_{L/F}(a) = a^{|L:F|}$ if $a \in F$;
- (iii) $N_{L/F}(\alpha) = 0$ if and only if $\alpha = 0$;
- (iv) If M is intermediate between L and F , then

$$N_{L/F} = N_{M/F} \circ N_{L/M}.$$

Proof. Properties (i) to (iii) are clear, so we will check (iv) only. We will start with a proof sketch in the special case that $\alpha \in M$; let $\omega_1, \dots, \omega_n$ be a basis for L/M . Then $L = M\omega_1 + \dots + M\omega_n$, and α acts block-diagonally. Thus, $N_{L/F}(\alpha) = N_{M/F}(\alpha)^n = N_{M/F} \circ N_{L/M}(\alpha)$. *Perhaps, this will be continued later?* $\square$

Proposition 4.10. $\text{Tr}_{L/F}$ has the following properties,

- (i) $\text{Tr}_{L/F}(\alpha_1 + \alpha_2) = \text{Tr}_{L/F}(\alpha_1) + \text{Tr}_{L/F}(\alpha_2)$;
- (ii) $\text{Tr}_{L/F}(a\alpha) = a \text{Tr}_{L/F}(\alpha)$ for $a \in F$ and $\alpha \in L$;
- (iii) $\text{Tr}_{L/F}(a) = |L:F|a$ if $a \in F$;
- (iv) If M is an intermediate field between L and F , then

$$\text{Tr}_{L/F} = \text{Tr}_{M/F} \circ \text{Tr}_{L/M}.$$

Proof. The proof is the same. $\square$

Remark 4.11. Let L/F be Galois. Then

$$N_{L/F}(\alpha) = \prod_{\sigma \in \text{Gal}(L/F)} \sigma(\alpha), \quad \text{Tr}_{L/F}(\alpha) = \sum_{\sigma \in \text{Gal}(L/F)} \sigma(\alpha).$$

In general, sum and products embed L into F . *I didn't catch what he said here; he'll probably restate this next class, so, we'll see then.*

We will do this next time.

Day 5: Norm and Trace (Sep. 22, 2026)

Recall, from last time, that we defined norm and trace for field extensions. Specifically, let K/F be a finite field extension; then, for $\alpha \in L$, the left-multiplication map $\phi_\alpha: L \rightarrow L$ given by $\beta \mapsto \alpha\beta$ is a K -linear map of K -vector spaces, for which

$$N_{L/F}(\alpha) = \det(\phi_\alpha), \quad \text{Tr}_{L/F}(\alpha) = \text{tr}(\phi_\alpha).$$

Note that this definition is different than that of the book (cf. p.15), so we will relate our definition to theirs now.

Proposition 5.1. Let $\bar{F}$ be an algebraic closure of F . Then, for $\alpha \in L$, we have that

$$\text{Tr}_{L/F}(\alpha) = \sum_{\substack{\sigma: L \rightarrow \bar{F} \\ F\text{-embeddings}}} \sigma(\alpha), \quad N_{L/F}(\alpha) = \prod_{\substack{\sigma: L \rightarrow \bar{F} \\ F\text{-embeddings}}} \sigma(\alpha).$$

Proof. Observe that $F \subset F(\alpha) \subset L$. Then $\text{Tr}_{L/F}(\alpha) = |L : F(\alpha)| \text{Tr}_{F(\alpha)/F}(\alpha)$, where $F(\alpha) = F[x]/p(x)$, with $p(x)$ being the minimal polynomial of α . $p(x)$ has $n = \deg p = |F(\alpha) : F|$ distinct roots, which we will call $\alpha_1, \dots, \alpha_n \in \bar{F}$, so there are exactly n embeddings of $F(\alpha) \hookrightarrow \bar{F}$. $\square$

In particular,

$$\text{Tr}_{F(\alpha)/F}(\alpha) = \sum_{i=1}^n \alpha_i, \quad N_{F(\alpha)/F}(\alpha) = \prod_{i=1}^n \alpha_i.$$

In the basis of $F(\alpha)$, i.e., $\{1, \alpha, \dots, \alpha^{n-1}\}$, the matrix of ϕ_α is given by

$$\begin{pmatrix} 0 & & & -a_0 \\ 1 & & & -a_1 \\ & \ddots & & \vdots \\ 0 & & 1 & -a_{n-1} \end{pmatrix},$$

i.e., the companion matrix for the minimal polynomial $p(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0$ of α .

In L/F , there are $|L : F|$ F -algebra embeddings $L \rightarrow \bar{F}$, where each one takes α to one of $\alpha_1, \dots, \alpha_n$.

Claim 5.2. Each α_i appears exactly $\frac{1}{n} |L : F|$ times.

Proof. For every embedding $F(\alpha) \rightarrow \bar{F}$, there exists $\frac{1}{n} |L : F|$ $F(\alpha)$ -algebra embeddings $L \rightarrow \bar{F}$. Specifically,

$$|L : F(\alpha)| = \frac{|L : \bar{F}|}{|F(\alpha) : F|} = \frac{|L : F|}{n}. \quad \square$$

In this way, we may write

$$\sum_{\substack{\sigma: L \rightarrow \bar{F} \\ F\text{-alg}}} \sigma(\alpha) = n \sum_{\substack{\sigma: L \rightarrow \bar{F} \\ F\text{-alg}}} \sigma(\alpha) = n \text{Tr}_{F(\alpha)/F}(\alpha) = \text{Tr}_{L/F}(\alpha).$$

The same applies to norm. As a corollary, if L/F is a finite Galois extension with Galois group G , then, for $\alpha \in L$,

$$\text{Tr}_{L/F}(\alpha) = \sum_{\sigma \in G} \sigma(\alpha), \quad N_{L/F}(\alpha) = \prod_{\sigma \in G} \sigma(\alpha).$$

Proposition 5.3. Let F be a number field. If $\alpha \in \mathcal{O}_F$, then $\text{Tr}_{F/\mathbb{Q}}(\alpha), N_{F/\mathbb{Q}}(\alpha) \in \mathbb{Z}$.

Proof. We saw, by Viète's formulas, that

$$\text{Tr}_{\mathbb{Q}(\alpha)/\mathbb{Q}}(\alpha) = -a_{n-1}, \quad N_{\mathbb{Q}(\alpha)/\mathbb{Q}}(\alpha) = (-1)^n a_0,$$

for which a_0, a_{n-1} are coefficients of the minimal polynomial of α , whence they are integers. It follows that

$$\text{Tr}_{F/\mathbb{Q}}(\alpha) = |F : \mathbb{Q}(\alpha)| \text{Tr}_{\mathbb{Q}(\alpha)/\mathbb{Q}}(\alpha), \quad N_{F/\mathbb{Q}}(\alpha) = |F : \mathbb{Q}(\alpha)| N_{\mathbb{Q}(\alpha)/\mathbb{Q}}(\alpha)$$

yields the desired result, as $|F : \mathbb{Q}(\alpha)|$ is an integer as well. $\square$

We now talk about another important invariant: the discriminant (cf. p.18).

Definition 5.4. Let L/F be a finite extension, and pick $\alpha_1, \dots, \alpha_n \in L$; then the associated discriminant is

$$\Delta(\alpha_1, \dots, \alpha_n) := \det(\text{Tr}_{L/F}(\alpha_i \alpha_j))_{ij} \in F.$$

As an example, let $d \in \mathbb{Z} \setminus \{0, 1\}$ not be a cube, and consider $F = \mathbb{Q}(d^{1/3})$; then the minimal polynomial is $x^3 - d = 0$, so the basis for $F/\mathbb{Q}$ is $\{1, d^{1/3}, d^{2/3}\}$. We have that $\Delta(1, d^{1/3}, d^{2/3})$ may be computed as the determinant

$$\det \begin{pmatrix} \text{Tr}_{F/\mathbb{Q}} 1 & \text{Tr}_{F/\mathbb{Q}} d^{1/3} & \text{Tr}_{F/\mathbb{Q}} d^{2/3} \\ \text{Tr}_{F/\mathbb{Q}} d^{1/3} & \text{Tr}_{F/\mathbb{Q}} d^{2/3} & \text{Tr}_{F/\mathbb{Q}} d \\ \text{Tr}_{F/\mathbb{Q}} d^{2/3} & \text{Tr}_{F/\mathbb{Q}} d & \text{Tr}_{F/\mathbb{Q}} d^{4/3} \end{pmatrix} = \det \begin{pmatrix} 3 & 0 & 0 \\ 0 & 0 & 3d \\ 0 & 3d & 0 \end{pmatrix} = -27d^2.$$

Note that, if F is a number field, and $\alpha_1, \dots, \alpha_n \in \mathcal{O}_f$, then $\Delta(\alpha_1, \dots, \alpha_n) \in \mathbb{Z}$ (the determinant of an integer-valued matrix is an integer).

Proposition 5.5. The discriminant detects bases. Let L/F be a finite extension, and let $\alpha_1, \dots, \alpha_n \in L$.

(i) If $\Delta(\alpha_1, \dots, \alpha_n) \neq 0$, then $\alpha_1, \dots, \alpha_n$ are linearly independent over F .

(ii) If $\alpha_1, \dots, \alpha_n$ form a basis for L/F and $\text{char } F = 0$, then $\Delta(\alpha_1, \dots, \alpha_n) \neq 0$.

Proof. For (i), suppose $\alpha_1, \dots, \alpha_n$ are linearly dependent. Then there exists $a_1, \dots, a_n \in F$ such that $a_1 \alpha_1 + \dots + a_n \alpha_n = 0$, so for any j ,

$$0 = \text{Tr}_{L/F} \left(\alpha_j \left(\sum_i a_i \alpha_i \right) \right) = \sum_{i=1}^n a_i \text{Tr}_{L/F}(\alpha_i \alpha_j) \implies (\text{Tr}_{L/F}(\alpha_i \alpha_j))_{ij} \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} = 0.$$

Thus, $\det(\text{Tr}_{L/F}(\alpha_i \alpha_j)) = 0$.

For (ii), suppose $\{\alpha_1, \dots, \alpha_n\}$ is a basis of L/F and $\Delta(\alpha_1, \dots, \alpha_n) = c$. Then the F -linear map $\phi: L \rightarrow L$ given by $(\text{Tr}_{L/F}(\alpha_i \alpha_j))$ has nonzero kernel (by reading the matrix as a linear transformation), and we can run the same argument backwards to see that there is some choice of $a_1, \dots, a_n \in F$, not all zero, such that

$$\sum_{i=1}^n a_i \text{Tr}_{L/F}(\alpha_i \alpha_j) = 0$$

for all j . Now, set $\alpha = a_1 \alpha_1 + \dots + a_n \alpha_n \in L$, and consider $\beta \in L$ given by $\beta = b_1 \alpha_1 + \dots + b_n \alpha_n$; we have that

$$\text{Tr}_{L/F}(\beta \alpha) = \text{Tr}_{L/F} \left(\sum_{j=1}^n b_j \alpha_j \alpha \right) = \sum_{j=1}^n b_j \text{Tr}_{L/F}(\alpha_j \alpha) = \sum_{j=1}^n b_j \left(\sum_{i=1}^n a_i \text{Tr}_{L/F}(\alpha_i \alpha_j) \right) = 0.$$

If $\alpha \neq 0$, we may take $\beta = \alpha^{-1}$ to get $\text{Tr}_{L/F}(1) = 0$, which is a contradiction, and we are done. ⁴ $\square$

⁴this is why you don't rush proofs man.

Day 6: Integral Basis (Sep. 24, 2026)

Rozenblyum isn't particularly happy with how he introduced the discriminant, so we will be doing some linear algebra exposition for the start of this lecture. Consider a vector space V over F . A *symmetric bilinear form* is a map

$$\langle -, - \rangle : V \times V \rightarrow F$$

such that $\langle v, w \rangle = \langle w, v \rangle$ and, for all $v \in V$, the map $V \rightarrow F$ given by $w \mapsto \langle v, w \rangle$ is a linear map. $\langle -, - \rangle$ is *nondegenerate* if, for all $v \in V$, there exists some $w \in V$ such that $\langle v, w \rangle \neq 0$. Our main example is the following: consider L/F a finite field extension; then

$$L \times L \rightarrow F, \quad (\alpha, \beta) \mapsto \text{Tr}(\alpha\beta)$$

is a symmetric bilinear form. Moreover, if $\text{char } F = 0$, it's nondegenerate, since $\text{Tr}(\alpha\alpha^{-1}) = \text{Tr}(1) = |L : F|$. Suppose V/F is a vector space with symmetric bilinear form $\langle -, - \rangle$. If $\{v_1, \dots, v_n\}$ is a basis of V , we can form a matrix

$$Q = (\langle v_i, v_j \rangle)_{i,j}.$$

Observe that if $v = \sum_i a_i v_i$ and $w = \sum_j b_j v_j$, then

$$\langle v, w \rangle = \left\langle \sum_i a_i v_i, \sum_j b_j v_j \right\rangle = \sum_{i,j} a_i b_j \langle v_i, v_j \rangle = (a_1, \dots, a_n) Q \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}.$$

We may also observe that $\det Q \neq 0$ if and only if $\langle -, - \rangle$ is nondegenerate. Now, suppose $\{w_1, \dots, w_n\}$ is another basis of V . We have a change of basis matrix $P = (P_{ij})$, where

$$w_j = \sum_{i=1}^n P_{ij} v_i,$$

and we have $Q' = (\langle w_i, w_j \rangle)$ (i.e., representing the same symmetric bilinear form, but in a different basis). How are Q and Q' related?

$$\langle w_i, w_j \rangle = \left\langle \sum_{k=1}^n P_{ki} v_k, \sum_{\ell=1}^n P_{\ell j} v_\ell \right\rangle = \sum_{k,\ell} P_{ki} P_{\ell j} \langle v_k, v_\ell \rangle = \sum_{k,\ell} P_{ki} P_{\ell j} Q_{k\ell} = (P^\top Q P)_{ij},$$

i.e., $Q' = P^\top Q P$.

Corollary 6.1. Let L/F be a finite field extension, and let $\{\alpha_1, \dots, \alpha_n\}$ be a basis of L/F . Let $\{\beta_1, \dots, \beta_n\}$ be a different basis of L/F . Let P be the transition matrix from α to β ; then

$$\beta_j = \sum_{i=1}^n P_{ij} \alpha_i,$$

and $\Delta(\beta_1, \dots, \beta_n) = (\det P)^2 \Delta(\alpha_1, \dots, \alpha_n)$.

Proof. Let Q be the matrix corresponding to the form $\text{Tr}(\alpha\beta)$ in the basis $\{\alpha_1, \dots, \alpha_n\}$, i.e., $Q_{ij} = \text{Tr}(\alpha_i \alpha_j)$. Then $\Delta(\alpha_1, \dots, \alpha_n) = \det Q$. Now, let Q' be the corresponding matrix to $\{\beta_1, \dots, \beta_n\}$; since $Q' = P^\top Q P$, we have that

$$\Delta(\beta_1, \dots, \beta_n) = \det Q' = \det P^\top \det Q \det P = (\det P)^2 \det Q = (\det P)^2 \Delta(\alpha_1, \dots, \alpha_n). \quad \square$$

Recall that, if $p(x)$ is a polynomial of degree n with roots $\alpha_1, \dots, \alpha_n$, then

$$\Delta(p) = \left(\prod_{i < j} (\alpha_j - \alpha_i) \right)^2$$

is a symmetric polynomial in the α_i . In particular, $\Delta(p)$ is a polynomial in $a_0, \dots, a_{n-1}$ (i.e., the coefficients of p), and is called the *discriminant* of p .

We will detour a little here to talk about elementary symmetric polynomials. Suppose $f(x_1, \dots, x_n) \in F[x_1, \dots, x_n]$. We say that f is symmetric if $f(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = f(x_1, \dots, x_n)$ for all $\sigma \in S_n$. Specifically, let us define $\sigma_i(x_1, \dots, x_n)$ to be the sums of products of x_i 's; i.e.,

$$\sigma_1 = x_1 + \dots + x_n, \quad \sigma_2 = \sum_{i < j} x_i x_j, \quad \dots \quad \sigma_n = x_1 \dots x_n.$$

These are called the “elementary symmetric polynomials”, for which the fundamental theorem of symmetric polynomials states that any symmetric $f(x_1, \dots, x_n)$ can be expressed as $g(\sigma_1, \dots, \sigma_n)$ for a unique g . This is why we're able to write $\Delta(p)$ as a polynomial in $a_0, \dots, a_{n-1}$.

Theorem 6.2. Consider a field extension $L = F(\alpha)/F$, where α has minimal polynomial $p(x)$ of degree n . Then

$$\Delta(1, \alpha, \dots, \alpha^{n-1}) = \Delta(p(x)).$$

Proof. Write $p(x) = (x - \alpha_1) \dots (x - \alpha_n)$. We see $\text{Tr}(\alpha) = \sum_{i=1}^n \alpha_i$ and $\text{Tr}(\alpha^k) = \sum_{i=1}^n \alpha_i^k$, so we may use this fact to try and write the discriminant in terms of these traces. Recall that the discriminant of $\{1, \alpha, \dots, \alpha^{n-1}\}$ is given by

$$\Delta(1, \alpha, \dots, \alpha^{n-1}) = \det \left(\text{Tr}_{L/F}(\alpha^{(i+1)+(j-1)}) \right)_{1 \leq i, j \leq n} = \det \left(\sum_{k=1}^n \alpha_k^{i+j-2} \right).$$

Denote the entries as $M_{i,j}$ for convenience. We wish to decompose M as $M = V^T V$, where

$$V = \begin{pmatrix} 1 & \alpha_1 & \alpha_1^2 & \dots & \alpha_1^{n-1} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & \alpha_n & \alpha_n^2 & \dots & \alpha_n^{n-1} \end{pmatrix};$$

note that V is called the *Vandermonde matrix*. Indeed, since $V_{ij} = \alpha_i^{j-1}$, we see that

$$(V^T V)_{ij} = \sum_{k=1}^n V_{ik}^T V_{kj} = \sum_{k=1}^n V_{ki} V_{kj} = \sum_{k=1}^n \alpha_k^{i-1} \alpha_k^{j-1} = M_{ij}$$

as desired. In this manner, it follows that 6.1 yields

$$\Delta(1, \alpha, \dots, \alpha^{n-1}) = (\det M)(\det V)^2,$$

where $\det V = \prod_{i < j} (\alpha_j - \alpha_i)$, so the right hand side is precisely the discriminant of the polynomial $p(x)$ as seen earlier. $\square$

As an observation,

$$\Delta(1, \alpha, \dots, \alpha^{n-1}) = N_{L/F}(p'(\alpha)).$$

Proof. Let $p(x) = (x - \alpha_1) \dots (x - \alpha_n)$, and denote the latter $n - 1$ terms as $g(x)$. We have that $p'(x)$ can be written $(x - \alpha_1)g'(x) + g(x)$, where

$$p'(\alpha_1) = g(\alpha_1) = \prod_{i>1} (\alpha_1 - \alpha_i).$$

We may let g take on $(x - \alpha_1) \dots \widehat{(x - \alpha_i)} \dots (x - \alpha_n)$ to also see that

$$p'(\alpha_i) = \prod_{j \neq i} (\alpha_i - \alpha_j),$$

so

$$N_{L/F}(p'(\alpha)) = \prod_{i=1}^n p'(\alpha_i) = \prod_{i \neq j} (\alpha_i - \alpha_j) = (-1)^{\frac{n(n-1)}{2}} \prod_{i < j} (\alpha_i - \alpha_j)^2,$$

where we picked up a term of $(-1)^{\#\{i < j\}}$ by counting. $\square$

Definition 6.3. Let F be a number field. Let $I \subset \mathcal{O}_F$ be a nonzero ideal. An *integral basis* of I is a basis of $F/\mathbb{Q}$, $\{\omega_1, \dots, \omega_n\} \in F$, such that

- (i) each $\omega_i \in I$,
- (ii) $I = \mathbb{Z}\omega_1 + \dots + \mathbb{Z}\omega_n$.

An integral basis of F is an integral basis of $I = \mathcal{O}_F$.

Lemma 6.4. Let F be a number field. If $I \subset \mathcal{O}_F$ is a nonzero ideal, then I contains a basis of F .

Proof. Let $\alpha_1, \dots, \alpha_n \in F$ be a basis; then there exists $m \in \mathbb{Z}$ such that $m\alpha_i \in \mathcal{O}_F$. Let $p \in I$ be a nonzero element; then the left-multiplication by β map, $\alpha \mapsto \beta\alpha$ from $F \rightarrow F$ is invertible. In particular, this means $\{\beta(m\alpha_i)\}_i$ is a basis of F . $\square$

Proposition 6.5. Let F be a number field and $I \subset \mathcal{O}_F$ a nonzero ideal. Then I has an integral basis.

Proof. Recall that, if $\omega_1, \dots, \omega_n \in \mathcal{O}_F$, then $\Delta(\omega_1, \dots, \omega_n) \in \mathbb{Z}$. Choose a basis of F , $\{\omega_1, \dots, \omega_n\} \in I$, such that $|\Delta(\omega_1, \dots, \omega_n)|$ is minimal (it can't be zero by linear independence). We claim that $\{\omega_1, \dots, \omega_n\}$ constitutes an integral basis.

Suppose not. Then there exists some element $\alpha \in I$ such that α is a rational linear combination of the basis elements, i.e., $\sum_i a_i \omega_i$ for $a_i \in \mathbb{Q}$ (but not all $a_i \in \mathbb{Z}$). Without loss of generality, suppose a_1 is not an integer. This means $a_1 = \lfloor a_1 \rfloor + \theta$ for some rational $\theta \in (0, 1)$. Consider

$$\omega'_1 := \alpha - \lfloor a_1 \rfloor \omega_1 = \theta \omega_1 + a_2 \omega_2 + \dots + a_n \omega_n,$$

which is clearly in I , as it is still a $\mathbb{Q}$ -linear combination. Clearly, $\{\omega'_1, \omega_2, \dots, \omega_n\}$ is a basis for F . We now check that this contradicts our supposition: indeed,

$$\Delta(\omega'_1, \omega_2, \dots, \omega_n) = \det(P)^2 \Delta(\omega_1, \dots, \omega_n), \quad P = \begin{pmatrix} \theta & & & 0 \\ a_2 & & & \\ \vdots & & \ddots & \\ a_n & & & 1 \end{pmatrix},$$

for which $\det P = \theta < 1$, whence $|\Delta(\omega'_1, \omega_2, \dots, \omega_n)| < |\Delta(\omega_1, \dots, \omega_n)|$, so we are done. $\square$

Proposition 6.6. Suppose $\{\omega_1, \dots, \omega_n\}$ and $\{\omega'_1, \dots, \omega'_n\}$ are integral bases of I . Then the absolute values of the discriminants of both bases are equal, i.e.,

$$|\Delta(\omega_1, \dots, \omega_n)| = |\Delta(\omega'_1, \dots, \omega'_n)|.$$

In particular, $\omega_1, \dots, \omega_n \in I$ is an integral basis if and only if it is a basis and the absolute value of the discriminant is minimal.

Proof. Without loss of generality, $|\Delta(\omega_1, \dots, \omega_n)|$ is minimal. We have that $\mathbb{Z}^n \rightarrow I$ is an isomorphism under the maps

$$(a_1, \dots, a_n) \mapsto \sum_i a_i \omega_i, \quad (b_1, \dots, b_n) \mapsto \sum_i b_i \omega'_i,$$

for which the change of basis matrix P gives an isomorphism $\mathbb{Z}^n \rightarrow \mathbb{Z}^n$. Any such change of basis must have $\det P = \pm 1$, so it follows that

$$\Delta(\omega'_1, \dots, \omega'_n) = (\det P)^2 \Delta(\omega_1, \dots, \omega_n) = \Delta(\omega_1, \dots, \omega_n). \quad \square$$

“An easy way to do exercise 5 on the homework is to use the above property.”

Definition 6.7. Let F be a number field, and let $\{\omega_1, \dots, \omega_n\} \in \mathcal{O}_F$ be an integral basis. The *absolute discriminant* of F is

$$\Delta_F = \Delta(\omega_1, \dots, \omega_n).$$

By the 6.6, this is well-defined. As an example, take d to be a squarefree integer, and consider $F = \mathbb{Q}(\sqrt{d})$. We have that

$$\Delta_F = \begin{cases} 4d & d \equiv 2, 3 \pmod{4}, \\ d & d \equiv 1 \pmod{4}, \end{cases}$$

since $\Delta(1, \sqrt{d}) = 4d$ for the first case, and d for the second case because $\{1, \sqrt{d}\}$ is not an integral basis for F (we may consider $\frac{1}{2}(1 + \sqrt{d})$), and so Δ_F for $d \equiv 1 \pmod{4}$ must be a square apart from $4d$, whence the only candidate is d (cf. p.143?).

This number has meaning... something called “ramification”.

Definition 6.8. Let F be a number field, and let $I \subset \mathcal{O}_F$ be a nonzero ideal. The *norm* of I , $N(I)$, is defined as the number of elements in the quotient $\mathcal{O}_F/I$.

In order for the definition to make sense, we need to know that $\#\{\mathcal{O}_F/I\}$ is finite.

Proposition 6.9. Let F be a number field and $I \subset \mathcal{O}_F$ a nonzero ideal. Then $\mathcal{O}_F/I$ is finite, i.e., a ring with finitely many elements.

Proof. Let A, B be finitely generated abelian groups such that $A \subset B$. Then $\text{rank } B/A = \text{rank } B - \text{rank } A$.

Alternatively, let $m \in I \cap \mathbb{Z}$ be nonzero (e.g., $\alpha \in I$ nonzero with minimal polynomial $x^n + a_{n-1}x^{n-1} + \dots + a_0$; then $a_0 \in I$). Let $\alpha \in \mathcal{O}_F/I$. We can write $\alpha = \sum_i c_i \overline{w_i}$, where $c_i \in \mathbb{Z}$, with $c_i = mq_i + r_i$. Then

$$\alpha = \sum_i r_i \overline{w_i} \pmod{I}.$$

Since $m \in I$, we see that any element of $\mathcal{O}_F/I$ can be represented as $\sum_i r_i \overline{w_i}$, for $r_i \in \mathbb{Z}$ and $|r_i| < m$. There are finitely such elements, so we are done. $\square$

Day 7: Dedekind Domains (Sep. 29, 2026)

Recall that $\mathbb{Z}[\sqrt{-5}] = \mathcal{O}_{\mathbb{Q}(\sqrt{-5})}$ is not a PID, since $3, 7, 1 + 2\sqrt{-5}, 1 - 2\sqrt{-5}$ are all irreducible elements, and

$$21 = 3 \cdot 7 = (1 + 2\sqrt{-5})(1 - 2\sqrt{-5}),$$

with ± 1 being units. In number rings, ideals generalize elements, and considering ideals in number rings restores unique factorization. We may think of these as “ideal numbers”.

Recall, that for a ring R , with $I, J \subset R$ ideals, we can consider $I + J = \{a + b \in R \mid a \in I, b \in J\}$ and $IJ = \langle ab \mid a \in I, b \in J \rangle$. If $I = \langle a_1, \dots, a_n \rangle$ and $J = \langle b_1, \dots, b_m \rangle$, then $IJ = \langle a_i b_j \rangle$ for $i = 1, \dots, n$ and $j = 1, \dots, m$. In $\mathbb{Z}[\sqrt{-5}]$,

$$\begin{aligned} \langle 3, 1 + 2\sqrt{-5} \rangle \langle 3, 1 - 2\sqrt{-5} \rangle &= \langle 3 \rangle, \\ \langle 7, 1 + 2\sqrt{-5} \rangle \langle 7, 1 - 2\sqrt{-5} \rangle &= \langle 7 \rangle, \\ \langle 3, 1 + 2\sqrt{-5} \rangle \langle 7, 1 + 2\sqrt{-5} \rangle &= \langle 1 + 2\sqrt{-5} \rangle, \\ \langle 7, 1 - 2\sqrt{-5} \rangle \langle 3, 1 - 2\sqrt{-5} \rangle &= \langle 1 - 2\sqrt{-5} \rangle. \end{aligned}$$

In the above, the ideals on the left hand side are prime ideals in R . From this, we may multiply the top two examples to get

$$\langle 21 \rangle = \langle 3, 1 + 2\sqrt{-5} \rangle \langle 3, 1 - 2\sqrt{-5} \rangle \langle 7, 1 + 2\sqrt{-5} \rangle \langle 7, 1 - 2\sqrt{-5} \rangle.$$

We want some general context in which the above is true.

Definition 7.1. A Dedekind domain is an integral domain R such that

- (i) every nonzero prime ideal of R is maximal,
- (ii) R is Noetherian, and
- (iii) R is integrally closed, i.e., $R \subset \text{Frac}(R)$. This means that, supposing $\alpha \in K$ such that $a_0, \dots, a_{n-1} \in R$ satisfy

$$\alpha^n + a_{n-1}\alpha^{n-1} + \dots + a_0 = 0,$$

i.e., α is a root of a monic polynomial $p(x) \in R[x]$, then $\alpha \in R$.

We now discuss some key facts of Dedekind domains.

Theorem 7.2. Every number ring $\mathcal{O}_F$ is a Dedekind domain.

Theorem. If R is a Dedekind domain and $I \subset R$ is a nonzero (proper) ideal, then there is a unique set of prime ideals, up to reordering, $p_1, \dots, p_n \subset R$ such that

$$I = p_1 p_2 \dots p_n.$$

As an example, if R is a PID, then R is a Dedekind domain.

Proof of 7.2. We proved that, if $I \subset \mathcal{O}_F$ is a nonzero ideal, then $\mathcal{O}_F/I$ is finite. This followed from the existence of integral bases. We now claim that if R is a finite integral domain, then it is a field.

Subproof. Take a nonzero element $x \in R$. Since R is an integral domain, the right-multiplication map $a \mapsto ax$ from $R \rightarrow R$ is injective automatically, and surjective since R is finite. Thus, there exists $y \in R$ such that $ay = 1$. ■

From here, if $I \subset \mathcal{O}_F$ is a nonzero prime ideal, then $\mathcal{O}_F/I$ is a finite integral domain, which is a field, which means I is maximal. *I prime implies that $\mathcal{O}_F/I$ is a finite PID.*

Suppose $I_1 \subset I_2 \subset \dots$ is an ascending chain of ideals. If $I_i = 0$ for all i , then the chain is stationary. Otherwise, without loss of generality, suppose I_1 is nonzero. Then

$$R/I_1 \twoheadrightarrow R/I_2 \twoheadrightarrow R/I_3 \twoheadrightarrow \dots,$$

which must eventually be stationary, since we start with some finite underlying set in R/I_1 , and the surjections $\twoheadrightarrow$ either decrease or keep the number of elements the same; in either case, there can only be so many maps decreasing the number of elements, meaning that the ascending chain must become stationary at some point.

To conclude that $\mathcal{O}_F$ is a Dedekind domain, we need to check that it is integrally closed. We will leave it as a homework exercise. To do this, you need to show that, for $\alpha \in \mathcal{O}_F$ such that $p(\alpha) = 0$, then $p \in \mathcal{O}_F[x]$ is monic, and so α is an algebraic integer. $\square$

To prove 8.7, we want to do the following first:

Theorem. Let $I \subset R$ be a nonzero ideal in a Dedekind domain. Then there exists a nonzero ideal J such that IJ is principal.

Lemma 7.3. Let R be a Dedekind domain and $I \subset R$ a nonzero ideal. Then I contains a product of nonzero prime ideals.

Proof. Suppose not. Let S be the set of ideals of R which don't contain a product of prime ideals. Since R is Noetherian, S contains a maximal element $M \in S$. Observe that M is not prime, since M does not contain a product of prime ideals (i.e., we would have $M \subset M$ otherwise). In this way, there exists $a, b \in M$ such that $a, b \notin M$, but $ab \in M$. Consider $I = M + (a)$ and $J = M + (b)$, for which $I, J \not\subset M$. We have that $IJ \subset M$, since every element of IJ is either in M or (ab) ; in particular, this means $I, J \neq R$. Since M is maximal with $I, J \not\subset S$, we see that I, J must contain products of prime ideals, i.e.,

$$I \supset p_1 \dots p_n, \quad J \supset q_1 \dots q_m,$$

and so $M \supset IJ \supset (p_1 \dots p_n)(q_1 \dots q_m)$. $\square$

We will prove 8.7 and 8.2 (these were relabeled because they were proved later) next lecture.

Day 8: Properties of Dedekind Domains, Ideal Class Group (Oct. 1, 2026)

Last time, we defined Dedekind domains. For now, an ideal will denote a *nonzero* ideal (just so Rozenblyum doesn't have to write nonzero every single time). Last lecture, we proved that every ideal in a Dedekind domain contains a product of prime ideals.

Lemma 8.1. R is a Dedekind domain with field of fractions K and $A \subsetneq R$ a proper ideal, then there exists some element $\gamma \in K \setminus R$ such that $\gamma A \subset R$.

Proof. Let $a \in A$ be a nonzero element; then $(a) \subset A$. By the earlier lemma,

$$A \supset (a) \supset p_1 \dots p_k,$$

with $p_1, \dots, p_k$ prime ideals (and k being chosen to be minimal). Every ideal is contained in a maximal ideal, so $A \subset P$, where P is a prime ideal (since maximal ideals are prime); so $p_1 \dots p_k \subset P$ (when you multiply ideals, you get a smaller ideal, which is why there is no contradiction).

We claim that $P = p_i$ for some i . Otherwise, for all i , there exists $a_i \in P_i \setminus P$; but $a_1 \dots a_k \in P$, so one of the a_i must be in P itself; from here, we would get that $P_i \subset P$, but then $P_i = P$, since prime ideals are maximal in Dedekind domains.

Without loss of generality, let $P = p_1$ per the above. This means we have

$$p_1 \dots p_i \subset (a) \subset A \subset P = p_1,$$

where, by minimality of k , $p_2 \dots p_k \not\subset (a)$; in other words, there exists some element $b \in (p_2 \dots p_k) \setminus (a)$. Take $\gamma = b/a \in K \setminus R$; we want to show that, for all $x \in A$, that $\gamma x \in R$. Equivalently, $bx \in (a)$. This is true for all $x \in p_1$, since $b \in p_2 \dots p_k$, so $bx \in p_1 \dots p_k \subset (a)$, so we are done. $\square$

Theorem 8.2. If R is a Dedekind domain and $I \subset R$ an ideal, then there exists an ideal J such that IJ is principal.

Proof. Let $\alpha \in I$ be nonzero; then $(\alpha) \subset I$. Define $J = \{x \in R \mid xI \subset (\alpha)\}$; then $J \neq \emptyset$ is an ideal with $\alpha \in J$. We want to show that $IJ = (\alpha)$. It suffices to check the opposite direction, since $IJ \subset (\alpha)$ by construction; in this manner, let

$$A = \frac{1}{\alpha} IJ \subset R.$$

We want to check that $A = R$. Supposing not, then A would be a proper ideal, and by 8.1, there would exist $\gamma \in A \setminus R$ such that $\gamma A \subset R$. We'll show that $\gamma \in R$ to get a contradiction. It is enough to check that γ satisfies a monic polynomial with coefficients in R , since Dedekind domains are integrally closed.

Since $\alpha \in I$, we have that $J \subset A = \frac{1}{\alpha} IJ$, for which $\gamma J \subset \gamma A \subset R$. We claim that $\gamma J \subset J$, which we would see by demonstrating $\gamma xy \in (\alpha)$ for $x \in J$ and $y \in I$. Indeed,

$$\frac{1}{\alpha} \gamma xy = \gamma \left(\frac{1}{\alpha} xy \right) \in R,$$

for which the bracketed portion is an element of A . We know that $\gamma A \subset R$ from earlier, so $\alpha^{-1} \gamma xy \in R$, which is precisely what we wanted to check.

In this way, $\gamma: J \rightarrow J$ is a map of abelian groups. Let $\alpha_1, \dots, \alpha_m$ be a generating set for the ideal J . We get

$$\gamma \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_m \end{pmatrix} = M \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_m \end{pmatrix}$$

for M an $m \times m$ matrix over R , i.e., $\gamma\alpha_i = \sum_{j=1}^m M_{ij}\alpha_j$. γ is an eigenvalue of M , so γ is a root of the characteristic polynomial of M , i.e., a monic polynomial with coefficients in R . $\square$

Define an equivalence relation on ideals $I \sim J$ if there exist $\alpha, \beta \in R$ such that $\alpha I = \beta J$. If $I \sim J \sim K$, then $\alpha I = \beta J$ and $\gamma J = \delta K$, so $\gamma\alpha I = \beta\gamma J = \beta\delta K$. Observe that $\sim$ is compatible with products, i.e., $I \sim I'$ and $J \sim J'$ yields $IJ \sim I'J'$. The upshot being, we may write $C\ell_R$ to be the “ideal classes” to denote the quotient set. This has a binary operation $[I] \cdot [J] = [IJ]$.

Claim 8.3. $C\ell_R$ is a group.

Proof. For any nonzero $\alpha \in R$, we have that $[\alpha I] = [(\alpha) \cdot I] = [I]$, so (α) is clearly an identity element in $C\ell_R$. Note that, for any nonzero $\alpha, \beta \in R$, $[(\alpha)] = [(\beta)]$, since $\beta(\alpha) = \alpha(\beta)$. This means that the identity element of $C\ell_R$ is given by equivalence class of principal ideals. Specifically, this means we may use 8.2 to find inverses. $\square$

We call $C\ell_R$ the *ideal class group* of R . Note that, if R is a PID, then $C\ell_R = \{e\}$. Conversely, if the class group is trivial, then R is a PID as well; specifically, it is enough to show that, if $I \sim (\alpha)$, then I is principal. $I \sim (\alpha)$ is equivalent to there existing nonzero $\alpha, \beta \in R$ such that $\alpha I = \beta(\alpha)$; but then

$$R \supset I = \frac{1}{\alpha}\beta(\alpha) \implies \frac{\beta\alpha}{\alpha} = \beta \in R.$$

We can regard $C\ell_R$ as a measure of how non-principal ideals in R can be.

Theorem 8.4. If $R = \mathcal{O}_F$, where F is a number field, then $C\ell_R$ is finite.

This will be proved next week. I will change the label and indexing when it is done.

We start by checking two properties of Dedekind domains.

Proposition 8.5. If A, B, C are ideals of a Dedekind ring R and $AB = AC$, then $B = C$.

Proof. There exists some ideal J such that $JA = (\alpha)$, for $\alpha \in R$. This means $JAB = JAC$, which means $\alpha B = \alpha C$, which we may divide throughout by α to get $B = C$. $\square$

Proposition 8.6. If $A, B \subset R$ are ideals of a Dedekind domain R , then $A \mid B$ (i.e., there exists C such that $B = AC$) if and only if $B \subset A$.

Proof. If $A \mid B$, then $B = CA \subset A$. Suppose $B \subset A$; then there exists an ideal $J \subset R$ such that $JA = (\alpha)$ is principal. Take $C = \alpha^{-1}JB$; then clearly, $CA = B$. Since $B \subset A$, we have $JB \subset JA = (\alpha)$, so $\alpha^{-1}JB \subset R$. $\alpha^{-1}JB$ is an ideal, so we are done. $\square$

Theorem 8.7. Every ideal in a Dedekind domain R is uniquely representable as a product of prime ideals.

Proof. We first check existence, i.e., that every ideal can be expressed as a product of prime ideals. Suppose not; then, since R is Noetherian, there exists a largest proper ideal I that cannot be expressed as a product of prime ideals. Every proper ideal is contained in a maximal ideal, so, in particular, $I \subset M$, for which M being maximal means it is prime. This means that there must exist some J such that $I = JM$. Note that $I \subsetneq J$, since, if $I = J$, then $J = RJ = MJ = 1$, meaning $R = M$, which is a contradiction.

By definition of I , we would have that $J = p_1 \dots p_k$ for prime ideals $p_1, \dots, p_k$. This would mean $I = p_1 \dots p_k M$ as a product of prime ideals, so we have our desired contradiction of our earlier supposition. This means every ideal *can* be expressed as a product of prime ideals.

We now check for uniqueness. Suppose $p_1 \dots p_k = q_1 \dots q_\ell$. Since, for all the prime ideals $p_1, \dots, p_k$, we have $p_1 \dots p_k \subset p_1$, we have $q_1 \dots q_\ell \subset p_1$, so there exists i such that $q_i = p_1$. Without loss of generality, let $i = 1$ by rearranging. By cancellation, $p_2 \dots p_k = q_2 \dots q_\ell$, and we may just induct on the maximum of $\{k, \ell\}$ to finish off the proof. $\square$

With this, we can “do arithmetic with ideals”. Given $I, J \subset R$, we can define things like the greatest common divisor and least common multiple,

$$\gcd(I, J) := I + J, \quad \text{lcm}(I, J) := I \cap J.$$

Theorem 8.8. Let $I \subset R$ be an ideal in a Dedekind domain. Let $\alpha \in I$ be nonzero; then there exists $\beta \in I$ such that $I = (\alpha, \beta)$.

Proof. It is enough to find β such that $I = \gcd((\alpha), (\beta))$. Let $I = p_1^{n_1} \dots p_k^{n_k}$, where $p_1, \dots, p_k$ are distinct primes. Then, let $Q_1, \dots, Q_r$ be other distinct prime ideals that divide (α) . We want $\beta \in \bigcap_i (p_i^{n_i} - p_i^{n_i+1})$, since this would imply that $\beta \in \bigcap_j (R - Q_j)$. Per the chinese remainder theorem, fix $\beta_i \in p_i^{n_i} - p_i^{n_i+1}$, and let β satisfy

$$\beta = \beta_i \pmod{p_i^{n_i+1}};$$

then $\beta = 1 \pmod{Q_j}$. (Recall that, in order to apply the chinese remainder theorem, we would need to show that any pair of ideals is comaximal; but $p_i^{n_i+1} + p_j^{n_j+1} = R$, for which $\gcd(p_i^{n_i+1}, p_j^{n_j+1}) = R$). $\square$

Day 9: Ramification (Oct. 6, 2026)

Theorem 9.1. A Dedekind domain R is a UFD if and only if it is a PID.

Proof. If R is a PID, then it is a UFD. We will check the forwards direction. Suppose R is a UFD and $I \subset R$ is a nonzero ideal. We know that there exists $J \subset R$ such that $IJ = (a)$. Indeed, take the prime factorization $a = p_1 \dots p_r$ (and so $(a) = (p_1) \dots (p_r)$); since p_i being prime implies that the (p_i) are prime ideals, we see that

$$IJ = (p_1) \dots (p_r).$$

By uniqueness of prime factors of ideals, we see that $I = (p_{i_1}) \dots (p_{i_k}) = (p_{i_1} \dots p_{i_k})$, where $\{i_1, \dots, i_k\} \subset \{1, \dots, r\}$, so I is principal. $\square$

We now discuss the splitting of primes. First, we already saw that primes in $\mathbb{Z}$ might factor in other number rings, i.e., in $R = \mathbb{Z}[i]$, $5 = (2 - i)(2 + i)$, but $3 \in \mathbb{Z}[i]$ is prime. There are three kinds of primes in the Gaussian integers. For example, 3 is prime in $\mathbb{Z}[i]$, 5 factors into distinct primes, and 2 factors as the *square* of a prime $(1 + i)^2$.

$R = \mathbb{Z}[\sqrt{-5}]$ is not a PID. 2, 3 are irreducible in R , but they're not prime. Indeed,

$$(2) = (2, 1 + \sqrt{-5})(2, 1 - \sqrt{-5}), \quad (3) = (3, 1 + \sqrt{-5})(3, 1 - \sqrt{-5}).$$

We now discuss the more general setup. Let $K \subset L$ be number fields, and let $R := \mathcal{O}_K$ and $S := \mathcal{O}_L$ (so $R \subset S$).

Theorem 9.2. Let $P \subset R$, $Q \subset S$ be prime ideals; then the following are equivalent.

- (i) Q divides into $PS = \{\alpha_1\beta_1 + \dots + \alpha_r\beta_k \mid \alpha_i \in P, \beta_j \in S\}$;
- (ii) $PS \subset Q$;
- (iii) $P \subset Q$;
- (iv) $Q \cap R = P$;
- (v) $Q \cap K = P$.

Proof. (i) is equivalent to (ii) because divisibility implies containment. (ii) is equivalent to (iii) since PS is the smallest ideal of S containing P . (iv) is equivalent to (v) because $Q \cap K \subset R$. Clearly, (iv) implies (iii), so the only thing that remains to show is (iii) implies (iv). $P \subset Q$ implies $P \subset Q \cap R$; we may observe that $Q \cap R \subset R$ is a proper ideal, i.e., $1 \notin Q \cap R$, since 1 is not in Q . Since P is a nonzero prime, it is maximal, so $P = Q \cap R$. $\square$

Definition 9.3. When conditions of the above theorem are satisfied, we say “ Q lies over P ”, or “ P lies under Q ”.

Theorem 9.4. In the context above,

- (i) Every prime $Q \subset S$ lies over a unique prime $P \subset R$.
- (ii) Every prime $P \subset R$ lies under at least one prime Q of S .

Proof. For (i), we need to show that $Q \cap R$ is a prime ideal. $ab \in Q \cap R$ implies that a or b is in $Q \cap R$. Since $1 \notin Q \cap R$, it follows that $Q \cap R$ is a proper ideal.

For (ii), take Q to be any prime ideal in the factorization of PS . We need to show that $PS \neq S$, i.e., $1 \notin PS$. Recall that there exists $\gamma \in K \setminus R$ such that $\gamma P \subset R$. This means $\gamma PS \subset S$; if $1 \in PS$, then $\gamma \in S$, meaning γ is an algebraic integer (by definition of S being a number field). γ is an algebraic integer contained in K , meaning it must be contained in $R = \mathcal{O}_K$, which is a contradiction. $\square$

We now have that, if $p \subset R$ is prime, then $pS = q_1^{e_1} \dots q_r^{e_r}$, where the q_i 's are distinct prime ideals over p , and the e_i are called *ramification indices*. p is said to be *ramified* if there are exponents greater than 1, i.e., some prime appears with multiplicity higher than 1.

If q lies over p , then the map $R \rightarrow S$ descends to a map

$$R/p \rightarrow S/q$$

in the quotient. Both R/p and S/q are fields, since p, q are maximal, meaning S/q is a field extension of R/p . We denote the *inertial degree* of q over p as $|S/q : R/p|$, usually written $f(q/p)$.

As an example, consider $R = \mathbb{Z}$ and $S = \mathbb{Z}[i]$. 2 lies under $(1 - i)$, and $S/2S$ has order 4, so $|S/(1 - i)|$ is a proper divisor of 4, i.e., $|S/(1 - i)| = 2$. On the other hand, 3 is a prime in $\mathbb{Z}[i]$. $3S \subset S$ is a prime ideal, and $|S/3S| = 9 = 3^2$, so $f(3S/3) = 2$.

With these examples, we see that

$$\sum_{i=1}^r e_i f_i = |L : K|,$$

where $f_i = f(q_i/p)$. As an example, for $\mathbb{Z} \subset \mathbb{Z}[i]$, $(2) = (1 - i)^2$, so $e = 2$ and $f = 1$. Here, $|\mathbb{Q}(i) : \mathbb{Q}| = 2$. We also have $(3) = (3)$, so $e = 1$ and $f = 2$.