

a
Symmetric nested validation
n = 266 paired complexes
median difference = 0.25
95% bootstrap CI 0.175 to 0.308
one-sided Wilcoxon P = 1.03 × 10−19
Number of complexes
Held-out concordance: curated complex − matched null
b
Independent cryo-EM model-quality audit
Global Q-score percentile
versus same-resolution entries
67.5%
Q=0.526
6/6 ARID1A contacts retained
• in both structures
• at 4.0, 4.5 and 5.0 Å
• after bottom-decile residue-Q filtering
0.6%
Q=0.007
6LTJ
3.7 Å
6LTH
3.0 Å
c
Structure–co-dependency sensitivity
Original partition
7 vs 8
Experimental contacts
6 vs 8
Present subunits
6 vs 2
Lung
Pan-cancer
Median ρ difference, 95% bootstrap CI
d
ARID1A-low modules remain null after correction
ARID1A-contacting cBAF
Core shared
cBAF specific
PBAF specific
ncBAF specific
ARID1B paralogue
SMARCA2 ATPase
Positive control:
SMARCA4-low → SMARCA2
q=0.187
q=0.187
q=0.822
q=0.187
q=0.761
q=0.187
q=0.96
q=3.05e-09
Mean within-lineage dependency-z difference, 95% bootstrap
negative = stronger dependency in low-expression group