
a
Deposited structure quality
Same-resolution Q-score percentile
100
80
60
40
20
0
Global Q: 0.007 / 0.526
67.5%
0.6%
6LTJ
3.7 Å
6LTH
3.0 Å
All 6 contacts retained in both deposits at 4.0–5.0 Å,
including removal of bottom-decile residue-Q values.
b
Contact-group sensitivity
lung
pancancer
Genes per group
7 / 8
6 / 8
6 / 2
0.0
0.1
0.2
0.3
0.4
0.5
0.6
Contact − comparison median ρ
Same-assembly (6 / 2): exact P = 0.286
in both cohorts; not statistically supported.
c
Protein-low dependency contrasts
Contact-associated (7 genes)
Core-shared
cBAF-associated
PBAF-associated
ncBAF-associated
ARID1B
SMARCA2
SMARCA4-low → SMARCA2
q = 0.187
q = 0.187
q = 0.822
q = 0.187
q = 0.761
q = 0.187
q = 0.960
q = 3.05 × 10⁻⁹
−1.0
−0.5
0.0
0.5
Mean lineage-standardized dependency difference (low − rest)
Bottom abundance decile; bars: 95% bootstrap intervals. Negative values indicate stronger dependency in the low group.
ARID1A groups use protein abundance; the SMARCA4 positive control uses RNA abundance.