

# MDI343

## Time series : an introduction

François Roueff

<http://perso.telecom-paristech.fr/~roueff/>

Telecom ParisTech

Jan. 7, 2020

# Outline

- 1 Example of time series
- 2 Reminders: i.i.d. models
  - Univariate models
  - Multivariate models
  - Regression model
  - Hidden variables
- 3 Introducing dynamics
  - What's wrong with i.i.d. models ?
  - Univariate models
  - Multivariate models
  - Partially observed multivariate time series
- 4 Stationary Time series
  - The statistical approach
  - Classical steps of statistical inference
  - Stationary and ergodic models
- 5 Weakly stationary time series
  - $L^2$  processes
  - Weak stationarity
  - Spectral measure
  - Empirical estimation

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series

## Example : USD vs EUR currency exchange rate





Figure: Daily currency exchange rate : price of 1 Euro in US Dollars.

## Example : USD vs EUR currency exchange rate (cont.)

Compare with an IID  $\mathcal{N}(0, 1)$  sequence:



## Example : USD vs EUR currency exchange rate (cont.)

Applying the **differencing operator**, we obtain the increment process

$$Y = \Delta X \quad \text{defined by} \quad Y_t = X_t - X_{t-1}, \quad t \in \mathbb{Z}.$$

Makes the “local” mean “more constant”.





Figure: Increments of daily USD-EUR currency exchange rate.

## Example : USD vs EUR currency exchange rate (cont.)

Applying the **differencing operator** of the **logs**, we obtain the **log returns**

$$Y = \Delta \log X \quad \text{defined by} \quad Y_t = \log X_t - \log X_{t-1}, \quad t \in \mathbb{Z}.$$


Makes the “local” mean **and the variance** “more constant”.



**Figure:** Log returns of daily USD-EUR currency exchange rate.

## Example : USD vs EUR currency exchange rate (cont.)

Looking at things “locally” ...



**Figure:** Daily currency exchange rate : price of 1 Euro in US Dollars, on a shorter observation window: between 1999-05-21 and 1999-12-17.

The mean and variance does not appear to vary too much, but still not i.i.d.



- 1 Example of time series
- 2 Reminders: i.i.d. models**
  - Univariate models
  - Multivariate models
  - Regression model
  - Hidden variables
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series

- 1 Example of time series
- 2 Reminders: i.i.d. models
  - Univariate models
  - Multivariate models
  - Regression model
  - Hidden variables
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series

# Discrete observations

- ▷ If we observe i.i.d. **discrete** observations  $X_1, \dots, X_n$ , then the **log-likelihood** can be defined as

$$L_n(\theta) = \sum_{k=1}^n \log p_{\theta}(X_k) ,$$

where, for all  $x$  in the discrete observation space and parameter  $\theta$

$$p_{\theta}(x) = \mathbb{P}_{\theta}(X_1 = x) .$$

# Discrete observations

- ▷ If we observe i.i.d. **discrete** observations  $X_1, \dots, X_n$ , then the **log-likelihood** can be defined as

$$L_n(\theta) = \sum_{k=1}^n \log p_\theta(X_k) ,$$

where, for all  $x$  in the discrete observation space and parameter  $\theta$

$$p_\theta(x) = \mathbb{P}_\theta(X_1 = x) .$$

- ▷ We denote the marginal distribution of  $X_1$  under  $\mathbb{P}_\theta$  by  $\mathbb{P}_\theta^{X_1}$ .

# Discrete observations

- ▷ If we observe i.i.d. **discrete** observations  $X_1, \dots, X_n$ , then the **log-likelihood** can be defined as

$$L_n(\theta) = \sum_{k=1}^n \log p_\theta(X_k) ,$$

where, for all  $x$  in the discrete observation space and parameter  $\theta$

$$p_\theta(x) = \mathbb{P}_\theta(X_1 = x) .$$

- ▷ We denote the marginal distribution of  $X_1$  under  $\mathbb{P}_\theta$  by  $\mathbb{P}_\theta^{X_1}$ .
- ▷ Setting the definition of  $\mathbb{P}_\theta^{X_1}$  or  $p_\theta$  for all  $\theta$  provides a **statistical model** for the observations  $X_1, \dots, X_n$ .

# Examples

▷ Bernoulli model:

$$p_{\theta}(x) = \theta^x (1 - \theta)^{1-x}, \quad \theta \in (0, 1), \quad x \in \{0, 1\}.$$

# Examples

- ▶ Bernoulli model:

$$p_{\theta}(x) = \theta^x (1 - \theta)^{1-x}, \quad \theta \in (0, 1), \quad x \in \{0, 1\}.$$

- ▶ Binomial model (with  $p \geq 1$  known):

$$p_{\theta}(x) = \binom{p}{x} \theta^x (1 - \theta)^{p-x}, \quad \theta \in (0, 1), \quad x \in \{0, \dots, p\}.$$

# Examples

- ▶ Bernoulli model:

$$p_{\theta}(x) = \theta^x(1 - \theta)^{1-x}, \quad \theta \in (0, 1), \quad x \in \{0, 1\}.$$

- ▶ Binomial model (with  $p \geq 1$  known):

$$p_{\theta}(x) = \binom{p}{x} \theta^x (1 - \theta)^{p-x}, \quad \theta \in (0, 1), \quad x \in \{0, \dots, p\}.$$

- ▶ Geometric model:

$$p_{\theta}(x) = \theta(1 - \theta)^{x-1}, \quad \theta \in (0, 1), \quad x \in \mathbb{N}^*.$$



# Examples

- ▶ Bernoulli model:

$$p_{\theta}(x) = \theta^x (1 - \theta)^{1-x}, \quad \theta \in (0, 1), \quad x \in \{0, 1\}.$$

- ▶ Binomial model (with  $p \geq 1$  known):

$$p_{\theta}(x) = \binom{p}{x} \theta^x (1 - \theta)^{p-x}, \quad \theta \in (0, 1), \quad x \in \{0, \dots, p\}.$$

- ▶ Geometric model:

$$p_{\theta}(x) = \theta(1 - \theta)^{x-1}, \quad \theta \in (0, 1), \quad x \in \mathbb{N}^*.$$

- ▶ Negative binomial, Poisson, ...

# Continuous observations

- If we observe i.i.d. **real valued** observations  $X_1, \dots, X_n$ , then the **log-likelihood** can be defined as

$$L_n(\theta) = \sum_{k=1}^n \log p_{\theta}(X_k) ,$$

where, for all  $x$  in the discrete observation space and parameter  $\theta$ ,  $p_{\theta}$  is the density of  $\mathbb{P}_{\theta}^{X_1}$ :

$$\mathbb{P}_{\theta}^{X_1}(A) = \mathbb{P}_{\theta}(X_1 \in A) = \int_A p_{\theta}(x) \, dx .$$

# Continuous observations

- ▷ If we observe i.i.d. **real valued** observations  $X_1, \dots, X_n$ , then the **log-likelihood** can be defined as

$$L_n(\theta) = \sum_{k=1}^n \log p_\theta(X_k) ,$$

where, for all  $x$  in the discrete observation space and parameter  $\theta$ ,  $p_\theta$  is the density of  $\mathbb{P}_\theta^{X_1}$ :

$$\mathbb{P}_\theta^{X_1}(A) = \mathbb{P}_\theta(X_1 \in A) = \int_A p_\theta(x) \, dx .$$

- ▷ Again, setting the definition of  $\mathbb{P}_\theta^{X_1}$  or  $p_\theta$  for all  $\theta$  provides a **statistical model** for the observations  $X_1, \dots, X_n$ .

# Examples

▷ Exponential model:

$$p_{\lambda}(x) = \lambda e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x) , \quad \lambda > 0 .$$

# Examples

- ▷ Exponential model:

$$p_{\lambda}(x) = \lambda e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \lambda > 0.$$

- ▷ Gamma model

$$p_{\theta}(x) = \frac{\lambda^{\alpha}}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \theta = (\alpha, \lambda) \in \mathbb{R}_+^{*2}.$$

# Examples

- ▷ Exponential model:

$$p_{\lambda}(x) = \lambda e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \lambda > 0.$$

- ▷ Gamma model

$$p_{\theta}(x) = \frac{\lambda^{\alpha}}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \theta = (\alpha, \lambda) \in \mathbb{R}_+^{*2}.$$

- ▷ Beta model :

$$p_{\theta}(x) = B(\alpha, \beta) x^{\alpha-1} (1-x)^{\beta-1} \mathbb{1}_{[0,1]}(x), \quad \theta = (\alpha, \beta) \in \mathbb{R}_+^{*2}.$$

# Examples

- ▶ Exponential model:

$$p_{\lambda}(x) = \lambda e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \lambda > 0.$$

- ▶ Gamma model

$$p_{\theta}(x) = \frac{\lambda^{\alpha}}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} \mathbb{1}_{\mathbb{R}_+}(x), \quad \theta = (\alpha, \lambda) \in \mathbb{R}_+^{*2}.$$

- ▶ Beta model :

$$p_{\theta}(x) = B(\alpha, \beta) x^{\alpha-1} (1-x)^{\beta-1} \mathbb{1}_{[0,1]}(x), \quad \theta = (\alpha, \beta) \in \mathbb{R}_+^{*2}.$$

- ▶ Gaussian model:

$$p_{\theta}(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)}, \quad \theta = (\mu, \sigma^2) \in \mathbb{R} \times \mathbb{R}_+^{*}.$$

- 1 Example of time series
- 2 Reminders: i.i.d. models
  - Univariate models
  - **Multivariate models**
  - Regression model
  - Hidden variables
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series



# Multivariate data

- ▶ Most real life data is multivariate in the sense that it is doubly indexed, e.g.

$$\mathbf{X}_t = X_{i,t} \quad i = 1, \dots, p,$$

where  $t$  is the time index and  $i$  enumerates individuals, assets, features etc.

# Multivariate data

- ▶ Most real life data is multivariate in the sense that it is doubly indexed, e.g.

$$\mathbf{X}_t = X_{i,t} \quad i = 1, \dots, p,$$

where  $t$  is the time index and  $i$  enumerates individuals, assets, features etc.

- ▶ Examples: **portfolio** returns, **panel** data (or **longitudinal** data), Risk indices, ...

# Multivariate data

- ▶ Most real life data is multivariate in the sense that it is doubly indexed, e.g.

$$\mathbf{X}_t = X_{i,t} \quad i = 1, \dots, p,$$

where  $t$  is the time index and  $i$  enumerates individuals, assets, features etc.

- ▶ Examples: **portfolio** returns, **panel** data (or **longitudinal** data), Risk indices, ...
- ▶ To simplify the presentation, let us see the index  $i$  as a **spatial** index (as opposed to **time index**).

# Multivariate data

- ▶ Most real life data is multivariate in the sense that it is doubly indexed, e.g.

$$\mathbf{X}_t = X_{i,t} \quad i = 1, \dots, p,$$

where  $t$  is the time index and  $i$  enumerates individuals, assets, features etc.

- ▶ Examples: **portfolio** returns, **panel** data (or **longitudinal** data), Risk indices, ...
- ▶ To simplify the presentation, let us see the index  $i$  as a **spatial** index (as opposed to **time index**).
- ▶ A multivariate model will generally try to capture the *spatial* covariance structure through **random vector** models: e.g. **Gaussian vectors**, **Ising model**, or more general **graphical models**...

## Example: i.i.d. Gaussian vectors

- ▷ Consider a portfolio of  $n$  asset returns  $\mathbf{X}_t = X_{i,t} \quad i = 1, \dots, p$ .
- ▷ Suppose that  $\mathbf{X}_1, \dots, \mathbf{X}_n$  are i.i.d.  $\mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ , where
  - ▷  $\boldsymbol{\mu} \in \mathbb{R}^p$  is the unknown mean.
  - ▷  $\boldsymbol{\Sigma} \in \mathbb{R}^{p \times p}$  is the unknown covariance matrix
- ▷ Then the log-likelihood reads, for all  $\boldsymbol{\theta} = (\boldsymbol{\mu}, \boldsymbol{\Sigma})$ ,

$$\begin{aligned} L_n(\boldsymbol{\theta}) &= \sum_{k=1}^n \log p_{\boldsymbol{\theta}}(\mathbf{X}_k) \\ &= -\frac{1}{2n} \left( \log \det(2\pi \boldsymbol{\Sigma}) + \sum_{k=1}^n (\mathbf{X}_k - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{X}_k - \boldsymbol{\mu}) \right). \end{aligned}$$

## Example: i.i.d. Gaussian vectors, estimators

- ▶ Using a classical moment estimation method, we obtain the **empirical estimators**:
  - ▶ the empirical mean

$$\hat{\mu}_{n,i} = \frac{1}{n} \sum_{t=1}^n \mathbf{x}_{i,t} .$$

- ▶ the empirical covariance matrix

$$\hat{\Sigma}_n[i, j] = \frac{1}{n} \sum_{t=1}^n (\mathbf{x}_{i,t} - \hat{\mu}_{n,i})(\mathbf{x}_{j,t} - \hat{\mu}_{n,j}) .$$

## Example: i.i.d. Gaussian vectors, estimators

- ▶ Using a classical moment estimation method, we obtain the **empirical estimators**:
  - ▶ the empirical mean

$$\hat{\mu}_{n,i} = \frac{1}{n} \sum_{t=1}^n \mathbf{x}_{i,t} .$$

- ▶ the empirical covariance matrix

$$\hat{\Sigma}_n[i, j] = \frac{1}{n} \sum_{t=1}^n (\mathbf{x}_{i,t} - \hat{\mu}_{n,i})(\mathbf{x}_{j,t} - \hat{\mu}_{n,j}) .$$

- ▶ Maximizing the likelihood yields to much more complicated estimators...

## Example: i.i.d. Gaussian vectors, estimators

- ▶ Using a classical moment estimation method, we obtain the **empirical estimators**:
  - ▶ the empirical mean

$$\hat{\mu}_{n,i} = \frac{1}{n} \sum_{t=1}^n \mathbf{x}_{i,t} .$$

- ▶ the empirical covariance matrix

$$\hat{\Sigma}_n[i, j] = \frac{1}{n} \sum_{t=1}^n (\mathbf{x}_{i,t} - \hat{\mu}_{n,i})(\mathbf{x}_{j,t} - \hat{\mu}_{n,j}) .$$

- ▶ Maximizing the likelihood yields to much more complicated estimators...
- ▶ In **high dimension** ( $p$  and  $n$  are of similar order), it is sometimes advantageous to make a **sparse** or **low rank** assumption.



## Example: i.i.d. Gaussian vectors, estimators

- ▶ Using a classical moment estimation method, we obtain the **empirical estimators**:
  - ▶ the empirical mean

$$\hat{\mu}_{n,i} = \frac{1}{n} \sum_{t=1}^n \mathbf{x}_{i,t} .$$

- ▶ the empirical covariance matrix

$$\hat{\Sigma}_n[i, j] = \frac{1}{n} \sum_{t=1}^n (\mathbf{x}_{i,t} - \hat{\mu}_{n,i})(\mathbf{x}_{j,t} - \hat{\mu}_{n,j}) .$$

- ▶ Maximizing the likelihood yields to much more complicated estimators...
- ▶ In **high dimension** ( $p$  and  $n$  are of similar order), it is sometimes advantageous to make a **sparse** or **low rank** assumption.
- ▶ From a **regression** perspective, it is easier to use sparsity of the **precision matrix**  $\mathbf{M} = \Sigma^{-1}$  .

- 1 Example of time series
- 2 Reminders: i.i.d. models
  - Univariate models
  - Multivariate models
  - Regression model
  - Hidden variables
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series

# From bivariate distribution to conditional distribution

- ▶ In a regression model, each multivariate observation  $\mathbf{X}_i$  is split into a pair of variables :  $\mathbf{X}_i = (\mathbf{Z}_i, Y_i)$ , where, usually,  $\mathbf{Z}_i$  itself is multivariate, say valued in  $\mathbb{R}^p$ , and  $Y_i$  is univariate (discrete or continuous).

# From bivariate distribution to conditional distribution

- ▶ In a regression model, each multivariate observation  $\mathbf{X}_i$  is split into a pair of variables :  $\mathbf{X}_i = (\mathbf{Z}_i, Y_i)$ , where, usually,  $\mathbf{Z}_i$  itself is multivariate, say valued in  $\mathbb{R}^p$ , and  $Y_i$  is univariate (discrete or continuous).
- ▶ Thus we can see the distribution of  $\mathbf{X}_1$  as a **bivariate** distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$ .

# From bivariate distribution to conditional distribution

- ▶ In a regression model, each multivariate observation  $\mathbf{X}_i$  is split into a pair of variables :  $\mathbf{X}_i = (\mathbf{Z}_i, Y_i)$ , where, usually,  $\mathbf{Z}_i$  itself is multivariate, say valued in  $\mathbb{R}^p$ , and  $Y_i$  is univariate (discrete or continuous).
- ▶ Thus we can see the distribution of  $\mathbf{X}_1$  as a **bivariate** distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$ .
- ▶ Every bivariate distribution can be decomposed using
  - ▶ the **marginal** distribution of the **first** variable;
  - ▶ the **conditional** distribution of the **second** variable **given** the first variable.

# From bivariate distribution to conditional distribution

- ▶ In a regression model, each multivariate observation  $\mathbf{X}_i$  is split into a pair of variables :  $\mathbf{X}_i = (\mathbf{Z}_i, Y_i)$ , where, usually,  $\mathbf{Z}_i$  itself is multivariate, say valued in  $\mathbb{R}^p$ , and  $Y_i$  is univariate (discrete or continuous).
- ▶ Thus we can see the distribution of  $\mathbf{X}_1$  as a **bivariate** distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$ .
- ▶ Every bivariate distribution can be decomposed using
  - ▶ the **marginal** distribution of the **first** variable;
  - ▶ the **conditional** distribution of the **second** variable **given** the first variable.
- ▶ In a **regression model**, we see  $\mathbf{Z}_i$  as an **input** (**regression variable**) and  $Y_i$  as an **output** (**observation or response variable**) and are only interested on the conditional distribution of the output given the input.

# Likelihood of a regression model

- ▷ The decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$  then yields

$$p_{\theta}(\mathbf{x}) = q(\mathbf{z})p_{\theta}(y|\mathbf{z}) , \quad \mathbf{x} = (\mathbf{z}, y) ,$$

where  $q(\mathbf{z})$  denotes the density of  $\mathbf{Z}_1$  and  $p_{\theta}(y|\mathbf{z})$  denotes the conditional density of  $Y_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $\mathbf{Z}_1 = \mathbf{z}$  under parameter  $\theta$ .

# Likelihood of a regression model

- ▷ The decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$  then yields

$$p_{\theta}(\mathbf{x}) = q(\mathbf{z})p_{\theta}(y|\mathbf{z}) , \quad \mathbf{x} = (\mathbf{z}, y) ,$$

where  $q(\mathbf{z})$  denotes the density of  $\mathbf{Z}_1$  and  $p_{\theta}(y|\mathbf{z})$  denotes the conditional density of  $Y_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $\mathbf{Z}_1 = \mathbf{z}$  under parameter  $\theta$ .

- ▷ It follows that the log-likelihood takes the form, up to an additive constant:

$$\log L_n(\theta) = \sum_{k=1}^n \log p_{\theta}(Y_k | \mathbf{Z}_k)$$



# Likelihood of a regression model

- ▷ The decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1} = \mathbb{P}_{\theta}^{(\mathbf{Z}_1, Y_1)}$  then yields

$$p_{\theta}(\mathbf{x}) = q(\mathbf{z})p_{\theta}(y|\mathbf{z}) , \quad \mathbf{x} = (\mathbf{z}, y) ,$$

where  $q(\mathbf{z})$  denotes the density of  $\mathbf{Z}_1$  and  $p_{\theta}(y|\mathbf{z})$  denotes the conditional density of  $Y_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $\mathbf{Z}_1 = \mathbf{z}$  under parameter  $\theta$ .

- ▷ It follows that the log-likelihood takes the form, up to an additive constant:

$$\log L_n(\theta) = \sum_{k=1}^n \log p_{\theta}(Y_k | \mathbf{Z}_k)$$

- ▷ Estimating  $\theta$  allows one to propose a predictor of  $Y$  given a new input  $\mathbf{Z}$ , assuming that they are distributed according to the same bivariate distribution as the learning data set.

## Two examples

- ▶ The linear regression model:

$$p_{\boldsymbol{\theta}, \sigma^2}(y|\mathbf{z}) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(y - \boldsymbol{\theta}^T \mathbf{z})^2 / (2\sigma^2)}, \quad (\boldsymbol{\theta}, \sigma^2) \in \mathbb{R}^p \times \mathbb{R}_+^*, \quad y \in \mathbb{R}.$$

Optimizing the likelihood leads to the **least mean square** estimator.

## Two examples

- ▶ The linear regression model:

$$p_{\boldsymbol{\theta}, \sigma^2}(y|\mathbf{z}) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(y - \boldsymbol{\theta}^T \mathbf{z})^2 / (2\sigma^2)}, \quad (\boldsymbol{\theta}, \sigma^2) \in \mathbb{R}^p \times \mathbb{R}_+^*, \quad y \in \mathbb{R}.$$

Optimizing the likelihood leads to the **least mean square** estimator.

- ▶ The logit regression model:

$$p_{\boldsymbol{\theta}}(y|\mathbf{z}) = \left( \frac{e^{\boldsymbol{\theta}^T \mathbf{z}}}{1 + e^{\boldsymbol{\theta}^T \mathbf{z}}} \right)^y \left( \frac{1}{1 + e^{\boldsymbol{\theta}^T \mathbf{z}}} \right)^{1-y}, \quad \boldsymbol{\theta} \in \mathbb{R}^p, \quad y \in \{0, 1\}.$$

- 1 Example of time series
- 2 Reminders: i.i.d. models
  - Univariate models
  - Multivariate models
  - Regression model
  - **Hidden variables**
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series

# The mixture model

- ▶ In certain cases, each (univariate or multivariate) observation  $\mathbf{X}_i$  can be seen as a **partial observation** of a larger set of data. The unobserved part of the data is represented by a **hidden** variable  $V_i$ .

# The mixture model

- ▶ In certain cases, each (univariate or multivariate) observation  $\mathbf{X}_i$  can be seen as a **partial observation** of a larger set of data. The unobserved part of the data is represented by a **hidden** variable  $V_i$ .
- ▶ Again we can then decompose the **bivariate** distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  of the complete data  $(V_1, \mathbf{X}_1)$  using
  - ▶ the **marginal** distribution of the hidden variable  $V_1$ ;
  - ▶ the **conditional** distribution of the observed variable  $\mathbf{X}_1$  **given** the hidden variable  $V_1$ .

# The mixture model

- ▶ In certain cases, each (univariate or multivariate) observation  $\mathbf{X}_i$  can be seen as a **partial observation** of a larger set of data. The unobserved part of the data is represented by a **hidden** variable  $V_i$ .
- ▶ Again we can then decompose the **bivariate** distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  of the complete data  $(V_1, \mathbf{X}_1)$  using
  - ▶ the **marginal** distribution of the hidden variable  $V_1$ ;
  - ▶ the **conditional** distribution of the observed variable  $\mathbf{X}_1$  given the hidden variable  $V_1$ .
- ▶ The resulting marginal distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1}$  is called a **mixture model**.

# The mixture model

- ▶ In certain cases, each (univariate or multivariate) observation  $\mathbf{X}_i$  can be seen as a **partial observation** of a larger set of data. The unobserved part of the data is represented by a **hidden** variable  $V_i$ .
- ▶ Again we can then decompose the **bivariate** distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  of the complete data  $(V_1, \mathbf{X}_1)$  using
  - ▶ the **marginal** distribution of the hidden variable  $V_1$ ;
  - ▶ the **conditional** distribution of the observed variable  $\mathbf{X}_1$  given the hidden variable  $V_1$ .
- ▶ The resulting marginal distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1}$  is called a **mixture model**.
- ▶ The simplest case is that of a **finite mixture**, where the hidden variable takes its values in a finite set  $\{1, 2, \dots, K\}$ . This case amounts to see the data as being separated into  $K$  **clusters**, each of them following a different distribution, namely, the **conditional distribution** of  $\mathbf{X}_1$  given  $V_1 = k$ , for  $k = 1, 2, \dots, K$ .



# The mixture model

- ▶ In certain cases, each (univariate or multivariate) observation  $\mathbf{X}_i$  can be seen as a **partial observation** of a larger set of data. The unobserved part of the data is represented by a **hidden** variable  $V_i$ .
- ▶ Again we can then decompose the **bivariate** distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  of the complete data  $(V_1, \mathbf{X}_1)$  using
  - ▶ the **marginal** distribution of the hidden variable  $V_1$ ;
  - ▶ the **conditional** distribution of the observed variable  $\mathbf{X}_1$  given the hidden variable  $V_1$ .
- ▶ The resulting marginal distribution  $\mathbb{P}_{\theta}^{\mathbf{X}_1}$  is called a **mixture model**.
- ▶ The simplest case is that of a **finite mixture**, where the hidden variable takes its values in a finite set  $\{1, 2, \dots, K\}$ . This case amounts to see the data as being separated into  $K$  **clusters**, each of them following a different distribution, namely, the **conditional distribution** of  $\mathbf{X}_1$  given  $V_1 = k$ , for  $k = 1, 2, \dots, K$ .
- ▶ A standard example of hidden variable for **financial data** is the (conditional) **volatility**.

# Likelihood of a mixture model

- ▷ The natural decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  yields

$$p_{\theta}(v, \mathbf{x}) = q_{\theta}(v)p_{\theta}(\mathbf{x}|v) ,$$

where  $q_{\theta}(v)$  denotes the density of  $V_1$  (or the probability of  $V_1 = v$ ) and  $p_{\theta}(\mathbf{x}|v)$  denotes the conditional density of  $\mathbf{X}_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $V_1 = v$  under parameter  $\theta$ .

# Likelihood of a mixture model

- ▷ The natural decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  yields

$$p_{\theta}(v, \mathbf{x}) = q_{\theta}(v)p_{\theta}(\mathbf{x}|v) ,$$

where  $q_{\theta}(v)$  denotes the density of  $V_1$  (or the probability of  $V_1 = v$ ) and  $p_{\theta}(\mathbf{x}|v)$  denotes the conditional density of  $\mathbf{X}_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $V_1 = v$  under parameter  $\theta$ .

- ▷ It follows that the log-likelihood takes the form (in the case of continuous hidden variables):

$$\log L_n(\theta) = \sum_{k=1}^n \log \int q_{\theta}(v) p_{\theta}(\mathbf{X}_k|v) \, dv .$$

# Likelihood of a mixture model

- ▶ The natural decomposition of the bivariate distribution  $\mathbb{P}_{\theta}^{(V_1, \mathbf{X}_1)}$  yields

$$p_{\theta}(v, \mathbf{x}) = q_{\theta}(v) p_{\theta}(\mathbf{x}|v) ,$$

where  $q_{\theta}(v)$  denotes the density of  $V_1$  (or the probability of  $V_1 = v$ ) and  $p_{\theta}(\mathbf{x}|v)$  denotes the conditional density of  $\mathbf{X}_1$  (or the conditional probability of  $\mathbf{X}_1 = \mathbf{x}$ ) given  $V_1 = v$  under parameter  $\theta$ .

- ▶ It follows that the log-likelihood takes the form (in the case of continuous hidden variables):

$$\log L_n(\theta) = \sum_{k=1}^n \log \int q_{\theta}(v) p_{\theta}(\mathbf{X}_k|v) \, dv .$$

- ▶ For discrete mixtures, estimating  $\theta$  allows one to clustering the data by identifying those who most likely share the same hidden variable.

## Two examples

- ▷ Mixture of two Gaussian variables with parameter  $\theta = (\alpha, \mu_0, \mu_1, \sigma_0^2, \sigma_1^2) \in (0, 1) \times \mathbb{R}^2 \times \mathbb{R}_+^{*2}$ :  $V_1 \sim \text{Bernoulli}(\alpha)$  and given  $V_1 = v$ ,  $X_1 \sim \mathcal{N}(\mu_v, \sigma_v^2)$ . Hence

$$q_{\theta}(v) = \alpha^v (1 - \alpha)^{1-v}$$
$$p_{\theta}(x|v) = (2\pi\sigma_v^2)^{-1/2} e^{-(x-\mu_v)^2/(2\sigma_v^2)}.$$

- ▷ Discrete mixture of Gaussian vectors with parameter  $\theta = (\alpha_k, \mu_k, \Sigma_k)_{1 \leq k \leq K}$  :

$$q_{\theta}(v) = \alpha_v$$
$$p_{\theta}(\mathbf{x}|v) = (\det(2\pi\Sigma_v))^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{x} - \mu_v)^T \Sigma_v^{-1}(\mathbf{x} - \mu_v)\right)$$

Optimizing the likelihood is a difficult question (related to the  $k$ -means algorithm).

## Two examples (cont)





Figure: Density of the mixture of two Gaussian distributions

## Two examples (cont)




**Figure:** IID draws of the mixture of 5 bidimensional Gaussian distributions. Colors represent the (supposedly hidden) cluster variables.

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics**
  - What's wrong with i.i.d. models ?
  - Univariate models
  - Multivariate models
  - Partially observed multivariate time series
- 4 Stationary Time series
- 5 Weakly stationary time series



- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
  - What's wrong with i.i.d. models ?
  - Univariate models
  - Multivariate models
  - Partially observed multivariate time series
- 4 Stationary Time series
- 5 Weakly stationary time series

## Back to the USD vs EUR currency exchange rate.



**Figure:** Top : price of 1 Euro in US Dollars between 1999-05-21 and 1999-12-17;  
Bottom : the same in randomly shuffled order.

# Order of observations is not taken into account in i.i.d. models

- ▶ The log-likelihood of an i.i.d. model has the form

$$L_n(\theta) = \sum_{k=1}^n \log p_{\theta}(X_k) ,$$

where  $X_1, \dots, X_n$  are the  $n$  observations, hence is **invariant** through **permutation** of indices:  $(X_1, \dots, X_n) \mapsto (X_{\sigma(1)}, \dots, X_{\sigma(n)})$ , where  $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$  is a permutation.

- ▶ The two previous time series are the same **up to a permutation of time indices**.
- ▶ Hence they have the **same likelihood** for any i.i.d. model.

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics**
  - What's wrong with i.i.d. models ?
  - Univariate models**
  - Multivariate models
  - Partially observed multivariate time series
- 4 Stationary Time series
- 5 Weakly stationary time series

## Some useful notation

- ▷ For any integers  $k \geq l$  and sequence  $(x_t)$  we denote the subsample with indices between  $k$  and  $l$  by

$$x_{k:l} = (x_k, \dots, x_l)$$

- ▷ If  $(\mathbf{X}, \mathbf{Y})$  is valued in  $\mathbb{R}^p \times \mathbb{R}^n$  and admits a density, we denote
- ▷ by  $p^{(\mathbf{X}, \mathbf{Y})} : (x, y) \mapsto p^{(\mathbf{X}, \mathbf{Y})}(x, y)$  the density of  $(\mathbf{X}, \mathbf{Y})$ ,
  - ▷ by  $p^{\mathbf{X}}$  the density of  $\mathbf{X}$  :

$$p^{\mathbf{X}}(\mathbf{x}) = \int_{\mathbb{R}^n} p^{(\mathbf{X}, \mathbf{Y})}(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} = \int \cdots \int p^{(\mathbf{X}, \mathbf{Y})}(\mathbf{x}, y_{1:n}) \, dy_1 \cdots dy_n .$$

- ▷ by  $p^{\mathbf{Y}|\mathbf{X}}(\cdot|x)$  the conditional density of  $\mathbf{Y}$  given  $\mathbf{X} = x$  :

$$p^{\mathbf{Y}|\mathbf{X}}(y|x) = \frac{p^{(\mathbf{X}, \mathbf{Y})}(x, y)}{p^{\mathbf{X}}(x)}$$

- ▷ We add a subscript  $\theta$  if the density depends on the unknown parameter  $\theta$ :  $p_{\theta}^{(\mathbf{X}, \mathbf{Y})}$ ,  $p_{\theta}^{\mathbf{X}}$ ,  $p_{\theta}^{\mathbf{Y}|\mathbf{X}}$  ...

# General form of the likelihood

- ▶ How to generalize the product form of likelihood without the i.i.d. assumption ?

# General form of the likelihood

- ▶ How to generalize the product form of likelihood without the i.i.d. assumption ?
- ▶ Suppose that  $X_{1:n}$  admits a density  $p_{\theta}^{X_{1:n}}$ .

# General form of the likelihood

- ▷ How to generalize the product form of likelihood without the i.i.d. assumption ?
- ▷ Suppose that  $X_{1:n}$  admits a density  $p_{\theta}^{X_{1:n}}$ .
- ▷ Conditioning successively, we have

$$\begin{aligned} p_{\theta}^{X_{1:n}}(x_{1:n}) &= p_{\theta}^{X_n | X_{1:(n-1)}}(x_n | x_{1:n-1}) p_{\theta}^{X_{1:n-1}}(x_{1:n-1}) \\ &\dots \\ &= \prod_{k=2}^n p_{\theta}^{X_k | X_{1:(k-1)}}(x_k | x_{1:k-1}) p_{\theta}^{X_1}(x_1) . \end{aligned}$$



# General form of the likelihood

- ▶ How to generalize the product form of likelihood without the i.i.d. assumption ?
- ▶ Suppose that  $X_{1:n}$  admits a density  $p_{\theta}^{X_{1:n}}$ .
- ▶ Conditioning successively, we have

$$\begin{aligned} p_{\theta}^{X_{1:n}}(x_{1:n}) &= p_{\theta}^{X_n | X_{1:(n-1)}}(x_n | x_{1:n-1}) p_{\theta}^{X_{1:n-1}}(x_{1:n-1}) \\ &\quad \dots \\ &= \prod_{k=2}^n p_{\theta}^{X_k | X_{1:(k-1)}}(x_k | x_{1:k-1}) p_{\theta}^{X_1}(x_1) . \end{aligned}$$

- ▶ It is therefore of primary importance to understand the **dynamics** of the model through the **conditional distribution** of  $X_k$  given its **past**  $X_{1:(k-1)}$ .

## Two important particular cases

▷ The i.i.d. case :

In this case, by independence of  $X_k$  and  $X_{1:(k-1)}$ , we have that  $p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1})$  does not depend on  $x_{1:k-1}$ , so that

$$p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1}) = p_{\theta}^{X_k}(x_k) .$$

And, by the "i.d." property,

$$p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1}) = p_{\theta}^{X_k}(x_k) = p_{\theta}(x_k) ,$$

where  $p_{\theta}$  is the common density of all  $X_k$ 's.

## Two important particular cases (cont.)

▷ The **homogeneous Markov** case :


In this case, we have that  $p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1})$  **only** depends on  $x_{k-1}$ , so that

$$p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1}) = p_{\theta}^{X_k|X_{k-1}}(x_k|x_{k-1}) .$$

And “homogeneous” means that  $p_{\theta}^{X_k|X_{k-1}}$  does not depend on  $k$  and is given by a **common conditional density**, say  $q_{\theta}(\cdot|\cdot)$ , hence

$$p_{\theta}^{X_k|X_{1:(k-1)}}(x_k|x_{1:k-1}) = p_{\theta}^{X_k|X_{k-1}}(x_k|x_{k-1}) = q_{\theta}(x_k|x_{k-1}) .$$

# Graphical representation of a homogeneous Markov chain



- ▶ Arrows indicate the dependence structure: given all other variables, a **child** can be generated using only its own **parents**.
- ▶ Here, each child only has 1 parent: the generation of the child is carried out through the conditional density  $q_{\theta}$ .

# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

$$X_{k-1}$$

×

# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

$$\begin{array}{ccc} X_{k-1} & & \\ \times & & \\ & \phi X_{k-1} & \\ & \times & \end{array}$$

# Examples of conditional density


An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .





# Examples of conditional density


An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .



# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

$X_{k-1}$   
×

$X_k$   
×

# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

$X_{k-1}$   
×

$X_k$   
×

$\phi X_k$   
×



# Examples of conditional density


An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .



# Examples of conditional density


An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .



# Examples of conditional density

An homoscedastic model : AR(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(\phi x, \sigma^2)$ , with  $\theta = (\phi, \sigma^2) \in (-1, 1) \times \mathbb{R}_+^*$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \phi X_{k-1} + \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, \sigma^2)$ .

$X_{k-1}$   
×

$X_k$   
×

$X_{k+1}$   
×



## Examples of conditional density (cont.)

An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

# Examples of conditional density (cont.)

An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

$$X_{k-1}$$

×





# Examples of conditional density (cont.)


An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



# Examples of conditional density (cont.)


## An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



# Examples of conditional density (cont.)


## An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



# Examples of conditional density (cont.)

An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

$X_{k-1}$   
×

→  
 $X_k$   
×

# Examples of conditional density (cont.)


An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



# Examples of conditional density (cont.)


## An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



# Examples of conditional density (cont.)


## An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .



## Examples of conditional density (cont.)


An heteroscedastic model : ARCH(1).

In this case,  $q_{\theta}(\cdot|x)$  is the density of  $\mathcal{N}(0, a + bx^2)$ , with  $\theta = (a, b) \in \mathbb{R}_+^* \times \mathbb{R}_+$ .

Equivalently, this model is given by the dynamical equation

$$X_k = \sqrt{a + bX_{k-1}^2} \epsilon_k ,$$

with  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .





# Likelihood for univariate time series

- ▶ The simplest model beyond the i.i.d. model that contains some non-trivial dynamics is the **homogeneous Markov** model.

# Likelihood for univariate time series

- ▶ The simplest model beyond the i.i.d. model that contains some non-trivial dynamics is the **homogeneous Markov** model.
- ▶ Its log-likelihood takes the form

$$L_n(\theta) = \sum_{k=2}^n \log q_{\theta}(X_k | X_{k-1}) + \log p^{X_1}(X_1) .$$

# Likelihood for univariate time series

- ▶ The simplest model beyond the i.i.d. model that contains some non-trivial dynamics is the **homogeneous Markov** model.
- ▶ Its **conditional** log-likelihood takes the form

$$L_n(\theta) = \sum_{k=2}^n \log q_{\theta}(X_k | X_{k-1}) + \cancel{\log p^{X_1}(X_1)}.$$

# Likelihood for univariate time series

- ▶ The simplest model beyond the i.i.d. model that contains some non-trivial dynamics is the **homogeneous Markov** model.
- ▶ Its **conditional** log-likelihood takes the form

$$L_n(\theta) = \sum_{k=2}^n \log q_{\theta}(X_k | X_{k-1}) + \cancel{\log p^{X_1}(X_1)} .$$

- ▶ The likelihood is no longer invariant by permutation.

## Exemple: likelihood of the Gaussian AR(1) model

Consider the ► AR(1) model. Then we have

$$q_{\theta}(x_k|x_{k-1}) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x_k - \phi x_{k-1})^2 / (2\sigma^2)} .$$

It follows that the (conditional) negated log likelihood reads

$$-L_n(\theta) = \frac{n-1}{2} \log(2\pi\sigma^2) + \frac{1}{2\sigma^2} \sum_{k=2}^n (X_k - \phi X_{k-1})^2 ,$$

which leads to the estimators

$$\hat{\phi}_n = \frac{\sum_{k=2}^n X_{k-1} X_k}{\sum_{k=2}^n X_{k-1}^2} \quad \text{and} \quad \hat{\sigma}_n^2 = \frac{1}{n-1} \sum_{k=2}^n (X_k - \hat{\phi}_n X_{k-1})^2 .$$

## Exemple: likelihood of the conditionally Gaussian ARCH(1) model

Consider the ► ARCH(1) model. Then we have

$$q_{\theta}(x_k|x_{k-1}) = \frac{1}{\sqrt{2\pi(a + bx_{k-1}^2)}} e^{-x_k^2/(2(a + bx_{k-1}^2))} .$$

It follows that the (conditional) negated log likelihood reads

$$-L_n(\theta) = \frac{1}{2} \sum_{k=2}^n \left( \log(2\pi(a + bX_{k-1}^2)) + \frac{X_k^2}{a + bX_{k-1}^2} \right) ,$$

which can be minimized in  $\theta = (a, b)$  using a gradient descent algorithm.

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 **Introducing dynamics**
  - What's wrong with i.i.d. models ?
  - Univariate models
  - **Multivariate models**
  - Partially observed multivariate time series
- 4 Stationary Time series
- 5 Weakly stationary time series

# Multivariate time series

- ▶ Exactly as in the IID case, a time series  $(\mathbf{X}_t)$  can be multivariate, i.e.  $\mathbf{X}_t$  is valued in  $\mathbb{R}^p$  for some  $p \geq 2$ .



# Multivariate time series

- ▷ Exactly as in the IID case, a time series  $(\mathbf{X}_t)$  can be multivariate, i.e.  $\mathbf{X}_t$  is valued in  $\mathbb{R}^p$  for some  $p \geq 2$ .
- ▷ For instance, under the homogeneous Markov chain assumption, the (conditional) likelihood then reads

$$L_n(\theta) = \sum_{k=2}^n \log q_{\theta}(\mathbf{X}_k | \mathbf{X}_{k-1}) .$$

# Multivariate time series

- ▷ Exactly as in the IID case, a time series  $(\mathbf{X}_t)$  can be multivariate, i.e.  $\mathbf{X}_t$  is valued in  $\mathbb{R}^p$  for some  $p \geq 2$ .
- ▷ For instance, under the homogeneous Markov chain assumption, the (conditional) likelihood then reads

$$L_n(\theta) = \sum_{k=2}^n \log q_\theta(\mathbf{X}_k | \mathbf{X}_{k-1}) .$$

- ▷ In particular, consider a univariate  $p$ -order Markov time series with log likelihood

$$L_n(\theta) = \sum_{k=p+2}^n \log q_\theta(X_k | X_{k-p:k-1}) .$$

To obtain a multivariate (first order) Markov time series, one can set  $\mathbf{X}_k = X_{k-p+1:k}$ .

## Exemple of Multivariate time series: AR( $p$ ) time series

An AR( $p$ ) time series ( $X_t$ ) satisfies the AR( $p$ ) equation

$$X_t = \sum_{k=1}^p \phi_k X_{t-k} + \epsilon_t, \quad t \in \mathbb{Z}.$$

Setting  $\mathbf{X}_k = [X_k \ X_{k-1} \ \dots \ X_{k-p+1}]^T$ , this leads to the vector AR(1) equation:

$$\mathbf{X}_t = \Phi \mathbf{X}_{t-1} + \epsilon_t, \quad t \in \mathbb{Z}.$$


where

$$\Phi = \begin{bmatrix} \phi_1 & \phi_2 & \dots & \phi_p \\ 1 & 0 & \dots & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & \dots & 1 & 0 \end{bmatrix} \quad \text{and} \quad \epsilon_t = \begin{bmatrix} \epsilon_t \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

# Exemple of Multivariate time series: general bivariate case

Consider the bivariate case  $\mathbf{X}_t = (X_t(1), X_t(2))$ .


▷ IID case




# Exemple of Multivariate time series: general bivariate case

Consider the bivariate case  $\mathbf{X}_t = (X_t(1), X_t(2))$ .

▷ IID case



▷ Markov case:



- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 **Introducing dynamics**
  - What's wrong with i.i.d. models ?
  - Univariate models
  - Multivariate models
  - **Partially observed multivariate time series**
- 4 Stationary Time series
- 5 Weakly stationary time series

## Partially observed multivariate time series

- ▶ Exactly as in the IID case, adding **hidden variables** allows one to build a wild variety of models, while allowing the practitioner to provide intuitive interpretations of the model.

# Partially observed multivariate time series

- ▶ Exactly as in the IID case, adding **hidden variables** allows one to build a wild variety of models, while allowing the practitioner to provide intuitive interpretations of the model.
- ▶ The most widely used such time series model is the **linear state-space** model, or **dynamic linear model**, defined through two linear equations

$$\mathbf{X}_t = \Phi \mathbf{X}_{t-1} + \mathbf{U}_t \quad (\text{State Equation}) \quad (1a)$$

$$\mathbf{Y}_t = \mathbf{A} \mathbf{X}_t + \mathbf{V}_t \quad (\text{Observation Equation}), \quad (1b)$$

where  $(\mathbf{Y}_t)$  is the **observed** time series, and  $(\mathbf{X}_t)$  is the **hidden** time series (also called the state variables), and  $(\mathbf{U}_t)$  and  $(\mathbf{V}_t)$  are IID noise sequences.



# Partially observed multivariate time series

- ▶ Exactly as in the IID case, adding **hidden variables** allows one to build a wild variety of models, while allowing the practitioner to provide intuitive interpretations of the model.
- ▶ The most widely used such time series model is the **linear state-space** model, or **dynamic linear model**, defined through two linear equations

$$\mathbf{X}_t = \Phi \mathbf{X}_{t-1} + \mathbf{U}_t \quad (\text{State Equation}) \quad (1a)$$


$$\mathbf{Y}_t = \mathbf{A} \mathbf{X}_t + \mathbf{V}_t \quad (\text{Observation Equation}), \quad (1b)$$

where  $(\mathbf{Y}_t)$  is the **observed** time series, and  $(\mathbf{X}_t)$  is the **hidden** time series (also called the state variables), and  $(\mathbf{U}_t)$  and  $(\mathbf{V}_t)$  are IID noise sequences.

- ▶ This is a particular instance of the general class of the **partially observed Markov models**, where one has a bivariate Markov chain  $((\mathbf{X}_t, \mathbf{Y}_t))$ , where only the component  $(\mathbf{Y}_t)$  is observed.


# Examples of partially observed multivariate time series

▷ IID case




# Examples of partially observed multivariate time series

## ▷ IID case




## ▷ Partially observed Markov model: general case.



# Examples of partially observed multivariate time series (cont.)

▷ Hidden Markov model.




In this special case:

- ▷  $(X_t)$  alone is a Markov chain.

# Examples of partially observed multivariate time series (cont.)

## ▷ Hidden Markov model.




In this special case:

- ▷  $(X_t)$  alone is a Markov chain.
- ▷ Given  $(X_t)$ , the observations  $(Y_t)$  are **conditionally independent**.

# Examples of partially observed multivariate time series (cont.)

## ▶ Hidden Markov model.



In this special case:

- ▶  $(X_t)$  alone is a Markov chain.
- ▶ Given  $(X_t)$ , the observations  $(Y_t)$  are **conditionally independent**.
- ▶ Two highly popular special cases:
  - ▶ HMM with **finite** state space : when  $X_t$  takes values in  $\{1, \dots, K\}$ .
  - ▶ The **dynamic linear model**, see (1).

## Example : an HMM with two hidden states.




Figure: An HMM with two (supposedly) hidden states (red and black).

## Example : Noisy observations of an hidden AR(1) state variables.





Figure: Observations (black 'o') obtained by adding noise to a (supposedly) hidden AR(1) process (red lines).



# Observation driven models

- ▶ For most of the partially observed Markov models, there are no closed form formula for the **likelihood** and computational cost of  $L_n$  can be very high as  $n$  increases.
- ▶ **Observation driven models** stand as a popular exception. Their dependence structure takes the following form:



With the additional property that the **conditional distribution** of  $X_{k+1}$  given  $(X_k, Y_k)$  is **degenerate**.

## Exemple: GARCH(1,1) model

### GARCH(1,1) model

For parameter  $\theta = (a, b, c) \in (0, \infty)^3$ ,  $(Y_t)$  satisfies the GARCH(1,1) equation

$$\sigma_t^2 = a + bY_{t-1} + c\sigma_{t-1}^2 \quad (2a)$$

$$Y_t = \sigma_t \epsilon_t, \quad (2b)$$

where  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

Moreover it is assumed that  $(\sigma_t)$  is **non-anticipative** solution, in the sense that, for all  $t \in \mathbb{Z}$ ,  $\sigma_t$  only depends on  $(\epsilon_s)_{s < t}$

# Exemple: GARCH(1,1) model

## GARCH(1,1) model

For parameter  $\theta = (a, b, c) \in (0, \infty)^3$ ,  $(Y_t)$  satisfies the GARCH(1,1) equation

$$\sigma_t^2 = a + bY_{t-1} + c\sigma_{t-1}^2 \quad (2a)$$

$$Y_t = \sigma_t \epsilon_t, \quad (2b)$$

where  $(\epsilon_t)_{t \in \mathbb{Z}}$  i.i.d.  $\sim \mathcal{N}(0, 1)$ .

Moreover it is assumed that  $(\sigma_t)$  is **non-anticipative** solution, in the sense that, for all  $t \in \mathbb{Z}$ ,  $\sigma_t$  only depends on  $(\epsilon_s)_{s < t}$

The fact that  $(\sigma_t)$  is **non-anticipative** ensures that, for all  $t \in \mathbb{Z}$ , given  $(\epsilon_s)_{s < t}$ , the conditional distribution of  $Y_t$  is  $\mathcal{N}(0, \sigma_t)$ .

## Exemple: GARCH(1,1) model, likelihood

Iterating (2a) with a given  $\theta$ , for all  $k = 2, \dots, n$ , one can express  $\sigma_k^2$  as a **deterministic** function of  $Y_{1:k-1}$  and  $\sigma_1^2$ , say

$$\sigma_k^2 = \psi^\theta < Y_{1:k-1} > (\sigma_1^2). \quad (3)$$

Note that  $\psi^\theta < Y_{1:k-1} > (\sigma_1^2)$  is easy to compute iteratively.

## Exemple: GARCH(1,1) model, likelihood

Iterating (2a) with a given  $\theta$ , for all  $k = 2, \dots, n$ , one can express  $\sigma_k^2$  as a **deterministic** function of  $Y_{1:k-1}$  and  $\sigma_1^2$ , say

$$\sigma_k^2 = \psi^\theta < Y_{1:k-1} > (\sigma_1^2). \quad (3)$$

Note that  $\psi^\theta < Y_{1:k-1} > (\sigma_1^2)$  is easy to compute iteratively.

Using (3) and (2b), the (conditional) negated log likelihood (given  $\sigma_1^2 = s_1^2$  and  $Y_1$  for some arbitrary  $s_1^2$ ) is given by

$$-L_n(\theta) = \frac{1}{2} \sum_{k=2}^n \left( \log \left( 2\pi \psi^\theta < Y_{1:k-1} > (s_1^2) \right) + \frac{Y_k^2}{\psi^\theta < Y_{1:k-1} > (s_1^2)} \right),$$

which can be minimized in  $\theta = (a, b, c)$  using a **gradient descent algorithm**.

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series**
  - The statistical approach
  - Classical steps of statistical inference
  - Stationary and ergodic models
- 5 Weakly stationary time series

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series**
  - The statistical approach
    - Classical steps of statistical inference
    - Stationary and ergodic models
- 5 Weakly stationary time series

# Basic (important) definitions

## Definition : Data set

A data set is a collection of values, say  $X_{1:n} = X_1, \dots, X_n$ . Time series data sets are usually **sampled** from recorded measurements.



# Basic (important) definitions

## Definition : Data set

A data set is a collection of values, say  $X_{1:n} = X_1, \dots, X_n$ . Time series data sets are usually **sampled** from recorded measurements.

## Definition : Model

A **model** is a collection of probability distributions. The data set is assumed to be distributed according to one of them.

# Basic (important) definitions

## Definition : Data set

A data set is a collection of values, say  $X_{1:n} = X_1, \dots, X_n$ . Time series data sets are usually **sampled** from recorded measurements.

## Definition : Model

A **model** is a collection of probability distributions. The data set is assumed to be distributed according to one of them.

## Definition : Statistic

A **statistic** is any value which can be computed from the data.

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

For instance :

- ▶ a date in years and frequency= 12 corresponds to **monthly data**,

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

For instance :

- ▶ a date in years and frequency= 12 corresponds to **monthly data**,
- ▶ a date in years and frequency= 4 corresponds to **quarterly data**,

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

For instance :

- ▶ a date in years and frequency= 12 corresponds to **monthly data**,
- ▶ a date in years and frequency= 4 corresponds to **quarterly data**,
- ▶ a date in days and frequency= 1 corresponds to **daily data**,
- ▶  $\vdots$



# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

For instance :

- ▶ a date in years and frequency= 12 corresponds to **monthly data**,
- ▶ a date in years and frequency= 4 corresponds to **quarterly data**,
- ▶ a date in days and frequency= 1 corresponds to **daily data**,
- ▶  $\vdots$
- ▶ Remarks :
  - ▶ There may be **missing values** (usually expressed as 'NA')

# Raw data set

A time series  $X_1, \dots, X_n$  is usually presented as

- ▶ a list of real values  $X_1, \dots, X_n$  in a data or spreadsheet file,
- ▶ a corresponding list of dates (days, years, seconds...)
- ▶ or, **equivalently**, a starting date (in some unit), and a frequency.

For instance :

- ▶ a date in years and frequency= 12 corresponds to **monthly data**,
- ▶ a date in years and frequency= 4 corresponds to **quarterly data**,
- ▶ a date in days and frequency= 1 corresponds to **daily data**,
- ▶  $\vdots$
- ▶ Remarks :
  - ▶ There may be **missing values** (usually expressed as 'NA')
  - ▶ In the case of **multivariate time series**, each variable usually corresponds to a column (so each row corresponds to a date).

## Example : US GNP data set

```
# Title:           Gross National Product
# Source:          U.S. Department of Commerce
# Frequency:       Quarterly
```

```
DATE,VALUE
```

```
1947-01-01,238.1
```

```
1947-04-01,241.5
```

```
1947-07-01,245.6
```

```
1947-10-01,255.6
```

```
1948-01-01,261.7
```

```
1948-04-01,268.7
```

```
1948-07-01,275.3
```

```
1948-10-01,276.6
```

```
1949-01-01,271.3
```

```
1949-04-01,267.5
```

```
1949-07-01,268.9
```

```
⋮
```

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series**
  - The statistical approach
  - Classical steps of statistical inference**
  - Stationary and ergodic models
- 5 Weakly stationary time series

## First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .

## First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:

## First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:
  - ▶ a **polynomial trend**,

## First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:
  - ▶ a **polynomial trend**,
  - ▶ a **seasonal trend**.



# First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:
  - ▶ a **polynomial trend**,
  - ▶ a **seasonal trend**.
- ▶ Then  $X = D + Y$  where  $D$  belongs to a **finite dimensional** space  $V$  and  $Y$  is a centered random process.

# First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:
  - ▶ a **polynomial trend**,
  - ▶ a **seasonal trend**.
- ▶ Then  $X = D + Y$  where  $D$  belongs to a **finite dimensional** space  $V$  and  $Y$  is a centered random process.
- ▶ Trends can be estimated or removed by
  - ▶ Fitting the trend using **least squares**,

$$\hat{D} = \operatorname{argmin}_{d \in V} \sum_t |X_t - d_t|^2 .$$

# First step : remove an additive trend

- ▶ Consider a time series  $X_1, \dots, X_n$ .
- ▶ It often includes an **additive trend**, for instance:
  - ▶ a **polynomial trend**,
  - ▶ a **seasonal trend**.
- ▶ Then  $X = D + Y$  where  $D$  belongs to a **finite dimensional** space  $V$  and  $Y$  is a centered random process.
- ▶ Trends can be estimated or removed by
  - ▶ Fitting the trend using **least squares**,

$$\hat{D} = \underset{d \in V}{\operatorname{argmin}} \sum_t |X_t - d_t|^2 .$$

- ▶ Or applying a well chosen filter  $F_\psi$ , such that  $F_\psi(D) = 0$  and thus

$$F_\psi(X) = F_\psi(Y) .$$

# R code example: Johnson and Johnson trend adjustment

[trend-adjustment.html](#)

## Second step : choose a stochastic model on the random part

In time series analysis, one is interested in modeling the **time dependence** in the **trend adjusted** data  $Y_1, \dots, Y_n$ .

## Second step : choose a stochastic model on the random part

In time series analysis, one is interested in modeling the **time dependence** in the **trend adjusted** data  $Y_1, \dots, Y_n$ .

This can be done by using

- ▶ a **parametric model**.

## Second step : choose a stochastic model on the random part

In time series analysis, one is interested in modeling the **time dependence** in the **trend adjusted** data  $Y_1, \dots, Y_n$ .

This can be done by using

▷ a **parametric model**.

### Example

$Y_1, \dots, Y_n$  is the sample of a Gaussian ARMA( $p, q$ ) model with (unknown) **parameter**  $\vartheta = (\theta_1, \dots, \theta_q, \phi_1, \dots, \phi_p, \sigma^2)$ .

## Second step : choose a stochastic model on the random part

In time series analysis, one is interested in modeling the **time dependence** in the **trend adjusted** data  $Y_1, \dots, Y_n$ .

This can be done by using

- ▷ a **parametric model**.

### Example

$Y_1, \dots, Y_n$  is the sample of a Gaussian ARMA( $p, q$ ) model with (unknown) **parameter**  $\vartheta = (\theta_1, \dots, \theta_q, \phi_1, \dots, \phi_p, \sigma^2)$ .

- ▷ a **non-parametric model**.



## Second step : choose a stochastic model on the random part

In time series analysis, one is interested in modeling the **time dependence** in the **trend adjusted** data  $Y_1, \dots, Y_n$ .

This can be done by using

- ▷ a **parametric model**.

### Example

$Y_1, \dots, Y_n$  is the sample of a Gaussian ARMA( $p, q$ ) model with (unknown) **parameter**  $\vartheta = (\theta_1, \dots, \theta_q, \phi_1, \dots, \phi_p, \sigma^2)$ .

- ▷ a **non-parametric model**.

### Example

$Y_1, \dots, Y_n$  is the sample of a centered stationary Gaussian process with (unknown) autocovariance  $\gamma$  (or spectral density  $f$ ).

## Third step : estimate parameters, test hypotheses

Once a model is fixed for  $Y_1, \dots, Y_n$ , it can be used to

- ▶ **Estimate** a parameter of the model such as  $\vartheta$ ,  $\gamma(t)$ ,  $\sigma^2$ ,  $f$ , ...

## Third step : estimate parameters, test hypotheses

Once a model is fixed for  $Y_1, \dots, Y_n$ , it can be used to

▷ **Estimate** a parameter of the model such as  $\vartheta$ ,  $\gamma(t)$ ,  $\sigma^2$ ,  $f$ , ...

→ Define an **estimator**, say  $\hat{\vartheta}_n$ , which is a statistic based on the sample  $Y_1, \dots, Y_n$ .

## Third step : estimate parameters, test hypotheses

Once a model is fixed for  $Y_1, \dots, Y_n$ , it can be used to

▷ **Estimate** a parameter of the model such as  $\vartheta$ ,  $\gamma(t)$ ,  $\sigma^2$ ,  $f$ , ...

→ Define an **estimator**, say  $\hat{\vartheta}_n$ , which is a statistic based on the sample  $Y_1, \dots, Y_n$ .

▷ **Test** hypotheses, for instance

$$H_0 = \{Y \text{ is white noise}\} \quad \text{against} \quad H_1 = \{Y \text{ is ARMA}(p, q)\}$$

## Third step : estimate parameters, test hypotheses

Once a model is fixed for  $Y_1, \dots, Y_n$ , it can be used to

▷ **Estimate** a parameter of the model such as  $\vartheta$ ,  $\gamma(t)$ ,  $\sigma^2$ ,  $f$ , ...

→ Define an **estimator**, say  $\hat{\vartheta}_n$ , which is a statistic based on the sample  $Y_1, \dots, Y_n$ .

▷ **Test** hypotheses, for instance

$$H_0 = \{Y \text{ is white noise}\} \quad \text{against} \quad H_1 = \{Y \text{ is ARMA}(p, q)\}$$

→ Define a **statistical test**, say

$$\delta = \begin{cases} 1 & \text{if } T_n > t_n, \\ 0 & \text{otherwise,} \end{cases}$$

where  $T_n$  is a **statistic** based on the sample  $Y_1, \dots, Y_n$  and  $t_n$  is a **threshold**.

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series**
  - The statistical approach
  - Classical steps of statistical inference
  - **Stationary and ergodic models**
- 5 Weakly stationary time series

# Stationary and ergodic models

- ▶ We see  $Y_1, \dots, Y_n$  as a **finite sample** of a **stochastic process**  $(Y_t)_{t \in T}$ , with  $T = \mathbb{N}$  or  $\mathbb{Z}$ , distributed according to a **statistical model**.

# Stationary and ergodic models

- ▶ We see  $Y_1, \dots, Y_n$  as a finite sample of a stochastic process  $(Y_t)_{t \in T}$ , with  $T = \mathbb{N}$  or  $\mathbb{Z}$ , distributed according to a statistical model.
- ▶ It is crucial to work with stationary and ergodic models.



# Stationary and ergodic models

- ▶ We see  $Y_1, \dots, Y_n$  as a **finite sample** of a **stochastic process**  $(Y_t)_{t \in T}$ , with  $T = \mathbb{N}$  or  $\mathbb{Z}$ , distributed according to a **statistical model**.
- ▶ It is crucial to work with **stationary** and **ergodic** models.
- ▶ **Stationary** means that the model is **shift invariant**: for all  $n \geq 1$ , and all  $t_1, \dots, t_n \in T$ , we have

$$(X_{t_1}, \dots, X_{t_n}) \stackrel{d}{=} (X_{t_1+1}, \dots, X_{t_n+1}) .$$

# Stationary and ergodic models

- ▶ We see  $Y_1, \dots, Y_n$  as a **finite sample** of a **stochastic process**  $(Y_t)_{t \in T}$ , with  $T = \mathbb{N}$  or  $\mathbb{Z}$ , distributed according to a **statistical model**.
- ▶ It is crucial to work with **stationary** and **ergodic** models.
- ▶ **Stationary** means that the model is **shift invariant**: for all  $n \geq 1$ , and all  $t_1, \dots, t_n \in T$ , we have

$$(X_{t_1}, \dots, X_{t_n}) \stackrel{d}{=} (X_{t_1+1}, \dots, X_{t_n+1}) .$$

- ▶ **Ergodic** means that observing one **path**  $(Y_t)_{t \in T}$  allows one to recover the distribution **entirely**.

# Examples

- ▶ An IID process is stationary and ergodic;

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:
  - ▶ removing trends,

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:
  - ▶ removing trends,
  - ▶ linear filters such as taking the increments (random walks),

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:
  - ▶ removing trends,
  - ▶ linear filters such as taking the increments (random walks),
  - ▶ non-linear filters such as taking the logs.



# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:
  - ▶ removing trends,
  - ▶ linear filters such as taking the increments (random walks),
  - ▶ non-linear filters such as taking the logs.
- ▶ A sequence of variables  $(Y_t)_{t \in \mathbb{Z}}$  that is constant, i.e.  $Y_t = Y_0$  for all  $t$ , is stationary but is not ergodic;

# Examples

- ▶ An IID process is stationary and ergodic;
- ▶ A sequence of independent variables that are not identically distributed is not stationary;
- ▶ some models are not stationary but can be made so through some basic transformations:
  - ▶ removing trends,
  - ▶ linear filters such as taking the increments (random walks),
  - ▶ non-linear filters such as taking the logs.
- ▶ A sequence of variables  $(Y_t)_{t \in \mathbb{Z}}$  that is constant, i.e.  $Y_t = Y_0$  for all  $t$ , is stationary but is not ergodic;
- ▶ A Markov chain on a finite state space can be made stationary by choosing the initial state adequately. If it is irreducible, then it is ergodic.

# R code example: dependent data

[non-iid-data.html](#)

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series**
  - $L^2$  processes
  - Weak stationarity
  - Spectral measure
  - Empirical estimation

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series**
  - $L^2$  processes
  - Weak stationarity
  - Spectral measure
  - Empirical estimation

# $L^2$ space

We denote

$$L^2(\Omega, \mathcal{F}, \mathbb{P}) = \{X \text{ } \mathbb{C}\text{-valued r.v. such that } \mathbb{E}[|X|^2] < \infty\} .$$

$(L^2, \langle, \rangle)$  is a Hilbert space with

$$\langle X, Y \rangle = \mathbb{E}[X\overline{Y}] .$$

## Definition : $L^2$ Processes

The process  $X = (X_t)_{t \in T}$  defined on  $(\Omega, \mathcal{F}, \mathbb{P})$  with values in  $\mathbb{C}$  is an  $L^2$  process if  $X_t \in L^2(\Omega, \mathcal{F}, \mathbb{P})$  for all  $t \in T$ .

# Mean and covariance functions

Let  $X = (X_t)_{t \in T}$  be an  $L^2$  process.

# Mean and covariance functions

Let  $X = (X_t)_{t \in T}$  be an  $L^2$  process.

▷ Its **mean function** is defined by  $\mu(t) = \mathbb{E}[X_t]$ ,



# Mean and covariance functions

Let  $X = (X_t)_{t \in T}$  be an  $L^2$  process.

- ▷ Its **mean function** is defined by  $\mu(t) = \mathbb{E}[X_t]$ ,
- ▷ Its **covariance function** is defined by

$$\gamma(s, t) = \text{cov}(X_s, X_t) = \mathbb{E}[X_s \overline{X_t}] - \mathbb{E}[X_s] \mathbb{E}[\overline{X_t}] .$$

# Mean and covariance functions

Let  $X = (X_t)_{t \in T}$  be an  $L^2$  process.

- ▷ Its **mean function** is defined by  $\mu(t) = \mathbb{E}[X_t]$ ,
- ▷ Its **covariance function** is defined by

$$\gamma(s, t) = \text{cov}(X_s, X_t) = \mathbb{E}[X_s \overline{X_t}] - \mathbb{E}[X_s] \mathbb{E}[\overline{X_t}] .$$

## Hermitian symmetry, non-negative definiteness

For all finite subset  $I \subset T$ ,  $\Gamma_I = \text{Cov}([X(t)]_{t \in I}) = [\gamma(s, t)]_{s, t \in I}$  is a **hermitian non-negative definite** matrix.

# Examples

- ▷  $L^2$  independent random variables  $(X_t)_{t \in \mathbb{Z}}$  have mean  $\mu(t) = \mathbb{E}[X_t]$  and covariance

$$\gamma(s, t) = \begin{cases} \text{var}(X_t) & \text{if } s = t, \\ 0 & \text{otherwise.} \end{cases}$$

# Examples

- ▷  $L^2$  independent random variables  $(X_t)_{t \in \mathbb{Z}}$  have mean  $\mu(t) = \mathbb{E}[X_t]$  and covariance

$$\gamma(s, t) = \begin{cases} \text{var}(X_t) & \text{if } s = t, \\ 0 & \text{otherwise.} \end{cases}$$

- ▷ A Gaussian process is an  $L^2$  process whose law is entirely determined by its mean and covariance functions: for all  $I = \{t_1, \dots, t_n\}$ ,

$$(X_s)_{s \in I} \sim \mathcal{N}((\mu_s)_{s \in I}, \Gamma_I) .$$

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series**
  - $L^2$  processes
  - **Weak stationarity**
  - Spectral measure
  - Empirical estimation

# Weakly stationary processes

- ▶ Let  $T = \mathbb{Z}$ . Let  $X$  be an  $L^2$  strictly stationary process with mean function  $\mu$  and covariance function  $\gamma$ .

# Weakly stationary processes

- ▷ Let  $T = \mathbb{Z}$ . Let  $X$  be an  $L^2$  strictly stationary process with mean function  $\mu$  and covariance function  $\gamma$ .
- ▷ Then  $\mu(t) = \mu(0)$  and  $\gamma(s, t) = \gamma(s - t, 0)$  for all  $s, t \in T$ .

# Weakly stationary processes

- ▶ Let  $T = \mathbb{Z}$ . Let  $X$  be an  $L^2$  strictly stationary process with mean function  $\mu$  and covariance function  $\gamma$ .
- ▶ Then  $\mu(t) = \mu(0)$  and  $\gamma(s, t) = \gamma(s - t, 0)$  for all  $s, t \in T$ .

## Definition : Weak stationarity

We say that a random process  $X$  is **weakly stationary** with **mean**  $\mu \in \mathbb{C}$  and **autocovariance function**  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$  if it is  $L^2$  with mean function  $t \mapsto \mu$  and covariance function  $(s, t) \mapsto \gamma(s - t)$ .



# Weakly stationary processes

- ▶ Let  $T = \mathbb{Z}$ . Let  $X$  be an  $L^2$  strictly stationary process with mean function  $\mu$  and covariance function  $\gamma$ .
- ▶ Then  $\mu(t) = \mu(0)$  and  $\gamma(s, t) = \gamma(s - t, 0)$  for all  $s, t \in T$ .

## Definition : Weak stationarity

We say that a random process  $X$  is **weakly stationary** with **mean**  $\mu \in \mathbb{C}$  and **autocovariance function**  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$  if it is  $L^2$  with mean function  $t \mapsto \mu$  and covariance function  $(s, t) \mapsto \gamma(s - t)$ .


- ▶ The **autocorrelation function** is then defined (when  $\gamma(0) > 0$ ) by

$$\rho(h) = \frac{\gamma(h)}{\gamma(0)} \in [-1, 1] .$$

# Autocorrelation=slope of regression line

We have, for all  $t \in \mathbb{Z}$  and  $h = 1, 2, \dots$ ,

$$X_t = \text{Constant} + \rho(h)X_{t-h} + \epsilon_{t,h} \quad \text{with} \quad \epsilon_{t,h} \perp \text{Span}(1, X_{t-h}) .$$



# Partial Autocorrelation

▷ We can also write, for all  $t \in \mathbb{Z}$  and  $h = 1, 2, \dots$ ,

$$X_t = \text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} + \epsilon_{t,h}$$

where

$$\epsilon_{t,h} \perp \text{Span}(1, X_{t-1}, \dots, X_{t-h}) .$$

# Partial Autocorrelation

- ▷ We can also write, for all  $t \in \mathbb{Z}$  and  $h = 1, 2, \dots$ ,

$$X_t = \text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} + \epsilon_{t,h}$$

where

$$\epsilon_{t,h} \perp \text{Span}(1, X_{t-1}, \dots, X_{t-h}) .$$

- ▷ In the following, we plot successively
- ▷  $X_t$  as a function of  $X_{t-h}$ , compared to the regression line;

# Partial Autocorrelation

- ▷ We can also write, for all  $t \in \mathbb{Z}$  and  $h = 1, 2, \dots$ ,

$$X_t = \text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} + \epsilon_{t,h}$$

where

$$\epsilon_{t,h} \perp \text{Span}(1, X_{t-1}, \dots, X_{t-h}) .$$

- ▷ In the following, we plot successively
- ▷  $X_t$  as a function of  $X_{t-h}$ , compared to the regression line;
  - ▷  $X_t$  as a function of  $X_{t-h}$ , compared to the predictor

$$\text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} .$$

# Partial Autocorrelation

- ▷ We can also write, for all  $t \in \mathbb{Z}$  and  $h = 1, 2, \dots$ ,

$$X_t = \text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} + \epsilon_{t,h}$$

where


$$\epsilon_{t,h} \perp \text{Span}(1, X_{t-1}, \dots, X_{t-h}) .$$

- ▷ In the following, we plot successively
- ▷  $X_t$  as a function of  $X_{t-h}$ , compared to the regression line;
  - ▷  $X_t$  as a function of  $X_{t-h}$ , compared to the predictor

$$\text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} + \kappa(h) X_{t-h} .$$

- ▷  $X_t - \left( \text{Constant} + \sum_{k=1}^{h-1} \phi_k X_{t-k} \right)$  as a function of  $X_{t-h}$ ,  
compared to the regression line  $X_{t-h} \mapsto \kappa(h) X_{t-h}$ .

# Partial Autocorrelation=slope of partial regression



# Examples

- ▶ An  $L^2$  strictly stationary process is weakly stationary.



# Examples

- ▷ An  $L^2$  strictly stationary process is weakly stationary.
- ▷ The constant  $L^2$  process has **constant autocovariance function**.

# Examples

- ▶ An  $L^2$  strictly stationary process is weakly stationary.
- ▶ The constant  $L^2$  process has **constant autocovariance function**.

## Strong and weak white noise

- ▶ A sequence of  $L^2$  i.i.d. random variables is called a **strong white noise**, denoted by  $X \sim \text{IID}(\mu, \sigma^2)$ .

# Examples

- ▶ An  $L^2$  strictly stationary process is weakly stationary.
- ▶ The constant  $L^2$  process has **constant autocovariance function**.

## Strong and weak white noise

- ▶ A sequence of  $L^2$  i.i.d. random variables is called a **strong white noise**, denoted by  $X \sim \text{IID}(\mu, \sigma^2)$ .
- ▶ An  $L^2$  process  $X$  with constant mean  $\mu$  and **constant diagonal covariance function** equal to  $\sigma^2$  is called a **weak white noise**. It is denoted by  $X \sim \text{WN}(\mu, \sigma^2)$ . (It does not have to be i.i.d.)

## Examples based on stationarity preserving linear filters

- ▶ Let  $X$  be weakly stationary with mean  $\mu$  and autocovariance  $\gamma$ .

# Examples based on stationarity preserving linear filters

- ▷ Let  $X$  be weakly stationary with mean  $\mu$  and autocovariance  $\gamma$ .
- ▷ Let  $\psi \in \ell^1$ . We define  $Y = F_\psi(X)$  by

$$Y_t = \sum_k \psi_k X_{t-k}, \quad t \in \mathbb{Z}.$$

# Examples based on stationarity preserving linear filters

- ▶ Let  $X$  be weakly stationary with mean  $\mu$  and autocovariance  $\gamma$ .
- ▶ Let  $\psi \in \ell^1$ . We define  $Y = F_\psi(X)$  by

$$Y_t = \sum_k \psi_k X_{t-k}, \quad t \in \mathbb{Z}.$$

- ▶ Then  $Y$  is weakly stationary with mean  $\mu'$  and autocovariance  $\gamma'$  given by

$$\begin{aligned} \mu' &= \mu \sum_k \psi_k \\ \gamma'(\tau) &= \sum_{\ell, k} \psi_k \overline{\psi_\ell} \gamma(\tau + \ell - k) \end{aligned} \tag{4}$$

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series**
  - $L^2$  processes
  - Weak stationarity
  - Spectral measure**
  - Empirical estimation

# Spectral measure

- ▶ Given a function  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$ , does there exist a weakly stationary process  $(X_t)_{t \in \mathbb{Z}}$  with autocovariance  $\gamma$ ?



# Spectral measure

- ▶ Given a function  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$ , does there exist a weakly stationary process  $(X_t)_{t \in \mathbb{Z}}$  with autocovariance  $\gamma$ ?
- ▶ A necessary and sufficient condition is that  $\Gamma_I = [\gamma(s - t)]_{s, t \in I}$  is a hermitian non-negative definite matrix for all finite subset  $I \subset \mathbb{Z}$ .

# Spectral measure

- ▶ Given a function  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$ , does there exist a weakly stationary process  $(X_t)_{t \in \mathbb{Z}}$  with autocovariance  $\gamma$ ?
- ▶ A necessary and sufficient condition is that  $\Gamma_I = [\gamma(s - t)]_{s, t \in I}$  is a hermitian non-negative definite matrix for all finite subset  $I \subset \mathbb{Z}$ .
- ▶ We say that  $\gamma$  is hermitian symmetric and non-negative definite.

# Spectral measure

- ▶ Given a function  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$ , does there exist a weakly stationary process  $(X_t)_{t \in \mathbb{Z}}$  with autocovariance  $\gamma$ ?
- ▶ A necessary and sufficient condition is that  $\Gamma_I = [\gamma(s - t)]_{s, t \in I}$  is a hermitian non-negative definite matrix for all finite subset  $I \subset \mathbb{Z}$ .
- ▶ We say that  $\gamma$  is hermitian symmetric and non-negative definite.

## Herglotz Theorem

Let  $\gamma : \mathbb{Z} \rightarrow \mathbb{C}$ . Then the two following assertions are equivalent:

- (i)  $\gamma$  is hermitian symmetric and non-negative definite.
- (ii) There exists a finite non-negative measure  $\nu$  on  $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$  such that,

$$\text{for all } t \in \mathbb{Z}, \quad \gamma(t) = \int_{\mathbb{T}} e^{i\lambda t} \nu(d\lambda). \quad (5)$$

When these two assertions hold,  $\nu$  is uniquely defined by (5).

# Spectral density

If moreover  $\gamma \in \ell^1(\mathbb{Z})$ , these assertions are equivalent to

$$f(\lambda) := \frac{1}{2\pi} \sum_{t \in \mathbb{Z}} e^{-i\lambda t} \gamma(t) \geq 0 \text{ for all } \lambda \in \mathbb{R},$$

and  $\nu$  has density  $f$  (that is,  $\nu(d\lambda) = f(\lambda)d\lambda$ ).

# Spectral density

If moreover  $\gamma \in \ell^1(\mathbb{Z})$ , these assertions are equivalent to

$$f(\lambda) := \frac{1}{2\pi} \sum_{t \in \mathbb{Z}} e^{-i\lambda t} \gamma(t) \geq 0 \text{ for all } \lambda \in \mathbb{R},$$

and  $\nu$  has density  $f$  (that is,  $\nu(d\lambda) = f(\lambda)d\lambda$ ).

## Definition : spectral measure and spectral density

If  $\gamma$  is the autocovariance of a weakly stationary process  $X$ , the corresponding measure  $\nu$  is called the **spectral measure** of  $X$ . Whenever the spectral measure  $\nu$  admits a density  $f$ , it is called the **spectral density function**.

## Examples

▷ Let  $X \sim \text{WN}(\mu, \sigma^2)$ . Then  $f(\lambda) = \frac{\sigma^2}{2\pi}$ .

## Examples

- ▷ Let  $X \sim \text{WN}(\mu, \sigma^2)$ . Then  $f(\lambda) = \frac{\sigma^2}{2\pi}$ .
- ▷ Let  $X$  be a weakly stationary process with covariance function  $\gamma$ /spectral measure  $\nu$ . Define

$$Y_t = \sum_k \psi_k X_{t-k}$$

for a sequence  $\psi \in \ell^1$ .

## Examples

- ▶ Let  $X \sim \text{WN}(\mu, \sigma^2)$ . Then  $f(\lambda) = \frac{\sigma^2}{2\pi}$ .
- ▶ Let  $X$  be a weakly stationary process with covariance function  $\gamma$ /spectral measure  $\nu$ . Define

$$Y_t = \sum_k \psi_k X_{t-k}$$

for a sequence  $\psi \in \ell^1$ .

- ▶ Recall that  $Y$  is a weakly stationary process with covariance function

$$\gamma'(\tau) = \sum_{\ell, k} \psi_k \overline{\psi_\ell} \gamma(\tau + \ell - k) .$$



## Examples

- ▶ Let  $X \sim \text{WN}(\mu, \sigma^2)$ . Then  $f(\lambda) = \frac{\sigma^2}{2\pi}$ .
- ▶ Let  $X$  be a weakly stationary process with covariance function  $\gamma$ /spectral measure  $\nu$ . Define

$$Y_t = \sum_k \psi_k X_{t-k}$$

for a sequence  $\psi \in \ell^1$ .

- ▶ Recall that  $Y$  is a weakly stationary process with covariance function

$$\gamma'(\tau) = \sum_{\ell, k} \psi_k \overline{\psi_\ell} \gamma(\tau + \ell - k) .$$

- ▶ Then  $Y$  is a weakly stationary process with spectral measure  $\nu'$  having density  $\lambda \mapsto \left| \sum_k \psi_k e^{-i\lambda k} \right|^2$  with respect to  $\nu$ ,

$$\nu'(d\lambda) = \left| \sum_k \psi_k e^{-i\lambda k} \right|^2 \nu(d\lambda) .$$

## A special one : the harmonic process

Let  $(A_k)_{1 \leq k \leq N}$  be  $N$  real valued  $L^2$  random variables. Denote  $\sigma_k^2 = \mathbb{E}[A_k^2]$ . Let  $(\Phi_k)_{1 \leq k \leq N}$  be  $N$  i.i.d. random variables with a uniform distribution on  $[0, 2\pi]$ , and independent of  $(A_k)_{1 \leq k \leq N}$ . Define

$$X_t = \sum_{k=1}^N A_k \cos(\lambda_k t + \Phi_k), \quad (6)$$

where  $(\lambda_k)_{1 \leq k \leq N} \in [-\pi, \pi]$  are  $N$  frequencies. The process  $(X_t)$  is called a **harmonic process**. It satisfies  $\mathbb{E}[X_t] = 0$  and, for all  $s, t \in \mathbb{Z}$ ,

$$\mathbb{E}[X_s X_t] = \frac{1}{2} \sum_{k=1}^N \sigma_k^2 \cos(\lambda_k(s - t)).$$

Hence  $X$  is weakly stationary with autocovariance

$$\gamma(t) = \frac{1}{2} \sum_{k=1}^N \sigma_k^2 \cos(\lambda_k t) = \int_{\mathbb{T}} e^{i\lambda t} \left( \frac{1}{4} \sum_{k=1}^N \sigma_k^2 (\delta_{-\lambda_k}(d\lambda) + \delta_{\lambda_k}(d\lambda)) \right).$$

- 1 Example of time series
- 2 Reminders: i.i.d. models
- 3 Introducing dynamics
- 4 Stationary Time series
- 5 Weakly stationary time series**
  - $L^2$  processes
  - Weak stationarity
  - Spectral measure
  - Empirical estimation

# Empirical estimates

- ▶ Suppose you want to estimate the mean and the autocovariance from a sample  $X_1, \dots, X_n$ .

# Empirical estimates

- ▷ Suppose you want to estimate the mean and the autocovariance from a sample  $X_1, \dots, X_n$ .
- ▷ Define the **empirical mean** as

$$\hat{\mu}_n = \frac{1}{n} \sum_{k=1}^n X_k ,$$

# Empirical estimates

- ▷ Suppose you want to estimate the mean and the autocovariance from a sample  $X_1, \dots, X_n$ .
- ▷ Define the **empirical mean** as

$$\hat{\mu}_n = \frac{1}{n} \sum_{k=1}^n X_k ,$$

- ▷ Define the **empirical autocovariance** and **autocorrelation** functions as

$$\hat{\gamma}_n(h) = \frac{1}{n} \sum_{k=1}^{n-|h|} (X_k - \hat{\mu}_n)(X_{k+|h|} - \hat{\mu}_n) \quad \text{and}$$
$$\hat{\rho}_n(h) = \frac{\hat{\gamma}_n(h)}{\hat{\gamma}_n(0)} .$$

# Periodogram

- ▶ The previous formula only work for  $h = -n + 1, \dots, n - 1$ .

# Periodogram

- ▶ The previous formula only work for  $h = -n + 1, \dots, n - 1$ .
- ▶ Define  $\hat{\gamma}_n(h) = \hat{\rho}_n(h) = 0$  for all  $|h| \geq n$ .



# Periodogram

- ▶ The previous formula only work for  $h = -n + 1, \dots, n - 1$ .
- ▶ Define  $\hat{\gamma}_n(h) = \hat{\rho}_n(h) = 0$  for all  $|h| \geq n$ .
- ▶ Now  $\hat{\gamma}_n$  is defined on  $\mathbb{Z}$  and satisfies

$$\hat{\gamma}_n(h) = \int_{-\pi}^{\pi} e^{i\lambda h} I_n(\lambda) d\lambda ,$$

where  $I_n$  is called the (raw) periodogram and is defined by

$$I_n(\lambda) = \frac{1}{2\pi n} \left| \sum_{k=1}^n (X_k - \hat{\mu}_n) e^{-i\lambda k} \right|^2 .$$

# Periodogram

- ▶ The previous formula only work for  $h = -n + 1, \dots, n - 1$ .
- ▶ Define  $\hat{\gamma}_n(h) = \hat{\rho}_n(h) = 0$  for all  $|h| \geq n$ .
- ▶ Now  $\hat{\gamma}_n$  is defined on  $\mathbb{Z}$  and satisfies

$$\hat{\gamma}_n(h) = \int_{-\pi}^{\pi} e^{i\lambda h} I_n(\lambda) \, d\lambda ,$$

where  $I_n$  is called the (raw) **periodogram** and is defined by

$$I_n(\lambda) = \frac{1}{2\pi n} \left| \sum_{k=1}^n (X_k - \hat{\mu}_n) e^{-i\lambda k} \right|^2 .$$

- ▶  $I_n(\lambda)$  can be seen as a (bad) estimator of the spectral density  $f(\lambda)$ .