

# 2 Unit Bridging Course – Day 4

The derivative of a function

Emi Tanaka



# Derivative of a function

A derivative is concerned with how one quantity changes with respect to another quantity, in other words a rate of change.

The derivative of the function  $y = f(x)$  with respect to  $x$  will show us how  $y$  changes as the value  $x$  changes. It gives us the slope, or **gradient** of the function.

The derivative of  $y = f(x)$  with respect to  $x$  is represented by the following notations:

$$f'(x), \quad \frac{d}{dx}(f(x)), \quad \frac{df}{dx} \quad \text{or} \quad \frac{dy}{dx}.$$

The process of finding the derivative is called *differentiation*.

# Derivative of a function

A derivative is concerned with how one quantity changes with respect to another quantity, in other words a rate of change.

The derivative of the function  $y = f(x)$  with respect to  $x$  will show us how  $y$  changes as the value  $x$  changes. It gives us the slope, or **gradient** of the function.

The derivative of  $y = f(x)$  with respect to  $x$  is represented by the following notations:

$$f'(x), \quad \frac{d}{dx}(f(x)), \quad \frac{df}{dx} \quad \text{or} \quad \frac{dy}{dx}.$$

The process of finding the derivative is called *differentiation*.

# Derivative of a function

A derivative is concerned with how one quantity changes with respect to another quantity, in other words a rate of change.

The derivative of the function  $y = f(x)$  with respect to  $x$  will show us how  $y$  changes as the value  $x$  changes. It gives us the slope, or **gradient** of the function.

The derivative of  $y = f(x)$  with respect to  $x$  is represented by the following notations:

$$f'(x), \quad \frac{d}{dx}(f(x)), \quad \frac{df}{dx} \quad \text{or} \quad \frac{dy}{dx}.$$

The process of finding the derivative is called *differentiation*.

# Derivative of a function

A derivative is concerned with how one quantity changes with respect to another quantity, in other words a rate of change.

The derivative of the function  $y = f(x)$  with respect to  $x$  will show us how  $y$  changes as the value  $x$  changes. It gives us the slope, or **gradient** of the function.

The derivative of  $y = f(x)$  with respect to  $x$  is represented by the following notations:

$$f'(x), \quad \frac{d}{dx}(f(x)), \quad \frac{df}{dx} \quad \text{or} \quad \frac{dy}{dx}.$$

The process of finding the derivative is called *differentiation*.

# Differentiation of linear functions

Differentiating a linear function is to simply find the gradient or slope of that function. You learnt about gradients of linear functions in Day 2.

For example:

1. Differentiate  $f(x) = 3x - 2$ .

Since  $f(x)$  is a linear function with gradient 3,  $f'(x) = 3$ .

2. Differentiate  $f(x) = 9$ .

Since  $f(x)$  is a constant (horizontal line),  $f'(x) = 0$ .

3. Differentiate  $y = 4 - 5x$ .

The slope of  $y$  is  $-5$ , so  $\frac{dy}{dx} = -5$ .

# Differentiation of linear functions

Differentiating a linear function is to simply find the gradient or slope of that function. You learnt about gradients of linear functions in Day 2.

For example:

1. Differentiate  $f(x) = 3x - 2$ .

Since  $f(x)$  is a linear function with gradient 3,  $f'(x) = 3$ .

2. Differentiate  $f(x) = 9$ .

Since  $f(x)$  is a constant (horizontal line),  $f'(x) = 0$ .

3. Differentiate  $y = 4 - 5x$ .

The slope of  $y$  is  $-5$ , so  $\frac{dy}{dx} = -5$ .

# Differentiation of linear functions

Differentiating a linear function is to simply find the gradient or slope of that function. You learnt about gradients of linear functions in Day 2.

For example:

1. Differentiate  $f(x) = 3x - 2$ .

Since  $f(x)$  is a linear function with gradient 3,  $f'(x) = 3$ .

2. Differentiate  $f(x) = 9$ .

Since  $f(x)$  is a constant (horizontal line),  $f'(x) = 0$ .

3. Differentiate  $y = 4 - 5x$ .

The slope of  $y$  is  $-5$ , so  $\frac{dy}{dx} = -5$ .



# Differentiation of linear functions

Differentiating a linear function is to simply find the gradient or slope of that function. You learnt about gradients of linear functions in Day 2.

For example:

1. Differentiate  $f(x) = 3x - 2$ .

Since  $f(x)$  is a linear function with gradient 3,  $f'(x) = 3$ .

2. Differentiate  $f(x) = 9$ .

Since  $f(x)$  is a constant (horizontal line),  $f'(x) = 0$ .


3. Differentiate  $y = 4 - 5x$ .

The slope of  $y$  is  $-5$ , so  $\frac{dy}{dx} = -5$ .

# Differentiation of a quadratic

The graph of a quadratic functions such as  $y = x^2$  is a parabola. This means that the gradient is changing as  $x$  changes and we cannot calculate the derivative with methods used previously.


So instead we must find the gradient of a tangent to the curve at a certain point.



# Differentiation of a quadratic

The graph of a quadratic functions such as  $y = x^2$  is a parabola. This means that the gradient is changing as  $x$  changes and we cannot calculate the derivative with methods used previously.


So instead we must find the gradient of a tangent to the curve at a certain point.



# Differentiation of a quadratic

The graph of a quadratic functions such as  $y = x^2$  is a parabola. This means that the gradient is changing as  $x$  changes and we cannot calculate the derivative with methods used previously.


So instead we must find the gradient of a tangent to the curve at a certain point.



# Differentiation of a quadratic

To find the gradient of the tangent to the curve  $y = x^2$  we first take an arbitrary point  $(x, y)$  that is on the curve.


Then take another point on the curve with the x-coordinate  $x + h$ , where  $h$  is a small number. Its corresponding y-coordinate is  $(x + h)^2$ .



# Differentiation of a quadratic


To find the gradient of the tangent to the curve  $y = x^2$  we first take an arbitrary point  $(x, y)$  that is on the curve.

Then take another point on the curve with the x-coordinate  $x + h$ , where  $h$  is a small number. Its corresponding y-coordinate is  $(x + h)^2$ .



# Differentiation of a quadratic

Draw a line through the 2 points.




The gradient of this line is:

$$\frac{\text{change in } y}{\text{change in } x} = \frac{(x+h)^2 - x^2}{(x+h) - x} = \frac{2xh + h^2}{h} = 2x + h.$$

# Differentiation of a quadratic

Draw a line through the 2 points.



The gradient of this line is:



$$\frac{\text{change in } y}{\text{change in } x} = \frac{(x+h)^2 - x^2}{(x+h) - x} = \frac{2xh + h^2}{h} = 2x + h.$$



# Differentiation of a quadratic

So, the gradient of the line through the 2 points =  $2x + h$ . To find the gradient of the tangent we need another step.



Notice that as the value of  $h$  becomes smaller the line through the 2 points becomes closer to the tangent at  $(x, y)$ .



# Differentiation of a quadratic

So, the gradient of the line through the 2 points  $= 2x + h$ . To find the gradient of the tangent we need another step.

Notice that as the value of  $h$  becomes smaller the line through the 2 points becomes closer to the tangent at  $(x, y)$ .



# Differentiation of a quadratic

As the value of  $h$  approaches 0, the gradient of the line through the two points approaches the gradient of the tangent to the curve, i.e.  $2x + h$  approaches  $2x$ .

Hence the gradient of the tangent to  $y = x^2$  at point  $(x, y)$  is  $2x$  or  $\frac{dy}{dx} = 2x$ .

This method is called the *Differentiation By First Principles*. In general, the derivative of a function  $f$  at  $x$  is given by:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

if it exists.

# Differentiation of a quadratic

As the value of  $h$  approaches 0, the gradient of the line through the two points approaches the gradient of the tangent to the curve, i.e.  $2x + h$  approaches  $2x$ .

Hence the gradient of the tangent to  $y = x^2$  at point  $(x, y)$  is  $2x$  or  $\frac{dy}{dx} = 2x$ .

This method is called the *Differentiation By First Principles*. In general, the derivative of a function  $f$  at  $x$  is given by:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

if it exists.

# Differentiation of a quadratic

As the value of  $h$  approaches 0, the gradient of the line through the two points approaches the gradient of the tangent to the curve, i.e.  $2x + h$  approaches  $2x$ .

Hence the gradient of the tangent to  $y = x^2$  at point  $(x, y)$  is  $2x$  or  $\frac{dy}{dx} = 2x$ .

This method is called the *Differentiation By First Principles*. In general, the derivative of a function  $f$  at  $x$  is given by:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

if it exists.

# Differentiation of a quadratic

As the value of  $h$  approaches 0, the gradient of the line through the two points approaches the gradient of the tangent to the curve, i.e.  $2x + h$  approaches  $2x$ .

Hence the gradient of the tangent to  $y = x^2$  at point  $(x, y)$  is  $2x$  or  $\frac{dy}{dx} = 2x$ .

This method is called the *Differentiation By First Principles*. In general, the derivative of a function  $f$  at  $x$  is given by:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

if it exists.

# Differentiation of a quadratic

So, putting it all together, if  $f(x) = x^2$

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} \\ &= \lim_{h \rightarrow 0} 2x + h \\ &= 2x. \end{aligned}$$

# Some differentiation rules

The following are some rules of differentiation:

►  $\frac{d}{dx}(\text{constant}) = 0, \quad \frac{d}{dx}(x) = 1, \quad \frac{d}{dx}(x^2) = 2x.$

►  $\frac{d}{dx}(kf(x)) = k \frac{d}{dx}(f(x)),$  where  $k$  is a constant.

►  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x)).$



# Some differentiation rules

The following are some rules of differentiation:

- ▶  $\frac{d}{dx}(\text{constant}) = 0$ ,  $\frac{d}{dx}(x) = 1$ ,  $\frac{d}{dx}(x^2) = 2x$ .
- ▶  $\frac{d}{dx}(kf(x)) = k\frac{d}{dx}(f(x))$ , where  $k$  is a constant.
- ▶  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x))$ .

# Some differentiation rules

The following are some rules of differentiation:

- ▶  $\frac{d}{dx}(\text{constant}) = 0$ ,  $\frac{d}{dx}(x) = 1$ ,  $\frac{d}{dx}(x^2) = 2x$ .
- ▶  $\frac{d}{dx}(kf(x)) = k\frac{d}{dx}(f(x))$ , where  $k$  is a constant.
- ▶  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x))$ .

# Some differentiation rules

The following are some rules of differentiation:

- ▶  $\frac{d}{dx}(\text{constant}) = 0$ ,  $\frac{d}{dx}(x) = 1$ ,  $\frac{d}{dx}(x^2) = 2x$ .
- ▶  $\frac{d}{dx}(kf(x)) = k\frac{d}{dx}(f(x))$ , where  $k$  is a constant.
- ▶  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x))$ .

# Some differentiation rules

The following are some rules of differentiation:

- ▶  $\frac{d}{dx}(\text{constant}) = 0$ ,  $\frac{d}{dx}(x) = 1$ ,  $\frac{d}{dx}(x^2) = 2x$ .
- ▶  $\frac{d}{dx}(kf(x)) = k\frac{d}{dx}(f(x))$ , where  $k$  is a constant.
- ▶  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x))$ .

# Some differentiation rules

The following are some rules of differentiation:

- ▶  $\frac{d}{dx}(\text{constant}) = 0$ ,  $\frac{d}{dx}(x) = 1$ ,  $\frac{d}{dx}(x^2) = 2x$ .
- ▶  $\frac{d}{dx}(kf(x)) = k\frac{d}{dx}(f(x))$ , where  $k$  is a constant.
- ▶  $\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}(f(x)) + \frac{d}{dx}(g(x))$ .

## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

$$\text{Hence } \frac{dy}{dx} = 10x + 3.$$

## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

$$\text{Hence } \frac{dy}{dx} = 10x + 3.$$



## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

$$\text{Hence } \frac{dy}{dx} = 10x + 3.$$





## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

$$\text{Hence } \frac{dy}{dx} = 10x + 3.$$

## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

Hence  $\frac{dy}{dx} = 10x + 3$ .

## Example

Differentiate  $y = 5x^2 + 3x - 4$ .

$$\frac{dy}{dx} = \frac{d}{dx}(5x^2 + 3x - 4) = \frac{d}{dx}(5x^2) + \frac{d}{dx}(3x) - \frac{d}{dx}(4).$$

Now

$$\frac{d}{dx}(5x^2) = 5 \frac{d}{dx}(x^2) = 5 \times 2x = 10x,$$

$$\frac{d}{dx}(3x) = 3 \frac{d}{dx}(x) = 3 \times 1 = 3,$$

$$\frac{d}{dx}(4) = 0.$$

$$\text{Hence } \frac{dy}{dx} = 10x + 3.$$

## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .

## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .

## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .

## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .

## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .



## Example

Differentiate  $f(x) = 2 - 4x - x^2$ .

$$f'(x) = \frac{d}{dx}(2) - \frac{d}{dx}(4x) - \frac{d}{dx}(x^2).$$

Now

$$\frac{d}{dx}(2) = 0,$$

$$\frac{d}{dx}(4x) = 4$$

$$\frac{d}{dx}(x^2) = 2x.$$

Hence  $f'(x) = -4 - 2x$ .

## Practice Questions

Differentiate the following:

1.  $f(x) = 2x - 5$

2.  $y = 9 - 2x$

3.  $f(x) = 3x^2 + 4x - 5$

4.  $f(x) = x^2 - 4x - 6$

5.  $y = x^2 - 5x$

6.  $m = 2n^2 - 2n + 1$

7.  $y = 7$

8.  $q = p - 6p^2$

9.  $f(a) = 4a^2 + 5a - 9$

10.  $f(x) = 6x - 4x^2$ .

## Answers to practice questions:

1.  $f'(x) = 2$

2.  $\frac{dy}{dx} = -2$

3.  $f'(x) = 6x + 4$

4.  $f'(x) = 2x - 4$

5.  $\frac{dy}{dx} = 2x - 5$

6.  $\frac{dm}{dn} = 4n - 2$

7.  $\frac{dy}{dx} = 0$

8.  $\frac{dq}{dp} = 1 - 12p$

9.  $f'(a) = 8a + 5$

10.  $f'(x) = 6 - 8x$ .