

2 Unit Bridging Course – Day 10

Circular Functions I – The sine function

Clinton Boys



THE UNIVERSITY OF
SYDNEY

We're going to study two new functions called **sin** and **cos**.

You can calculate various values of these functions using the `sin` and `cos` buttons on your calculator, providing you remember:

Radians

When using `sin` and `cos` on your calculator, **always** make sure you are in **radians mode**.

We're going to study two new functions called **sin** and **cos**.

You can calculate various values of these functions using the **sin** and **cos** buttons on your calculator, providing you remember:

Radians

When using **sin** and **cos** on your calculator, **always** make sure you are in **radians mode**.

We're going to study two new functions called **sin** and **cos**.

You can calculate various values of these functions using the `sin` and `cos` buttons on your calculator, providing you remember:

Radians

When using `sin` and `cos` on your calculator, **always** make sure you are in **radians mode**.

For example, use your calculator to find the following values of the sin and cos functions:

$$\sin(0) = 0$$

$$\sin(\pi) = 0$$

$$\sin(\pi/2) = 1$$

$$\cos(0) = 1$$

$$\cos(\pi) = -1$$

$$\cos(\pi/2) = 0.$$

Let's start by studying sin in more detail.

The **sine function** is abbreviated to sin, but is still pronounced “sine”.

The most important property of the sin function is that it is **periodic**.

This means that the graph of $y = \sin(x)$ **repeats** itself after a certain x -value is passed.

Check by using your calculator that

$$\sin(-2\pi) = \sin(0) = \sin(2\pi) = \sin(4\pi).$$

Indeed, the graph of $y = \sin(x)$ repeats itself every 2π .

Let's start by studying \sin in more detail.

The **sine function** is abbreviated to \sin , but is still pronounced “sine”.

The most important property of the \sin function is that it is **periodic**.

This means that the graph of $y = \sin(x)$ **repeats** itself after a certain x -value is passed.

Check by using your calculator that

$$\sin(-2\pi) = \sin(0) = \sin(2\pi) = \sin(4\pi).$$

Indeed, the graph of $y = \sin(x)$ repeats itself every 2π .

Let's start by studying \sin in more detail.

The **sine function** is abbreviated to \sin , but is still pronounced “sine”.

The most important property of the \sin function is that it is **periodic**.

This means that the graph of $y = \sin(x)$ **repeats** itself after a certain x -value is passed.

Check by using your calculator that

$$\sin(-2\pi) = \sin(0) = \sin(2\pi) = \sin(4\pi).$$

Indeed, the graph of $y = \sin(x)$ repeats itself every 2π .

Let's start by studying \sin in more detail.

The **sine function** is abbreviated to \sin , but is still pronounced “sine”.

The most important property of the \sin function is that it is **periodic**.

This means that the graph of $y = \sin(x)$ **repeats** itself after a certain x -value is passed.

Check by using your calculator that

$$\sin(-2\pi) = \sin(0) = \sin(2\pi) = \sin(4\pi).$$

Indeed, the graph of $y = \sin(x)$ repeats itself every 2π .

Let's start by studying \sin in more detail.

The **sine function** is abbreviated to \sin , but is still pronounced “sine”.

The most important property of the \sin function is that it is **periodic**.

This means that the graph of $y = \sin(x)$ **repeats** itself after a certain x -value is passed.


Check by using your calculator that

$$\sin(-2\pi) = \sin(0) = \sin(2\pi) = \sin(4\pi).$$

Indeed, the graph of $y = \sin(x)$ repeats itself every 2π .

Graph of $y = \sin(x)$

Below is the graph of $y = \sin(x)$ between $x = -4\pi$ and $x = 4\pi$.



The graph continues forever in both directions.

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.

- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.

- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.

- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.

- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.

- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .

We can read all of the following important properties off the graph of $y = \sin(x)$:

- (i) $-1 \leq \sin x \leq 1$ for all x .

$\sin(x)$ never goes above 1 or below -1 . We say that 1 is the **amplitude** of the function.


- (ii) $\sin(x + 2\pi) = \sin x$ for all x .

The function repeats itself every 2π – the value of the function 2π units further along the x -axis from some point a is the same as the value at a . We say the function is **periodic** with **period** 2π .



Properties of sine

(iii) $\sin(-x) = -\sin(x)$ for all x .




(For example here, $\sin(\pi/2) = 1$ while $\sin(-\pi/2) = -1$). We say the function is **odd**.




Properties of sine

(iii) $\sin(-x) = -\sin(x)$ for all x .




(For example here, $\sin(\pi/2) = 1$ while $\sin(-\pi/2) = -1$). We say the function is **odd**.

(iii) $\sin(-x) = -\sin(x)$ for all x .



(For example here, $\sin(\pi/2) = 1$ while $\sin(-\pi/2) = -1$). We say the function is **odd**.

(iii) $\sin(-x) = -\sin(x)$ for all x .



(For example here, $\sin(\pi/2) = 1$ while $\sin(-\pi/2) = -1$). We say the function is **odd**.