

# 2 Unit Bridging Course – Day 2

## Linear functions II: Finding equations

Clinton Boys



THE UNIVERSITY OF  
SYDNEY

# Finding equations of lines

If we have the information of

- (i) the gradient of a line
- (ii) the coordinates of a point on the line

then we can find its equation.

# Finding equations of lines

If we have the information of

- (i) the gradient of a line
- (ii) the coordinates of a point on the line

then we can find its equation.

# Finding equations of lines

If we have the information of

- (i) the gradient of a line
- (ii) the coordinates of a point on the line

then we can find its equation.

# Finding equations of lines

Let's think about how we would go about doing this.

## Example

Suppose we know a line has gradient 2 and passes through the point with co-ordinates  $(-1, 2)$ .

Since  $m = 2$ , we know the line must have the equation

$$y = 2x + b$$

for some number  $b$ .

# Finding equations of lines

Let's think about how we would go about doing this.

## Example

Suppose we know a line has gradient 2 and passes through the point with co-ordinates  $(-1, 2)$ .

Since  $m = 2$ , we know the line must have the equation

$$y = 2x + b$$

for some number  $b$ .

# Finding equations of lines

Let's think about how we would go about doing this.

## Example

Suppose we know a line has gradient 2 and passes through the point with co-ordinates  $(-1, 2)$ .

Since  $m = 2$ , we know the line must have the equation

$$y = 2x + b$$

for some number  $b$ .

## Example

We also know that  $(-1, 2)$  lies on the line, i.e. that this point **satisfies** the equation. This means

$$2 = 2 \times (-1) + b.$$

Solving this equation for  $b$  gives

$$b = 2 + 2 = 4,$$

so the equation of our line is

$$y = 2x + 4.$$



## Example

We also know that  $(-1, 2)$  lies on the line, i.e. that this point **satisfies** the equation. This means

$$2 = 2 \times (-1) + b.$$

Solving this equation for  $b$  gives

$$b = 2 + 2 = 4,$$

so the equation of our line is

$$y = 2x + 4.$$

## Example

We also know that  $(-1, 2)$  lies on the line, i.e. that this point **satisfies** the equation. This means


$$2 = 2 \times (-1) + b.$$

Solving this equation for  $b$  gives

$$b = 2 + 2 = 4,$$

so the equation of our line is

$$y = 2x + 4.$$



$$y = 2x + 4$$

## Practice questions

Find the equations of the following lines:

- (i) The line with gradient  $-1$  which passes through  $(3, 1)$ .
- (ii) The line with gradient  $2$  which passes through  $(-1, -4)$ .

## Answers to practice questions

(i)  $y = -x + 4$

(ii)  $y = 2x - 2$

# Rise-over-run formula

If we know two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on a line, we can use the following formula, known as the **rise-over-run formula** to compute the gradient of the line between the two points:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{y_2 - y_1}{x_2 - x_1}.$$

Let's see how we can use this formula to find the equation of a line.

# Rise-over-run formula

If we know two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on a line, we can use the following formula, known as the **rise-over-run formula** to compute the gradient of the line between the two points:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{y_2 - y_1}{x_2 - x_1}.$$

Let's see how we can use this formula to find the equation of a line.

# Rise-over-run formula

If we know two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on a line, we can use the following formula, known as the **rise-over-run formula** to compute the gradient of the line between the two points:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{y_2 - y_1}{x_2 - x_1}.$$

Let's see how we can use this formula to find the equation of a line.



# Equations of lines, continued

If we know the coordinates of **two** points which lie on a straight line, we can:

- (i) Use the rise-over-run formula to find the gradient.
- (ii) Use the gradient and one of the points to find the equation (as before).

# Equations of lines, continued

If we know the coordinates of **two** points which lie on a straight line, we can:

- (i) Use the rise-over-run formula to find the gradient.
- (ii) Use the gradient and one of the points to find the equation (as before).

## Example

Let's find the equation of the straight line which joins the points  $(-1, 2)$  and  $(1, 3)$ .

Using the rise-over-run formula we can calculate the gradient:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{2 - 3}{-1 - 1} = \frac{-1}{-2} = \frac{1}{2}.$$

## Example

Let's find the equation of the straight line which joins the points  $(-1, 2)$  and  $(1, 3)$ .

Using the rise-over-run formula we can calculate the gradient:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{2 - 3}{-1 - 1} = \frac{-1}{-2} = \frac{1}{2}.$$

## Example

Let's find the equation of the straight line which joins the points  $(-1, 2)$  and  $(1, 3)$ .

Using the rise-over-run formula we can calculate the gradient:

$$m = \frac{\text{change in } y}{\text{change in } x} = \frac{2 - 3}{-1 - 1} = \frac{-1}{-2} = \frac{1}{2}.$$

We can now use either of the points  $(-1, 2)$  or  $(1, 3)$  to find the equation. Let's use  $(1, 3)$ .

If the equation is  $y = \frac{1}{2}x + b$ , then the point  $(1, 3)$  must satisfy this equation, and so we must have

$$3 = \frac{1}{2} \times 1 + b,$$

i.e.  $b = 3 - \frac{1}{2} = 2\frac{1}{2} = \frac{5}{2}$ .

So the equation of the line is

$$y = \frac{1}{2}x + \frac{5}{2} = \frac{1}{2}(x + 5).$$

We can now use either of the points  $(-1, 2)$  or  $(1, 3)$  to find the equation. Let's use  $(1, 3)$ .

If the equation is  $y = \frac{1}{2}x + b$ , then the point  $(1, 3)$  must satisfy this equation, and so we must have

$$3 = \frac{1}{2} \times 1 + b,$$

i.e.  $b = 3 - \frac{1}{2} = 2\frac{1}{2} = \frac{5}{2}$ .

So the equation of the line is

$$y = \frac{1}{2}x + \frac{5}{2} = \frac{1}{2}(x + 5).$$

We can now use either of the points  $(-1, 2)$  or  $(1, 3)$  to find the equation. Let's use  $(1, 3)$ .

If the equation is  $y = \frac{1}{2}x + b$ , then the point  $(1, 3)$  must satisfy this equation, and so we must have

$$3 = \frac{1}{2} \times 1 + b,$$

i.e.  $b = 3 - \frac{1}{2} = 2\frac{1}{2} = \frac{5}{2}$ .

So the equation of the line is

$$y = \frac{1}{2}x + \frac{5}{2} = \frac{1}{2}(x + 5).$$



## Practice question

Check that you can also get this equation using the other point  $(-1, 2)$ .

# Graphs of linear functions

The graph of a linear function is a straight line, as we have already seen. In order to sketch the graph of a straight line from its equation, we need to know two points which lie on the line.

Two points which are particularly easy to find are the *x-intercept* and the *y-intercept*.

The *x-intercept*, which is where the line crosses the *x*-axis, is found by *setting*  $y = 0$  (because  $y = 0$  along the entire *x*-axis).

The *y-intercept*, which is where the line crosses the *y*-axis, is found by *setting*  $x = 0$  (because  $x = 0$  along the entire *y*-axis).

# Graphs of linear functions

The graph of a linear function is a straight line, as we have already seen. In order to sketch the graph of a straight line from its equation, we need to know two points which lie on the line.

Two points which are particularly easy to find are the **x-intercept** and the **y-intercept**.

The x-intercept, which is where the line crosses the x-axis, is found by *setting*  $y = 0$  (because  $y = 0$  along the entire x-axis).

The y-intercept, which is where the line crosses the y-axis, is found by *setting*  $x = 0$  (because  $x = 0$  along the entire y-axis).

# Graphs of linear functions

The graph of a linear function is a straight line, as we have already seen. In order to sketch the graph of a straight line from its equation, we need to know two points which lie on the line.

Two points which are particularly easy to find are the **x-intercept** and the **y-intercept**.

The x-intercept, which is where the line crosses the x-axis, is found by *setting*  $y = 0$  (because  $y = 0$  along the entire x-axis).

The y-intercept, which is where the line crosses the y-axis, is found by *setting*  $x = 0$  (because  $x = 0$  along the entire y-axis).

# Graphs of linear functions

The graph of a linear function is a straight line, as we have already seen. In order to sketch the graph of a straight line from its equation, we need to know two points which lie on the line.

Two points which are particularly easy to find are the ***x*-intercept** and the ***y*-intercept**.

The *x*-intercept, which is where the line crosses the *x*-axis, is found by *setting*  $y = 0$  (because  $y = 0$  along the entire *x*-axis).

The *y*-intercept, which is where the line crosses the *y*-axis, is found by *setting*  $x = 0$  (because  $x = 0$  along the entire *y*-axis).

## Example

Draw the graph of the straight line  $2y + x - 4 = 0$ .

The  $x$ -intercept is found by setting  $y = 0$ :

$$x - 4 = 0, \quad \text{i.e.} \quad x = 4.$$

The  $y$ -intercept is found by setting  $x = 0$ :

$$2y - 4 = 0, \quad \text{i.e.} \quad 2y = 4, \quad \text{i.e.} \quad y = 2.$$

# Graphs of linear functions

## Example

Draw the graph of the straight line  $2y + x - 4 = 0$ .

The  $x$ -intercept is found by setting  $y = 0$ :

$$x - 4 = 0, \quad \text{i.e.} \quad x = 4.$$

The  $y$ -intercept is found by setting  $x = 0$ :

$$2y - 4 = 0, \quad \text{i.e.} \quad 2y = 4, \quad \text{i.e.} \quad y = 2.$$

# Graphs of linear functions

## Example

Draw the graph of the straight line  $2y + x - 4 = 0$ .

The  $x$ -intercept is found by setting  $y = 0$ :

$$x - 4 = 0, \quad \text{i.e.} \quad x = 4.$$

The  $y$ -intercept is found by setting  $x = 0$ :

$$2y - 4 = 0, \quad \text{i.e.} \quad 2y = 4, \quad \text{i.e.} \quad y = 2.$$



## Example

Draw the graph of the straight line  $2y + x - 4 = 0$ .

The  $x$ -intercept is found by setting  $y = 0$ :

$$x - 4 = 0, \quad \text{i.e.} \quad x = 4.$$

The  $y$ -intercept is found by setting  $x = 0$ :

$$2y - 4 = 0, \quad \text{i.e.} \quad 2y = 4, \quad \text{i.e.} \quad y = 2.$$

## Example


Draw the graph of the straight line  $2y + x - 4 = 0$ .

The  $x$ -intercept is found by setting  $y = 0$ :


$$x - 4 = 0, \quad \text{i.e.} \quad x = 4.$$

The  $y$ -intercept is found by setting  $x = 0$ :

$$2y - 4 = 0, \quad \text{i.e.} \quad 2y = 4, \quad \text{i.e.} \quad y = 2.$$




$x$ -intercept  $x = 4$




$x$ -intercept  $x = 4$

$y$ -intercept  $y = 2$



$x$ -intercept  $x = 4$

$y$ -intercept  $y = 2$



$$2y + x - 4 = 0$$