

2 Unit Bridging Course – Day 6

Application of the second derivative: Curve sketching

Emi Tanaka



THE UNIVERSITY OF
SYDNEY

The second derivative: Curve sketching

In the last module, we saw how to use the second derivative to find points of inflexion.

We can also use the second derivative and concavity to determine the nature of stationary points.

Both can be used to help sketch the function.

The second derivative: Curve sketching

In the last module, we saw how to use the second derivative to find points of inflexion.

We can also use the second derivative and concavity to determine the nature of stationary points.

Both can be used to help sketch the function.

The second derivative: Curve sketching

In the last module, we saw how to use the second derivative to find points of inflexion.

We can also use the second derivative and concavity to determine the nature of stationary points.


Both can be used to help sketch the function.

Second derivative test: Minimum

Recall: If $f''(x) > 0$ over an interval, then the function f is concave up over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) > 0$, the curve is concave up at $x = p$, and f has a local minimum at p .




Second derivative test: Minimum

Recall: If $f''(x) > 0$ over an interval, then the function f is concave up over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) > 0$, the curve is concave up at $x = p$, and f has a local minimum at p .




Second derivative test: Minimum

Recall: If $f''(x) > 0$ over an interval, then the function f is concave up over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) > 0$, the curve is concave up at $x = p$, and f has a local minimum at p .




Second derivative test: Maximum

Recall: If $f''(x) < 0$ over an interval, then the function f is concave down over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) < 0$, the curve is concave down at $x = p$, and f has a local maximum at p .




Second derivative test: Maximum

Recall: If $f''(x) < 0$ over an interval, then the function f is concave down over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) < 0$, the curve is concave down at $x = p$, and f has a local maximum at p .




Second derivative test: Maximum

Recall: If $f''(x) < 0$ over an interval, then the function f is concave down over that interval.

Suppose that f has a stationary point at $x = p$, ie $f'(p) = 0$.

Then if $f''(p) < 0$, the curve is concave down at $x = p$, and f has a local maximum at p .



Second derivative test: Summary

- ▶ If $f'(p) = 0$ and $f''(p) > 0$, then the function f has a local minimum at p .
- ▶ If $f'(p) = 0$ and $f''(p) < 0$, then the function f has a local maximum at p .
- ▶ If $f'(p) = 0$ and $f''(p) = 0$, then the test tells us nothing.

So, if $f'(p) = 0$ and $f''(p) = 0$, then we must look at the first derivative to determine the nature of the stationary point like we did on Day 5.

Second derivative test: Summary

- ▶ If $f'(p) = 0$ and $f''(p) > 0$, then the function f has a local minimum at p .
- ▶ If $f'(p) = 0$ and $f''(p) < 0$, then the function f has a local maximum at p .
- ▶ If $f'(p) = 0$ and $f''(p) = 0$, then the test tells us nothing.

So, if $f'(p) = 0$ and $f''(p) = 0$, then we must look at the first derivative to determine the nature of the stationary point like we did on Day 5.

Second derivative test: Summary

- ▶ If $f'(p) = 0$ and $f''(p) > 0$, then the function f has a local minimum at p .
- ▶ If $f'(p) = 0$ and $f''(p) < 0$, then the function f has a local maximum at p .
- ▶ If $f'(p) = 0$ and $f''(p) = 0$, then the test tells us nothing.

So, if $f'(p) = 0$ and $f''(p) = 0$, then we must look at the first derivative to determine the nature of the stationary point like we did on Day 5.

Second derivative test: Summary

- ▶ If $f'(p) = 0$ and $f''(p) > 0$, then the function f has a local minimum at p .
- ▶ If $f'(p) = 0$ and $f''(p) < 0$, then the function f has a local maximum at p .
- ▶ If $f'(p) = 0$ and $f''(p) = 0$, then the test tells us nothing.

So, if $f'(p) = 0$ and $f''(p) = 0$, then we must look at the first derivative to determine the nature of the stationary point like we did on Day 5.

Example

Sketch the curve $y = f(x) = x^3 - 3x$.

First find any stationary points:

$$f'(x) = 3x^2 - 3 = 3(x - 1)(x + 1) = 0,$$

so the stationary points are $(1, -2)$ and $(-1, 2)$.

Next find the second derivative:

$$f''(x) = 6x.$$

Example

Sketch the curve $y = f(x) = x^3 - 3x$.

First find any stationary points:

$$f'(x) = 3x^2 - 3 = 3(x - 1)(x + 1) = 0,$$

so the stationary points are $(1, -2)$ and $(-1, 2)$.

Next find the second derivative:

$$f''(x) = 6x.$$

Example

Sketch the curve $y = f(x) = x^3 - 3x$.

First find any stationary points:

$$f'(x) = 3x^2 - 3 = 3(x - 1)(x + 1) = 0,$$

so the stationary points are $(1, -2)$ and $(-1, 2)$.

Next find the second derivative:

$$f''(x) = 6x.$$

The second derivative is

$$f''(x) = 6x.$$

Use the second derivative to test the stationary points:

$$f''(1) = 6,$$

since $f''(1) > 0$ the curve is concave up, hence $(1, -2)$ is a local minimum;

$$f''(-1) = -6$$

since $f''(-1) < 0$ the curve is concave down, hence $(-1, 2)$ is a local maximum.

The second derivative is

$$f''(x) = 6x.$$

Use the second derivative to test the stationary points:

$$f''(1) = 6,$$

since $f''(1) > 0$ the curve is concave up, hence $(1, -2)$ is a local minimum;

$$f''(-1) = -6$$

since $f''(-1) < 0$ the curve is concave down, hence $(-1, 2)$ is a local maximum.

The second derivative is

$$f''(x) = 6x.$$

Use the second derivative to test the stationary points:

$$f''(1) = 6,$$

since $f''(1) > 0$ the curve is concave up, hence $(1, -2)$ is a local minimum;

$$f''(-1) = -6$$

since $f''(-1) < 0$ the curve is concave down, hence $(-1, 2)$ is a local maximum.

Notice that $f''(x) = 6x = 0$ when $x = 0$, this means we have a **possible** point of inflexion.

To test if this is a point of inflexion, we check the second derivative slightly before and after the point.

When

$$x < 0, f''(x) < 0$$

and when

$$x > 0, f''(x) > 0.$$

This means that the concavity of the curve is changing at $x = 0$, hence there is a point of inflexion at $(0, 0)$.

Notice that $f''(x) = 6x = 0$ when $x = 0$, this means we have a **possible** point of inflexion.

To test if this is a point of inflexion, we check the second derivative slightly before and after the point.

When

$$x < 0, f''(x) < 0$$

and when

$$x > 0, f''(x) > 0.$$

This means that the concavity of the curve is changing at $x = 0$, hence there is a point of inflexion at $(0, 0)$.

Notice that $f''(x) = 6x = 0$ when $x = 0$, this means we have a **possible** point of inflexion.

To test if this is a point of inflexion, we check the second derivative slightly before and after the point.

When

$$x < 0, f''(x) < 0$$

and when

$$x > 0, f''(x) > 0.$$

This means that the concavity of the curve is changing at $x = 0$, hence there is a point of inflexion at $(0, 0)$.

Notice that $f''(x) = 6x = 0$ when $x = 0$, this means we have a **possible** point of inflexion.

To test if this is a point of inflexion, we check the second derivative slightly before and after the point.

When

$$x < 0, f''(x) < 0$$

and when

$$x > 0, f''(x) > 0.$$

This means that the concavity of the curve is changing at $x = 0$, hence there is a point of inflexion at $(0, 0)$.

The curve sketched:

