

# ASDS Statistics, YSU, Fall 2020

## Lecture 12

Michael Poghosyan

10 Oct 2020

# Contents

- ▶ QQ Plot
- ▶ Sample Covariance and Correlation Coefficient

## Q-Q Plots, Theoretical vs Theoretical Distribution

Assume now we have two Theoretical Distributions (say, given by their CDFs  $F$  and  $G$ ).

## Q-Q Plots, Theoretical vs Theoretical Distribution

Assume now we have two Theoretical Distributions (say, given by their CDFs  $F$  and  $G$ ). The Problem is to estimate visually which Distribution has fatter tails.

## Q-Q Plots, Theoretical vs Theoretical Distribution

Assume now we have two Theoretical Distributions (say, given by their CDFs  $F$  and  $G$ ). The Problem is to estimate visually which Distribution has fatter tails.

To answer this question, we again take some levels of quantiles, say, for some  $n$ ,

$$\alpha = \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}$$

and then draw the points  $(q_{\alpha}^F, q_{\alpha}^G)$ , where  $q_{\alpha}^F$  is the  $\alpha$ -quantile of the Theoretical Distribution with the CDF  $F$ , and  $q_{\alpha}^G$  is the  $\alpha$ -quantile of the Theoretical Distribution with the CDF  $G$ .

## Q-Q Plots, Theoretical vs Theoretical Distribution

Assume now we have two Theoretical Distributions (say, given by their CDFs  $F$  and  $G$ ). The Problem is to estimate visually which Distribution has fatter tails.

To answer this question, we again take some levels of quantiles, say, for some  $n$ ,



$$\alpha = \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}$$

and then draw the points  $(q_{\alpha}^F, q_{\alpha}^G)$ , where  $q_{\alpha}^F$  is the  $\alpha$ -quantile of the Theoretical Distribution with the CDF  $F$ , and  $q_{\alpha}^G$  is the  $\alpha$ -quantile of the Theoretical Distribution with the CDF  $G$ .

**Idea:** If  $G$  has fatter tails on both sides than  $F$ , then we will have graphically some cubic-function graph shape Quantiles.

# Some Experiments


```
par(mfrow = c(1,2))
x <- rnorm(100, mean=0, sd=1); y <- rt(100, df = 3)
qqplot(x,y, main = "Q-Q Plot")
t <- seq(-4,4,0.01)
plot(t, dnorm(t), type = "l", xlim = c(-4,4), ylim = c(0, 0.5), col ="blue", lwd = 2, main = "PDFs")
par(new = TRUE)
plot(t, dt(t, df = 3), type = "l", xlim = c(-4,4), ylim = c(0, 0.5), col ="red", lwd = 2)
```



## Some Experiments

The above Datasets, using BoxPlots:



```
boxplot(x,y, horizontal = T)
```





# Some Experiments

```
par(mfrow = c(1,2))
x <- rnorm(100, mean=2, sd=1); y <- rchisq(200, df = 3)
qqplot(x,y, main = "Q-Q Plot")
t <- seq(0,5,0.01)
plot(t, dnorm(t), type = "l", xlim = c(0,5), ylim = c(0, 0.5), col ="blue", lwd = 2, main = "PDFs")
par(new = TRUE)
plot(t, dchisq(t, df = 3), type = "l", xlim = c(0,5), ylim = c(0, 0.5), col ="red", lwd = 2)
```




## Addition, Q-Q Plot

here you can find some interpretations of different shapes of Q-Q Plots: [StackExchange Page](#).

# Addition, Q-Q Plot with a Confidence Band

```
require(qqplotr)
x <- data.frame(variable = rnorm(200))
ggplot(data = x, mapping = aes(sample = variable)) + stat_qq_band() +
  stat_qq_line() + stat_qq_point() + labs(x = "Theoretical Quantiles", y = "Sample Quantiles")
```



# Numerical Summaries for Bivariate Data

# Sample Covariance and the Correlation Coefficient

Assume now we have a bivariate Dataset

$$(x_1, y_1), \dots, (x_n, y_n),$$

or just two 1D Datasets of the same size:

$$x : x_1, \dots, x_n \quad \text{and} \quad y : y_1, \dots, y_n.$$

# Sample Covariance and the Correlation Coefficient

Assume now we have a bivariate Dataset

$$(x_1, y_1), \dots, (x_n, y_n),$$

or just two 1D Datasets of the same size:

$$x : x_1, \dots, x_n \quad \text{and} \quad y : y_1, \dots, y_n.$$

Our aim is to see if some linear relationship, association exists between  $x$  and  $y$ .

# Sample Covariance and the Correlation Coefficient

Assume now we have a bivariate Dataset

$$(x_1, y_1), \dots, (x_n, y_n),$$

or just two 1D Datasets of the same size:

$$x : x_1, \dots, x_n \quad \text{and} \quad y : y_1, \dots, y_n.$$

Our aim is to see if some linear relationship, association exists between  $x$  and  $y$ . Of course, the best way is to visualize our Dataset by a ScatterPlot.

# Sample Covariance and the Correlation Coefficient

Assume now we have a bivariate Dataset

$$(x_1, y_1), \dots, (x_n, y_n),$$

or just two 1D Datasets of the same size:

$$x : x_1, \dots, x_n \quad \text{and} \quad y : y_1, \dots, y_n.$$

Our aim is to see if some linear relationship, association exists between  $x$  and  $y$ . Of course, the best way is to visualize our Dataset by a ScatterPlot.

Now we want to answer, numerically, how strong/weak is the linear relationship between our variables  $x$  and  $y$ .



## Sample Covariance

The **Sample Covariance** of Variables (1D Datasets)  $x$  and  $y$  is

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n}$$

## Sample Covariance

The **Sample Covariance** of Variables (1D Datasets)  $x$  and  $y$  is

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n}$$

or

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n - 1}$$

## Sample Covariance

The **Sample Covariance** of Variables (1D Datasets)  $x$  and  $y$  is

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n}$$

or

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n - 1}$$

Here  $\bar{x}$  and  $\bar{y}$  are the Sample Means for the Datasets  $x$  and  $y$ .

## Sample Covariance

The **Sample Covariance** of Variables (1D Datasets)  $x$  and  $y$  is

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n}$$

or

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n - 1}$$

Here  $\bar{x}$  and  $\bar{y}$  are the Sample Means for the Datasets  $x$  and  $y$ .

**Note:** Recall that for a r.v.  $X$ ,  $\text{Cov}(X, X) = \text{Var}(X)$ .

## Sample Covariance

The **Sample Covariance** of Variables (1D Datasets)  $x$  and  $y$  is

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n}$$

or

$$\text{cov}(x, y) = s_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{n - 1}$$

Here  $\bar{x}$  and  $\bar{y}$  are the Sample Means for the Datasets  $x$  and  $y$ .

**Note:** Recall that for a r.v.  $X$ ,  $\text{Cov}(X, X) = \text{Var}(X)$ . Here, for Datasets, we have two definitions for the Sample Variance  $\text{var}(x)$ . And we give two definitions of the Sample Covariance, so the property  $\text{cov}(x, x) = \text{var}(x)$  will hold in both cases.

## Sample Covariance

**Definition:** We say that the Variables (Datasets)  $x$  and  $y$  are **uncorrelated**, if  $\text{cov}(x, y) = 0$ .

## Sample Covariance

**Definition:** We say that the Variables (Datasets)  $x$  and  $y$  are **uncorrelated**, if  $cov(x, y) = 0$ .

**Remark:** In Probability, we have 2 notions: *Independence* and *Correlation*. Here, in the case of Datasets, we do not have the notion of *Independence*.

## Sample Covariance

**Definition:** We say that the Variables (Datasets)  $x$  and  $y$  are **uncorrelated**, if  $cov(x, y) = 0$ .

**Remark:** In Probability, we have 2 notions: *Independence* and *Correlation*. Here, in the case of Datasets, we do not have the notion of *Independence*.

**Remark:** For almost all numerical summaries for 1D data, first step was sorting the Dataset to obtain Order Statistics. But please note that for calculating Covariance or Correlation Coefficient (as well as for ScatterPlotting), sorting the Datasets will give incorrect results.



# Sample Covariance

**Definition:** We say that the Variables (Datasets)  $x$  and  $y$  are **uncorrelated**, if  $cov(x, y) = 0$ .

**Remark:** In Probability, we have 2 notions: *Independence* and *Correlation*. Here, in the case of Datasets, we do not have the notion of *Independence*.

**Remark:** For almost all numerical summaries for 1D data, first step was sorting the Dataset to obtain Order Statistics. But please note that for calculating Covariance or Correlation Coefficient (as well as for ScatterPlotting), sorting the Datasets will give incorrect results. This is because we want to find a relationship between  $x_1$  and  $y_1$ ,  $x_2$  and  $y_2$ ,  $\dots$ , not the relationship between the minimal elements of Datasets etc.

## Example

Here is the **R** code to calculate the Covariance between the Speed and Dist variables in the cars Dataset:

```
cov(cars$speed, cars$dist)
```

```
## [1] 109.9469
```

## Sample Correlation Coefficient

Another measure of the linear relationship between the Variables  $x$  and  $y$  of Bivariate Dataset is the *Pearson's Correlation Coefficient*:

## Sample Correlation Coefficient

Another measure of the linear relationship between the Variables  $x$  and  $y$  of Bivariate Dataset is the *Pearson's Correlation Coefficient*:

**Definition:** The **Sample Correlation Coefficient** of  $x$  and  $y$  is

$$\text{cor}(x, y) = \rho_{xy} = \frac{\text{cov}(x, y)}{\sqrt{\text{Var}(x) \cdot \text{Var}(y)}} = \frac{\text{cov}(x, y)}{\text{sd}(x) \cdot \text{sd}(y)} = \frac{s_{xy}}{s_x \cdot s_y},$$

where  $s_x$  and  $s_y$  are the standard deviations for  $x$  and  $y$ , respectively.

## Sample Correlation Coefficient

Another measure of the linear relationship between the Variables  $x$  and  $y$  of Bivariate Dataset is the *Pearson's Correlation Coefficient*:

**Definition:** The **Sample Correlation Coefficient** of  $x$  and  $y$  is

$$\text{cor}(x, y) = \rho_{xy} = \frac{\text{cov}(x, y)}{\sqrt{\text{Var}(x) \cdot \text{Var}(y)}} = \frac{\text{cov}(x, y)}{\text{sd}(x) \cdot \text{sd}(y)} = \frac{s_{xy}}{s_x \cdot s_y},$$

where  $s_x$  and  $s_y$  are the standard deviations for  $x$  and  $y$ , respectively.

If  $s_x = 0$  or  $s_y = 0$ , then we take  $\text{cor}(x, y) = 0$  by definition.

## Sample Correlation Coefficient

Another measure of the linear relationship between the Variables  $x$  and  $y$  of Bivariate Dataset is the *Pearson's Correlation Coefficient*:

**Definition:** The **Sample Correlation Coefficient** of  $x$  and  $y$  is

$$\text{cor}(x, y) = \rho_{xy} = \frac{\text{cov}(x, y)}{\sqrt{\text{Var}(x) \cdot \text{Var}(y)}} = \frac{\text{cov}(x, y)}{\text{sd}(x) \cdot \text{sd}(y)} = \frac{s_{xy}}{s_x \cdot s_y},$$

where  $s_x$  and  $s_y$  are the standard deviations for  $x$  and  $y$ , respectively.

If  $s_x = 0$  or  $s_y = 0$ , then we take  $\text{cor}(x, y) = 0$  by definition.

**Note:** Please note that we need to calculate the Standard Deviations and Covariance by using the same denominator: either everywhere take  $n$ , or take everywhere  $n - 1$ .

## Sample Correlation Coefficient

In both cases, when one calculates Standard Deviations and Covariance by using  $n$  simultaneously or  $n - 1$  simultaneously in the denominator, we will obtain

$$\text{cor}(x, y) = \rho_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{\sqrt{\sum_{k=1}^n (x_k - \bar{x})^2 \cdot \sum_{k=1}^n (y_k - \bar{y})^2}}$$

## Sample Correlation Coefficient

In both cases, when one calculates Standard Deviations and Covariance by using  $n$  simultaneously or  $n - 1$  simultaneously in the denominator, we will obtain

$$\text{cor}(x, y) = \rho_{xy} = \frac{\sum_{k=1}^n (x_k - \bar{x}) \cdot (y_k - \bar{y})}{\sqrt{\sum_{k=1}^n (x_k - \bar{x})^2 \cdot \sum_{k=1}^n (y_k - \bar{y})^2}}$$

Another formula to calc the correlation coefficient is


$$\text{cor}(x, y) = \rho_{xy} = \frac{\sum_{k=1}^n x_k y_k - n \cdot \bar{x} \cdot \bar{y}}{\sqrt{\sum_{k=1}^n x_k^2 - n \cdot (\bar{x})^2} \cdot \sqrt{\sum_{k=1}^n y_k^2 - n \cdot (\bar{y})^2}}.$$



## Examples:

Now, the **R** code to calculate the Covariance between the Speed and Dist variables in the cars Dataset:

```
plot(dist~speed, data = cars)
```




```
cor(cars$speed, cars$dist)
```

```
## [1] 0.8068949
```

## Examples:

Some simulations:

```
x <- rnorm(100); y <- rnorm(100);  
plot(x,y, pch=16)
```




```
c(cor(x,y), cov(x,y))
```

```
## [1] -0.003809632 -0.003336435
```

## Examples:

Some simulations:

```
x <- rnorm(100); y <- -2.4*x + rnorm(100);  
plot(x,y, pch=16)
```



```
c(cor(x,y), cov(x,y))
```

```
## [1] -0.9072463 -2.1401902
```

## Examples:

Let us now use the `state.x77` Dataset from **R**:

```
head(state.x77)
```

##	Population	Income	Illiteracy	Life Exp	Murder	HS Gr
## Alabama	3615	3624	2.1	69.05	15.1	41
## Alaska	365	6315	1.5	69.31	11.3	66
## Arizona	2212	4530	1.8	70.55	7.8	58
## Arkansas	2110	3378	1.9	70.66	10.1	39
## California	21198	5114	1.1	71.71	10.3	62
## Colorado	2541	4884	0.7	72.06	6.8	63

## Examples:

Let us now use the `state.x77` Dataset from **R**:

```
head(state.x77)
```


##	Population	Income	Illiteracy	Life Exp	Murder	HS Gr
## Alabama	3615	3624	2.1	69.05	15.1	41
## Alaska	365	6315	1.5	69.31	11.3	66
## Arizona	2212	4530	1.8	70.55	7.8	58
## Arkansas	2110	3378	1.9	70.66	10.1	39
## California	21198	5114	1.1	71.71	10.3	62
## Colorado	2541	4884	0.7	72.06	6.8	63

It is not of the `DataFrame` format, so we change it to `DataFrame`:

```
state <- as.data.frame(state.x77)
```

## Examples:

```
plot(Murder~Illiteracy, data = state, pch=16)
```



```
cor(state$Illiteracy, state$Murder)
```

```
## [1] 0.7029752
```

## Examples:

**Question:** How to generate samples  $x, y$  with some given Correlation Coefficient?

## Examples:

**Question:** How to generate samples  $x, y$  with some given Correlation Coefficient?

**Answer:** Say, we want to have Datasets  $x, y$  of size  $n$  with  $\text{cor}(x, y) = \rho \in (-1, 1)$ .



## Examples:

**Question:** How to generate samples  $x, y$  with some given Correlation Coefficient?

**Answer:** Say, we want to have Datasets  $x, y$  of size  $n$  with  $cor(x, y) = \rho \in (-1, 1)$ .

One of the possible methods: take a Matrix

$$\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix},$$

which is **Positive Definite**, take any 2D vector, say  $\mu = [0, 0]^T$ , and generate a Sample of size  $n$  from the Bivariate Normal Distribution  $\mathcal{N}(\mu, \Sigma)$ .

## Examples:

**Question:** How to generate samples  $x, y$  with some given Correlation Coefficient?

**Answer:** Say, we want to have Datasets  $x, y$  of size  $n$  with  $\text{cor}(x, y) = \rho \in (-1, 1)$ .

One of the possible methods: take a Matrix


$$\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix},$$

which is **Positive Definite**, take any 2D vector, say  $\mu = [0, 0]^T$ , and generate a Sample of size  $n$  from the Bivariate Normal Distribution  $\mathcal{N}(\mu, \Sigma)$ .

Then, the  $\text{cor}(x, y)$  will be approximately  $\rho$  (and it will approach  $\rho$  as  $n \rightarrow +\infty$ ).

## Example

```
rho <- 0.35  
covmatrix <- matrix(c(1, rho, rho, 1), nrow = 2)  
mu <- c(0, 0)  
x <- mvtnorm::rmvnorm(100, mean = mu, sigma = covmatrix)  
plot(x, pch = 16)
```



```
cor(x)
```

```
##           [,1]      [,2]  
## [1,] 1.0000000 0.3069956  
## [2,] 0.3069956 1.0000000
```

## Properties of the Sample Covariance

- ▶  $\text{cov}(x, y) = \text{cov}(y, x);$

# Properties of the Sample Covariance

- ▶  $\text{cov}(x, y) = \text{cov}(y, x)$ ;
- ▶ For any Datasets  $x, y, z$  and real numbers  $\alpha, \beta$ ,

$$\text{cov}(\alpha \cdot x + \beta \cdot y, z) = \alpha \cdot \text{cov}(x, z) + \beta \cdot \text{cov}(y, z);$$

# Properties of the Sample Covariance

- ▶  $cov(x, y) = cov(y, x)$ ;
- ▶ For any Datasets  $x, y, z$  and real numbers  $\alpha, \beta$ ,

$$cov(\alpha \cdot x + \beta \cdot y, z) = \alpha \cdot cov(x, z) + \beta \cdot cov(y, z);$$

- ▶ For any Dataset  $x$ ,

$$cov(x, x) = var(x)$$

# Properties of the Sample Correlation Coefficient

►  $\text{cor}(x, y) = \text{cor}(y, x);$

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

# Properties of the Sample Correlation Coefficient

- ▶  $\text{cor}(x, y) = \text{cor}(y, x)$ ;
- ▶ If  $\alpha > 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = \text{cor}(x, y)$

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).



# Properties of the Sample Correlation Coefficient

- ▶  $\text{cor}(x, y) = \text{cor}(y, x)$ ;
- ▶ If  $\alpha > 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = \text{cor}(x, y)$
- ▶ If  $\alpha < 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = -\text{cor}(x, y)$

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

# Properties of the Sample Correlation Coefficient

- ▶  $cor(x, y) = cor(y, x)$ ;
- ▶ If  $\alpha > 0$  and  $\beta \in \mathbb{R}$ , then  $cor(\alpha \cdot x + \beta, y) = cor(x, y)$
- ▶ If  $\alpha < 0$  and  $\beta \in \mathbb{R}$ , then  $cor(\alpha \cdot x + \beta, y) = -cor(x, y)$
- ▶ For any Datasets  $x, y$ ,

$$-1 \leq \rho_{xy} \leq 1;$$

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

# Properties of the Sample Correlation Coefficient

- ▶  $\text{cor}(x, y) = \text{cor}(y, x)$ ;
- ▶ If  $\alpha > 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = \text{cor}(x, y)$
- ▶ If  $\alpha < 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = -\text{cor}(x, y)$
- ▶ For any Datasets  $x, y$ ,

$$-1 \leq \rho_{xy} \leq 1;$$

- ▶  $\rho_{xy} = 1$  iff there exists a constant  $a > 0$  and  $b \in \mathbb{R}$  such that<sup>1</sup>  
 $y_i = a \cdot x_i + b$  for any  $i = 1, \dots, n$ .

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

# Properties of the Sample Correlation Coefficient

- ▶  $\text{cor}(x, y) = \text{cor}(y, x)$ ;
- ▶ If  $\alpha > 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = \text{cor}(x, y)$
- ▶ If  $\alpha < 0$  and  $\beta \in \mathbb{R}$ , then  $\text{cor}(\alpha \cdot x + \beta, y) = -\text{cor}(x, y)$
- ▶ For any Datasets  $x, y$ ,

$$-1 \leq \rho_{xy} \leq 1;$$

- ▶  $\rho_{xy} = 1$  iff there exists a constant  $a > 0$  and  $b \in \mathbb{R}$  such that<sup>1</sup>  
 $y_i = a \cdot x_i + b$  for any  $i = 1, \dots, n$ .
- ▶  $\rho_{xy} = -1$  iff there exists a constant  $a < 0$  and  $b \in \mathbb{R}$  such  
that<sup>2</sup>  $y_i = a \cdot x_i + b$  for any  $i = 1, \dots, n$ .

---

<sup>1</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).

<sup>2</sup>Or  $x_i = a \cdot y_i + b$  for any  $i = 1, \dots, n$  (maybe for another  $a$  and  $b$ ).