ASDS Statistics, YSU, Fall 2020 Lecture 18

LCCture 10

Michael Poghosyan

31 Oct 2020

Contents

► Convergence Types of R.V. Sequences

Example: Assume X_n is a Discrete r.v. with the following PMF, defined on the same Probability Space:

$$\frac{X_n \mid 3 + \frac{1}{n^2} \mid n}{\mathbb{P}(X_n = x) \mid 1 - \frac{1}{n} \mid \frac{1}{n}.}$$

Example: Assume X_n is a Discrete r.v. with the following PMF, defined on the same Probability Space:

$$\begin{array}{c|c} X_n & 3 + \frac{1}{n^2} & n \\ \hline \mathbb{P}(X_n = x) & 1 - \frac{1}{n} & \frac{1}{n}. \end{array}$$

Which of the followings are true (use only the definitions):

- $X_n \stackrel{\mathbb{P}}{\longrightarrow} 3;$
- $\longrightarrow X_n \xrightarrow{qm} 3;$
- $X_n \xrightarrow{D} 3$?

Example: Assume

$$X_n \sim Unif\left[0, \frac{1}{n}\right]$$

and X_n are defined on the same Probability Space.

Example: Assume

$$X_n \sim Unif\left[0, \frac{1}{n}\right]$$

and X_n are defined on the same Probability Space. Which of the followings are true (use only the definitions):

- $X_n \stackrel{\mathbb{P}}{\longrightarrow} 0;$
- $\longrightarrow X_n \xrightarrow{qm} 0;$
- $\longrightarrow X_n \stackrel{D}{\longrightarrow} 0$?

Example: Show that if $X_n \sim Binom\left(n, \frac{\lambda}{n}\right)$, then $X_n \stackrel{D}{\longrightarrow} Pois(\lambda)$.


Example: Show that if $X_n \sim Binom\left(n, \frac{\lambda}{n}\right)$, then $X_n \stackrel{D}{\longrightarrow} Pois(\lambda)$.

Note: Note that when using $X_n \xrightarrow{D} Pois(\lambda)$ we mean $X_n \xrightarrow{D} X$, where $X \sim Pois(\lambda)$.

Example: Show that if $X_n \sim Binom\left(n, \frac{\lambda}{n}\right)$, then $X_n \stackrel{D}{\longrightarrow} Pois(\lambda)$.

Note: Note that when using $X_n \xrightarrow{D} Pois(\lambda)$ we mean $X_n \xrightarrow{D} X$, where $X \sim Pois(\lambda)$.

```
lambda <- 1.2; n <- 10; t <- seq(0,7, 0.1)
plot(t,pbinom(t, size = n, prob = lambda/n), type = "s", ylim = c(0,1), ylab = "CDF")
par(new = T)
plot(t, ppois(t, lambda = lambda), type = "s", col = "red", ylim = c(0,1), ylab = "CDF")</pre>
```




Example: Let $X_n \sim \textit{Unif}[0, n]$. Find the limit in Distributions of

 $Y_n = \frac{X_n}{n}$.

Example: Let $X_n \sim Unif[0, n]$. Find the limit in Distributions of $Y_n = \frac{X_n}{n}$. Let us visually show that $Y_n \stackrel{D}{\to} Y$, where $Y \sim Unif[0, 1]$.

Example: Let $X_n \sim Unif[0, n]$. Find the limit in Distributions of $Y_n = \frac{X_n}{n}$. Let us visually show that $Y_n \stackrel{D}{\to} Y$, where $Y \sim Unif[0, 1]$.

```
n <- 10000 ## We use Y_n
m <- 10000 ## No. of generated numbers
y <- runif(m, min = 0, max = n)/n
hist(y, freq = F)
abline(h = 1, col = "red", lwd = 2)</pre>
```



Theorem: Assume $X_n \overset{a.s.(P,qm,D)}{\longrightarrow} X$ and $Y_n \overset{a.s.(P,qm,D)}{\longrightarrow} Y$.

Theorem: Assume $X_n \stackrel{a.s.(P,qm,D)}{\longrightarrow} X$ and $Y_n \stackrel{a.s.(P,qm,D)}{\longrightarrow} Y$. Then

$$X_n + Y_n \xrightarrow{a.s.(P,qm)} X + Y;$$

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\triangleright X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\searrow X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $\blacktriangleright X_n \cdot Y_n \stackrel{a.s.(P)}{\longrightarrow} X \cdot Y$; and $\alpha \cdot X_n \stackrel{qm}{\longrightarrow} \alpha \cdot X$ for any constant α ;

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\triangleright X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $\blacktriangleright X_n \cdot Y_n \stackrel{a.s.(P)}{\longrightarrow} X \cdot Y$; and $\alpha \cdot X_n \stackrel{qm}{\longrightarrow} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \stackrel{a.s.(P,D)}{\longrightarrow} g(X)$

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\triangleright X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $\blacktriangleright X_n \cdot Y_n \stackrel{a.s.(P)}{\longrightarrow} X \cdot Y$; and $\alpha \cdot X_n \stackrel{qm}{\longrightarrow} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \stackrel{a.s.(P,D)}{\longrightarrow} g(X)$

Note: In the general case, if $X_n \stackrel{D}{\longrightarrow} X$ and $Y_n \stackrel{D}{\longrightarrow} Y$, then not necessarily $X_n + Y_n \stackrel{D}{\longrightarrow} X + Y$.

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\searrow X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $\blacktriangleright X_n \cdot Y_n \stackrel{a.s.(P)}{\longrightarrow} X \cdot Y$; and $\alpha \cdot X_n \stackrel{qm}{\longrightarrow} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \stackrel{a.s.(P,D)}{\longrightarrow} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.

But we have:

Theorem (Slutsky) Assume $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{P} c$, where $c \in \mathbb{R}$ is a constant. Then

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\searrow X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $\blacktriangleright X_n \cdot Y_n \stackrel{a.s.(P)}{\longrightarrow} X \cdot Y$; and $\alpha \cdot X_n \stackrel{qm}{\longrightarrow} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \stackrel{a.s.(P,D)}{\longrightarrow} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.

But we have:

Theorem (Slutsky) Assume $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{P} c$, where $c \in \mathbb{R}$ is a constant. Then

$$X_n + Y_n \xrightarrow{D} X + c;$$

Theorem: Assume $X_n \xrightarrow{a.s.(P,qm,D)} X$ and $Y_n \xrightarrow{a.s.(P,qm,D)} Y$. Then

- $\searrow X_n + Y_n \stackrel{a.s.(P,qm)}{\longrightarrow} X + Y;$
- $X_n \cdot Y_n \xrightarrow{a.s.(P)} X \cdot Y$; and $\alpha \cdot X_n \xrightarrow{qm} \alpha \cdot X$ for any constant α ;
- ▶ If $g \in C(\mathbb{R})$, then $g(X_n) \stackrel{a.s.(P,D)}{\longrightarrow} g(X)$

Note: In the general case, if $X_n \xrightarrow{D} X$ and $Y_n \xrightarrow{D} Y$, then not necessarily $X_n + Y_n \xrightarrow{D} X + Y$.


But we have:

Theorem (Slutsky) Assume $X_n \stackrel{D}{\longrightarrow} X$ and $Y_n \stackrel{P}{\longrightarrow} c$, where $c \in \mathbb{R}$ is a constant. Then

- $X_n + Y_n \xrightarrow{D} X + c;$
- $X_n \cdot Y_n \stackrel{D}{\longrightarrow} c \cdot X.$


Relationship between Convergence Types

Theorem: (Convergence Relationship Diagram)



Relationship between Convergence Types


Theorem: (Convergence Relationship Diagram)



Note: Inverse implications are not always correct.

Relationship between Convergence Types

Theorem: (Convergence Relationship Diagram)



Note: Inverse implications are not always correct. But, say, the following holds: If $X_n \xrightarrow{D} X$ and $X \equiv constant$, then $X_n \xrightarrow{P} X$ (X_n and X are defined on the same Probability space).