

# ASDS Statistics, YSU, Fall 2020

## Lecture 26

Michael Poghosyan

02 Dec 2020

# Contents

- ▶ Properties of MLE
- ▶ Confidence Intervals

## Some Notes about MLE

**Note:** The MLE does not always exist;

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Example:** Consider the Model  $Unif(a, b)$ , i.e., Uniform over the open interval  $(a, b)$ .

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Example:** Consider the Model  $Unif(a, b)$ , i.e., Uniform over the open interval  $(a, b)$ .

**Example:** Consider the Model with PDF

$$f(x|\theta) = \begin{cases} e^{\theta-x}, & x > \theta \\ 0, & x \leq \theta. \end{cases}$$

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Example:** Consider the Model  $Unif(a, b)$ , i.e., Uniform over the open interval  $(a, b)$ .

**Example:** Consider the Model with PDF

$$f(x|\theta) = \begin{cases} e^{\theta-x}, & x > \theta \\ 0, & x \leq \theta. \end{cases}$$

**Note:** The MLE is not always unique;

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Example:** Consider the Model  $Unif(a, b)$ , i.e., Uniform over the open interval  $(a, b)$ .

**Example:** Consider the Model with PDF

$$f(x|\theta) = \begin{cases} e^{\theta-x}, & x > \theta \\ 0, & x \leq \theta. \end{cases}$$

**Note:** The MLE is not always unique;

**Example:** Consider the Model  $Unif[\theta, \theta + 1]$ .

## Some Notes about MLE

**Note:** The MLE does not always exist;

**Example:** Consider the Model  $Unif(a, b)$ , i.e., Uniform over the open interval  $(a, b)$ .

**Example:** Consider the Model with PDF

$$f(x|\theta) = \begin{cases} e^{\theta-x}, & x > \theta \\ 0, & x \leq \theta. \end{cases}$$

**Note:** The MLE is not always unique;

**Example:** Consider the Model  $Unif[\theta, \theta + 1]$ .

**Note:** Sometimes it is necessary to find it numerically, say, using the NR Method



## Examples

**Example:** It is not possible to find the MLE Estimator in a closed form for  $\theta$  in the one-Parametric Cauchy Distribution  $Cauchy(\theta)$  Model. Here, the PDF of  $X \sim Cauchy(\theta)$  is given by

$$f(x|\theta) = \frac{1}{\pi(1 + (x - \theta)^2)}, \quad x \in \mathbb{R},$$

and  $\theta \in \mathbb{R}$  is called the *location parameter*.

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, as people usually use:

$$\sqrt{n} \cdot (\hat{\theta}_n^{MLE} - \theta) \xrightarrow{D} \mathcal{N}\left(0, \frac{1}{\mathcal{I}(\theta)}\right),$$

## Properties of the MLE

It can be proven that, under some regularity conditions on the Parametric Family  $\mathcal{F}_\theta$ ,

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Consistent, i.e.,

$$\hat{\theta}_n^{MLE} \xrightarrow{\mathbb{P}} \theta$$

- ▶ The MLE  $\hat{\theta}_n^{MLE}$  is Asymptotically Normal and Efficient:

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\theta)}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, as people usually use:

$$\sqrt{n} \cdot (\hat{\theta}_n^{MLE} - \theta) \xrightarrow{D} \mathcal{N}\left(0, \frac{1}{\mathcal{I}(\theta)}\right),$$

or, put in other way,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;



## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators.

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**.

## Properties of the MLE, Cont'd

Let us give an interpretation to this last property:

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right)$$

We can interpret this as:

- ▶  $\hat{\theta}_n^{MLE}$  is approximately (Asymptotically) Normal;
- ▶ The Mean of  $\hat{\theta}_n^{MLE}$  is approximately  $\theta$ , so it is Asymptotically Unbiased;
- ▶ The Variance of  $\hat{\theta}_n^{MLE}$  is approximately  $\frac{1}{n \cdot \mathcal{I}(\theta)}$ . And we know that, by CR LB, this is the possible smallest Variance we can have with Unbiased Estimators. So we interpret this as **MLE is Asymptotically Efficient**.

So, MLE is **Consistent** and **Asymptotically Efficient**. And this is why, for large Sample Size  $n$ , MLE is the Top 1 Choice, is (almost) unbeatable.

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

**Note:** This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of  $\mathcal{I}(\theta)$  we have  $\mathcal{I}(\hat{\theta}_n^{MLE})$ .



## Properties of the MLE, Cont'd

► Also,

$$\frac{\hat{\theta}_n^{MLE} - \theta}{\sqrt{\frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}}} \xrightarrow{D} \mathcal{N}(0, 1)$$

or, in more non-rigorous terms,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\hat{\theta}_n^{MLE})}\right)$$

**Note:** This is almost the above Property,

$$\hat{\theta}_n^{MLE} \stackrel{D}{\approx} \mathcal{N}\left(\theta, \frac{1}{n \cdot \mathcal{I}(\theta)}\right),$$

but, instead of  $\mathcal{I}(\theta)$  we have  $\mathcal{I}(\hat{\theta}_n^{MLE})$ .

**Note:** We will use this later, to construct an (approximate) Confidence Interval for  $\theta$  and for testing Hypotheses about  $\theta$ .

## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

## Properties of the MLE, Cont'd

- ▶ If  $\hat{\theta}$  is the MLE for  $\theta$ , then for any function  $g$ , the MLE of  $g(\theta)$  is  $g(\hat{\theta})$ , i.e.,

$$\widehat{g(\theta)}^{MLE} = g\left(\hat{\theta}^{MLE}\right).$$

This is sometimes called the **invariance Principle of MLE**.

**Example** Find the MLE for  $\sigma$  in  $\mathcal{N}(\mu, \sigma^2)$  Model.

**Solution:** OTB

## Some topics to consider by yourself

- ▶ Multivariate Normal and MLE for MVNormal
- ▶ Kullback-Leibler Divergence and its relation to MLE
- ▶ MLE for the Mixture Model, EM Algorithm
- ▶ Bayesian Estimation: MAP and Bayes Estimator

# Confidence Intervals

## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ .



## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v.

# Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v. If the True value of our Parameter is  $\theta^*$ , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess.

## Prelude

So far we have studied the Parametric Point Estimation Problem: we were assuming that our data is coming from a Parametric Family of Distribution, and our aim was to find a best guess for that Parameter.

Assume  $\theta$  is our Parameter to be Estimated, and  $\hat{\theta}$  is a good Estimator for  $\theta$ . Assume  $\hat{\theta}$  is a Continuous r.v. If the True value of our Parameter is  $\theta^*$ , then

$$\mathbb{P}(\hat{\theta} = \theta^*) = 0,$$

i.e., we will (almost) **never** be correct in our guess. Sad news!

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>,

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And we can ask questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large  $n$  needs to be to have a good estimate.

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.

## Prelude No. 2

But the good news is that even when we cannot exactly find the True value of our Parameter using  $\hat{\theta}$ , if  $\hat{\theta}$  possesses some good properties, we believe that the Estimate obtained is a good approximation/Estimate for  $\theta^*$ .

In fact, we think as

$$\theta^* = \hat{\theta} \pm \text{error},$$

where *error* is small<sup>1</sup>, or, rather, as

$$\theta^* \in (\hat{\theta} - \text{error}, \hat{\theta} + \text{error}).$$

And we can ask questions about how small is the *error* or how much sure are we in our Estimate (for the Unknown Parameter), and how large  $n$  needs to be to have a good estimate.

Here we want to develop the theory of Confidence Intervals, which will contain answers to these questions.

---

<sup>1</sup>Recall the  $\widehat{SE}$ , the Estimated Standard Error reporting story.



## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

# CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

# CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Our aim is to find an interval containing the Unknown Parameter value.

More rigorously, we want to have an interval

- ▶ which contains the True Parameter value with high Probability;
- ▶ which has the possible smallest length.

Let us state this in Mathematical terms. We will consider here only 1D case, i.e., we will assume  $\theta \in \Theta \subset \mathbb{R}$ .

# Random Intervals and CI

We start by **Random Intervals**:

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ .

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.



# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.

**Example:** Assume  $X \sim \text{Pois}(2.3)$ . Then

$$[X, X + 2)$$

is a Random Interval.

# Random Intervals and CI

We start by **Random Intervals**:

Assume  $X$  and  $Y$  are two r.v. on the same probability space and  $X \leq Y$  a.s., that is,  $\mathbb{P}(X > Y) = 0$ . We will call intervals of the form  $[X, Y]$  or  $(X, Y)$  or  $(X, Y]$  or  $[X, Y)$  or  $(X, +\infty), \dots$ , **random intervals**.

**Example:** Assume  $X \sim \text{Pois}(2.3)$ . Then

$$[X, X + 2)$$

is a Random Interval.

**Example:** Let  $X_1, X_2, \dots, X_n$  are IID r.v.s. Then

$$(\bar{X} - 0.1, \bar{X} + 0.1)$$

is a Random Interval.

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

# CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take  $\alpha \in (0, 1)$ , and call the number  $1 - \alpha$  the **confidence level**.

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take  $\alpha \in (0, 1)$ , and call the number  $1 - \alpha$  the **confidence level**.

**Problem:** Using the Random Sample, construct an interval containing the Unknown Parameter value with Probability not less than  $1 - \alpha$ .

## CI Problem Setting

Assume we have a Random Sample from a Parametric Model  $\mathcal{F}_\theta$ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

We take  $\alpha \in (0, 1)$ , and call the number  $1 - \alpha$  the **confidence level**.

**Problem:** Using the Random Sample, construct an interval containing the Unknown Parameter value with Probability not less than  $1 - \alpha$ .

The usual values of the confidence level are 90%, 95%, 99%, so the usual values of  $\alpha$  are 0.1, 0.05 and 0.01.

**Definition:** Assume  $0 < \alpha < 1$ , and let  $L = L(x_1, \dots, x_n, \alpha)$ ,  $U = U(x_1, \dots, x_n, \alpha)$  be two functions with  $L(x_1, \dots, x_n, \alpha) \leq U(x_1, \dots, x_n, \alpha)$  for all  $(x_1, \dots, x_n, \alpha)$ .

**Definition:** Assume  $0 < \alpha < 1$ , and let  $L = L(x_1, \dots, x_n, \alpha)$ ,  $U = U(x_1, \dots, x_n, \alpha)$  be two functions with  $L(x_1, \dots, x_n, \alpha) \leq U(x_1, \dots, x_n, \alpha)$  for all  $(x_1, \dots, x_n, \alpha)$ . The random interval

$$(L, U) = \left( L(X_1, \dots, X_n, \alpha), U(X_1, \dots, X_n, \alpha) \right)$$

is called a **confidence interval (or confidence interval estimator) for  $\theta$  of confidence level  $1 - \alpha$** , if for any  $\theta \in \Theta$ ,

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha.$$



In the case we have a realization/observation of  $X_1, \dots, X_n$ , say,  $x_1, \dots, x_n$ , then the interval

$$\left( L(x_1, \dots, x_n, \alpha), U(x_1, \dots, x_n, \alpha) \right)$$

will be an **interval estimate for  $\theta$  for the confidence level  $(1 - \alpha)$** .

# CI

In the case we have a realization/observation of  $X_1, \dots, X_n$ , say,  $x_1, \dots, x_n$ , then the interval

$$\left( L(x_1, \dots, x_n, \alpha), U(x_1, \dots, x_n, \alpha) \right)$$

will be an **interval estimate for  $\theta$  for the confidence level  $(1 - \alpha)$** .

Going back to our CI, CI of the confidence level  $1 - \alpha$  is a Random Interval that contains  $\theta$  in more than  $(1 - \alpha) \cdot 100\%$  of cases.

## CI, Interpretation

**Note:** It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

$\theta$  is not our r.v.,  $\theta$  is our unknown constant Parameter, so we do not read this as “with high Probability,  $\theta$  is in  $(L, U)$ ”.

---

## CI, Interpretation

**Note:** It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

$\theta$  is not our r.v.,  $\theta$  is our unknown constant Parameter, so we do not read this as “with high Probability,  $\theta$  is in  $(L, U)$ ”. Instead, the Interval  $(L, U)$  is Random, so we read this as **with high Probability, the interval  $(L, U)$  will contain  $\theta$ .**

---

<sup>2</sup>But not different  $\theta$ -s !!

# CI, Interpretation

**Note:** It is important to understand, that in the CI definition

$$\mathbb{P}(L < \theta < U) \geq 1 - \alpha$$

$\theta$  is not our r.v.,  $\theta$  is our unknown constant Parameter, so we do not read this as “with high Probability,  $\theta$  is in  $(L, U)$ ”. Instead, the Interval  $(L, U)$  is Random, so we read this as **with high Probability, the interval  $(L, U)$  will contain  $\theta$ .**

So, if we will have/generate different observations, we will have different Intervals<sup>2</sup>  $(L, U)$ , and we want to have that most of the time that interval contains our unknown Parameter value.

---

<sup>2</sup>But not different  $\theta$ -s !!

## CI, R Simulation

**Example:** Consider an example: our Model is  $Exp(\lambda)$ , and we have an observation from it. Let us take a Random Sample for the general case:  $X_1, X_2, \dots, X_n$  from  $Exp(\lambda)$ .

## CI, R Simulation

**Example:** Consider an example: our Model is  $Exp(\lambda)$ , and we have an observation from it. Let us take a Random Sample for the general case:  $X_1, X_2, \dots, X_n$  from  $Exp(\lambda)$ . Then good **Estimator** for  $\lambda$  is

## CI, R Simulation

**Example:** Consider an example: our Model is  $Exp(\lambda)$ , and we have an observation from it. Let us take a Random Sample for the general case:  $X_1, X_2, \dots, X_n$  from  $Exp(\lambda)$ . Then good **Estimator** for  $\lambda$  is

$$\hat{\lambda} = \frac{1}{\overline{X}}.$$



## CI, R Simulation

**Example:** Consider an example: our Model is  $Exp(\lambda)$ , and we have an observation from it. Let us take a Random Sample for the general case:  $X_1, X_2, \dots, X_n$  from  $Exp(\lambda)$ . Then good **Estimator** for  $\lambda$  is

$$\hat{\lambda} = \frac{1}{\bar{X}}.$$

Now, let us take as CI

## CI, R Simulation

**Example:** Consider an example: our Model is  $Exp(\lambda)$ , and we have an observation from it. Let us take a Random Sample for the general case:  $X_1, X_2, \dots, X_n$  from  $Exp(\lambda)$ . Then good **Estimator** for  $\lambda$  is

$$\hat{\lambda} = \frac{1}{\bar{X}}.$$


Now, let us take as CI

$$\left( \frac{1}{\bar{X}} - 0.1, \frac{1}{\bar{X}} + 0.1 \right)$$

and do some simulations:

# CI, R Simulation

Exponential Model, CI,  $(1/\text{mean} - 0.1, 1/\text{mean} + 0.1)$



## CI, R Simulation, Code

*#CI Idea, Exponential Model*

```
lambda <- 0.41
```

```
conf.level <- 0.95; a = 1 - conf.level
```

```
sample.size <- 50; no.of.intervals <- 100
```

```
epsilon <- 0.1
```

```
plot.new()
```

```
plot.window(xlim = c(0,no.of.intervals), ylim = c(0.2,0.8))
```

```
axis(1); axis(2)
```

```
title("Exponential Model, CI, (1/mean - 0.1, 1/mean + 0.1)")
```

```
for(i in 1:no.of.intervals){
```

```
  x <- rexp(sample.size, rate = lambda)
```

```
  lo <- 1/mean(x) - epsilon; up <- 1/mean(x) + epsilon
```

```
  if(lo > lambda || up < lambda){
```

```
    segments(c(i), c(lo), c(i), c(up), col = "red")
```

```
  }
```

```
  else{
```

```
    segments(c(i), c(lo), c(i), c(up))
```

```
  }
```

```
}
```

```
abline(h = lambda, lwd = 2, col = "blue")
```

# Methods to obtain Confidence Intervals

We will consider several methods to construct CIs:

- ▶ Chebyshev Inequality Based;
- ▶ Pivotal Quantity Based

## CI for the Mean, Variance is given, Cheby Method

**Example:** Assume  $X_1, X_2, \dots, X_n$  are Independent r.v. with the same Mean  $\mathbb{E}(X_k) = \mu$  and the same Variance  $\text{Var}(X_k) = \sigma^2$ .

## CI for the Mean, Variance is given, Cheby Method

**Example:** Assume  $X_1, X_2, \dots, X_n$  are Independent r.v. with the same Mean  $\mathbb{E}(X_k) = \mu$  and the same Variance  $\text{Var}(X_k) = \sigma^2$ . Assume  $\sigma^2$  is known, and  $\mu$  is unknown, and we want to estimate it using the Sample.

## CI for the Mean, Variance is given, Cheby Method

**Example:** Assume  $X_1, X_2, \dots, X_n$  are Independent r.v. with the same Mean  $\mathbb{E}(X_k) = \mu$  and the same Variance  $\text{Var}(X_k) = \sigma^2$ . Assume  $\sigma^2$  is known, and  $\mu$  is unknown, and we want to estimate it using the Sample. The problem is to construct a CI for  $\mu$  of given Confidence Level  $1 - \alpha$ .



## CI for the Mean, Variance is given, Cheby Method

**Example:** Assume  $X_1, X_2, \dots, X_n$  are Independent r.v. with the same Mean  $\mathbb{E}(X_k) = \mu$  and the same Variance  $\text{Var}(X_k) = \sigma^2$ . Assume  $\sigma^2$  is known, and  $\mu$  is unknown, and we want to estimate it using the Sample. The problem is to construct a CI for  $\mu$  of given Confidence Level  $1 - \alpha$ .

By the Chebyshev inequality method, we can obtain that the interval

$$\left( \bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}}, \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}} \right)$$

is a CI for  $\mu$  of Confidence Level  $1 - \alpha$ .

## CI for the Mean, Variance is given, Cheby Method

**Example:** Assume  $X_1, X_2, \dots, X_n$  are Independent r.v. with the same Mean  $\mathbb{E}(X_k) = \mu$  and the same Variance  $\text{Var}(X_k) = \sigma^2$ . Assume  $\sigma^2$  is known, and  $\mu$  is unknown, and we want to estimate it using the Sample. The problem is to construct a CI for  $\mu$  of given Confidence Level  $1 - \alpha$ .

By the Chebyshev inequality method, we can obtain that the interval

$$\left( \bar{X} - \frac{\sigma}{\sqrt{n \cdot \alpha}}, \bar{X} + \frac{\sigma}{\sqrt{n \cdot \alpha}} \right)$$

is a CI for  $\mu$  of Confidence Level  $1 - \alpha$ .

**Note:** Here

$$\frac{\sigma}{\sqrt{n \cdot \alpha}}$$

is called the **Margin of Error** (for the Interval Estimate of  $\mu$ , given  $\sigma^2$ ).

## Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

## Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

The CI length obtained above is

$$\frac{2\sigma}{\sqrt{n \cdot \alpha}}.$$

**Note:** If we increase  $n$ , the CI gets narrower. This is intuitive: if we collect more data, we can estimate the parameter more precisely, we can enclose it in a smaller length interval.

## Some Notes

Two notes about the obtained CI - in fact, these notes will work also for other cases too:

The CI length obtained above is

$$\frac{2\sigma}{\sqrt{n \cdot \alpha}}.$$

**Note:** If we increase  $n$ , the CI gets narrower. This is intuitive: if we collect more data, we can estimate the parameter more precisely, we can enclose it in a smaller length interval.

**Note:** If we increase the Confidence Level, i.e., if we decrease  $\alpha$ , then the length of CI increases. This is intuitive too: if we want to be more sure where our unknown Parameter is lying, we will get a larger interval.

## CI for the Proportion, Cheby Method

**Example:** Now, let us construct a CI of CLevel  $1 - \alpha$  for  $p$  in the *Bernoulli*( $p$ ) Model.

## CI for the Proportion, Cheby Method

**Example:** Now, let us construct a CI of CLevel  $1 - \alpha$  for  $p$  in the *Bernoulli*( $p$ ) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

## CI for the Proportion, Cheby Method

**Example:** Now, let us construct a CI of CLevel  $1 - \alpha$  for  $p$  in the *Bernoulli*( $p$ ) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

The CI for  $p$  by Chebyshev Inequality will be

$$\left( \bar{X} - \frac{1}{2\sqrt{n \cdot \alpha}}, \bar{X} + \frac{1}{2\sqrt{n \cdot \alpha}} \right)$$

is a CI for  $p$  of level  $1 - \alpha$ .



## CI for the Proportion, Cheby Method

**Example:** Now, let us construct a CI of CLevel  $1 - \alpha$  for  $p$  in the *Bernoulli*( $p$ ) Model. We assume we have a Random Sample

$$X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p).$$

The CI for  $p$  by Chebyshev Inequality will be

$$\left( \bar{X} - \frac{1}{2\sqrt{n \cdot \alpha}}, \bar{X} + \frac{1}{2\sqrt{n \cdot \alpha}} \right)$$


is a CI for  $p$  of level  $1 - \alpha$ .

**Note:** Here

$$\frac{1}{2\sqrt{n \cdot \alpha}}$$

is called the **Margin of Error** (for the Interval Estimate of  $p$ ).


# CI for Bernoulli, R Simulation



Sample Size = 50,  $CL = 95\%$


# CI for Bernoulli, R Simulation

Bernoulli Model, CI by Cheby



Sample Size = 150,  $CL = 95\%$


# CI for Bernoulli, R Simulation



Sample Size = 150,  $CL = 99\%$

# CI for Bernoulli, R Simulation

Bernoulli Model, CI by Cheby



Sample Size = 250,  $CL = 90\%$

## CI, R Simulation, Code

```
#CI Idea, Bernoulli Model
```

```
p <- 0.345
```

```
conf.level <- 0.9; a = 1 - conf.level
```

```
sample.size <- 250; no.of.intervals <- 150
```

```
ME <- 1/(2*sqrt(sample.size*a)) #Margin of Error
```

```
plot.new()
```

```
plot.window(xlim = c(0,no.of.intervals), ylim = c(0,1))
```

```
axis(1); axis(2)
```

```
title("Bernoulli Model, CI by Cheby")
```

```
for(i in 1:no.of.intervals){
```

```
  x <- rbinom(sample.size, size = 1, prob = p)
```

```
  lo <- mean(x) - ME
```

```
  up <- mean(x) + ME
```

```
  if(lo > p || up < p){
```

```
    segments(c(i), c(lo), c(i), c(up), col = "red")
```

```
  }
```

```
  else{
```

```
    segments(c(i), c(lo), c(i), c(up))
```

```
  }
```

```
}
```

```
abline(h = p, lwd = 2, col = "blue")
```

## Examples

**Example:** Assume we are interested in the proportion of smokers in AUA. We ask 120 persons at AUA and learn that 55 of them are smokers. Construct a CI for the proportion of smokers in AUA of 95% confidence level.

**Solution:** OTB

## Examples

**Example:** Assume we are interested in the proportion of smokers in AUA. We ask 120 persons at AUA and learn that 55 of them are smokers. Construct a CI for the proportion of smokers in AUA of 95% confidence level.

**Solution:** OTB

**Example:** Continuing the above Example: now assume we want to find that Proportion within the Error Margin 0.1, with the CL 95%. At least, how many persons at AUA we need to ask?

**Solution:** OTB