

ASDS Statistics, YSU, Fall 2020

Lecture 24

Michael Poghosyan

25 Nov 2020

Contents

- ▶ The Method of Moments

Methods to find (good) Estimators

The Problem

Here, in this Section, our Problem will be the following:

The Problem

Here, in this Section, our Problem will be the following:

We are given a Random Sample from a Parametric Family of Distributions,

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

The Problem

Here, in this Section, our Problem will be the following:

We are given a Random Sample from a Parametric Family of Distributions,

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Problem: The Problem is to find/construct a good Estimator for θ , using our Random Sample.

The Method of Moments

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$. The second Moment of \mathcal{F}_θ is $\mathbb{E}(X^2)$ etc.

Theoretical Moments

First we define the Theoretical and Sample (Empirical) Moments:

Assume we consider the Parametric Family of Distributions \mathcal{F}_θ , $\theta \in \Theta$. Let $X \sim \mathcal{F}_\theta$.

Definition: The **Theoretical Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ (or the r.v. X) is $\mathbb{E}(X^k)$.

Example: Say, the first Moment of \mathcal{F}_θ is just the Mean of X , $\mathbb{E}(X)$. The second Moment of \mathcal{F}_θ is $\mathbb{E}(X^2)$ etc.

Note: Note that, in general, the Theoretical Moments of \mathcal{F}_θ are functions of θ .

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Example: Say, the second order Empirical Moment is

$$\frac{(X_1)^2 + (X_2)^2 + \dots + (X_n)^2}{n}.$$

Empirical (Sample) Moments

Now assume we have a Random Sample from \mathcal{F}_θ :

$$X_1, X_2, \dots, X_n \sim \mathcal{F}_\theta, \quad \theta \in \Theta.$$

Definition: The **Empirical (Sample) Moment** of order $k \in \mathbb{N}$ of the Distribution \mathcal{F}_θ is

$$\frac{(X_1)^k + (X_2)^k + \dots + (X_n)^k}{n}.$$

Example: Say, the second order Empirical Moment is

$$\frac{(X_1)^2 + (X_2)^2 + \dots + (X_n)^2}{n}.$$

Note: The Empirical Moment is independent of the Parameter θ , it is just a Statistics, it is a function of X_1, X_2, \dots, X_n .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ .

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**. And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**. And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

If this is giving the value for θ , then you have the MoM Estimator.

The Method of Moments, 1D case

Now, the idea of the Method of Moments (MoM) is to match the corresponding Theoretical and Empirical Moments.

First we consider the case when our Parameter θ is 1D, i.e., $\theta \in \Theta \subset \mathbb{R}$. In that case, we need just one equation to find θ .

MoM suggests: Solve the Equation

1-st order Theoretical Moment = 1-st order Empirical Moment,

to find (express) θ . The value of θ , expressed by the Random Sample, is called **the MoM Estimator**. And if this will not give a value for θ , solve instead

2-nd order Theoretical Moment = 2-nd order Empirical Moment.

If this is giving the value for θ , then you have the MoM Estimator. Otherwise, continue with the 3-rd Moments, then with 4-th Moments, etc.

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$.

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$. Find the MoM Estimator and MoM Estimate for θ .

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for p in the *Bernoulli*(p) Model.

Solution: OTB

Example: Assume we have an observation

$$0, 1, 1, 2, 1, 0, 0, 1, 1$$

from the following Model:

X	0	1	2
$\mathbb{P}(X = x)$	$\frac{\theta}{10}$	$\frac{\theta}{5}$	$1 - \frac{3\theta}{10}$

where $\theta \in [0, \frac{10}{3}]$. Find the MoM Estimator and MoM Estimate for θ .

Solution: OTB

Example: Find the MoM Estimator for λ in the *Exp*(λ) Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for θ in the $Unif[0, \theta]$ Model.

Solution: OTB

Examples, MoM

Example: Find the MoM Estimator for θ in the $Unif[0, \theta]$ Model.

Solution: OTB

Example: Find the MoM Estimator for θ in the $Unif[-\theta, \theta]$ Model.

Solution: OTB

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 .

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The LHS of these equations, in general, will depend on θ_1 and θ_2 .

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The LHS of these equations, in general, will depend on θ_1 and θ_2 . If we can find from here θ_1 and θ_2 , we take them as the MoM Estimators.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The LHS of these equations, in general, will depend on θ_1 and θ_2 . If we can find from here θ_1 and θ_2 , we take them as the MoM Estimators. Otherwise, if we cannot express from this system θ_1 or θ_2 , then we try to solve another pair of this kind of equations, using the possible smallest order Moments.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The LHS of these equations, in general, will depend on θ_1 and θ_2 . If we can find from here θ_1 and θ_2 , we take them as the MoM Estimators. Otherwise, if we cannot express from this system θ_1 or θ_2 , then we try to solve another pair of this kind of equations, using the possible smallest order Moments. Say, if the 1st Moment equation is not giving a result, solve the system with 2nd and 3rd Moments.

The Method of Moments, m -D case

Now assume that our Parameter θ is m -Dimensional, or, in other way, we have m Parameters to Estimate. Let me give the 2D case, and you will be able to extend to the m -D case.

So assume $\theta = (\theta_1, \theta_2)$. Then we need 2 Equations to find θ_1 and θ_2 . The MoM says: first try to solve the following system:

$$\begin{cases} \text{1-st order Theoretical Moment} = \text{1-st order Empirical Moment} \\ \text{2-nd order Theoretical Moment} = \text{2-nd order Empirical Moment} \end{cases}$$

The LHS of these equations, in general, will depend on θ_1 and θ_2 . If we can find from here θ_1 and θ_2 , we take them as the MoM Estimators. Otherwise, if we cannot express from this system θ_1 or θ_2 , then we try to solve another pair of this kind of equations, using the possible smallest order Moments. Say, if the 1st Moment equation is not giving a result, solve the system with 2nd and 3rd Moments. Or, if in the initial system 2nd order Moment equation is not saying anything, use 1st and 3rd Moments, etc.

Example

Example: Find the MoM Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Example

Example: Find the MoM Estimator for (μ, σ^2) in the $\mathcal{N}(\mu, \sigma^2)$ Model.

Solution: OTB

Example: Find the MoM Estimator for (a, b) in the $Unif[a, b]$ Model.

Solution: OTB

Example

Example: Let us do an experiment in **R**, concerning the last example:

```
a <- 2.5; b <- 3.24
x <- runif(10, min = a, max = b)
x.bar <- mean(x)
z <- sqrt(3* (mean(x^2) - x.bar^2))
a.hat.MoM<- x.bar - z
b.hat.MoM <- x.bar + z
c(a.hat.MoM, b.hat.MoM)
```

```
## [1] 2.410272 3.052920
```

Example

Example: Let us do an experiment in **R**, concerning the last example:

```
a <- 2.5; b <- 3.24
x <- runif(10, min = a, max = b)
x.bar <- mean(x)
z <- sqrt(3* (mean(x^2) - x.bar^2))
a.hat.MoM<- x.bar - z
b.hat.MoM <- x.bar + z
c(a.hat.MoM, b.hat.MoM)
```

```
## [1] 2.410272 3.052920
```

Of course, we can just take $\hat{a} = X_{(1)}$ and $\hat{b} = X_{(n)}$:

```
c(min(x), max(x))
```

```
## [1] 2.515333 3.020296
```

Notes

Note: If you do not want to obtain the *MoM Estimator*, but just the *MoM Estimate*, just plug the Observations into the Empirical Moments, and then solve the Moments Equality Equations.

Notes

Note: If you do not want to obtain the *MoM Estimator*, but just the *MoM Estimate*, just plug the Observations into the Empirical Moments, and then solve the Moments Equality Equations.

Note: Sometimes we are interested not in the Estimate of the Parameter itself, but in the Estimate for some function of that Parameter.

Notes

Note: If you do not want to obtain the *MoM Estimator*, but just the *MoM Estimate*, just plug the Observations into the Empirical Moments, and then solve the Moments Equality Equations.

Note: Sometimes we are interested not in the Estimate of the Parameter itself, but in the Estimate for some function of that Parameter. Say, we want to Estimate $h(\theta)$, where θ is our unknown Parameter.

Notes

Note: If you do not want to obtain the *MoM Estimator*, but just the *MoM Estimate*, just plug the Observations into the Empirical Moments, and then solve the Moments Equality Equations.

Note: Sometimes we are interested not in the Estimate of the Parameter itself, but in the Estimate for some function of that Parameter. Say, we want to Estimate $h(\theta)$, where θ is our unknown Parameter. Then one of the approaches is the Plug-in Principle: find an Estimator $\hat{\theta}$ for θ , say, using the MoM, and then plug that in h , to obtain $h(\hat{\theta})$ as an Estimator for $h(\theta)$.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous. Then the MoM Estimator for θ will be Consistent.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous. Then the MoM Estimator for θ will be Consistent.

Indeed, to find the MoM Estimator for θ , we need to solve

$$e(\theta) = \bar{X}_n,$$

the solution of which we denote by $\hat{\theta}_n$.

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous. Then the MoM Estimator for θ will be Consistent.

Indeed, to find the MoM Estimator for θ , we need to solve

$$e(\theta) = \overline{X}_n,$$

the solution of which we denote by $\hat{\theta}_n$. This gives

$$\hat{\theta}_n = e^{-1}(\overline{X}_n).$$

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous. Then the MoM Estimator for θ will be Consistent.

Indeed, to find the MoM Estimator for θ , we need to solve

$$e(\theta) = \bar{X}_n,$$

the solution of which we denote by $\hat{\theta}_n$. This gives

$$\hat{\theta}_n = e^{-1}(\bar{X}_n).$$

Now, by the WLLN, $\bar{X}_n \xrightarrow{\mathbb{P}} \mathbb{E}(X_1) = e(\theta)$,

Consistency of the MoM Estimator

Let's consider the case when we use the 1st order Moments.

Assume that for the Model \mathcal{F}_θ , the Expectation $\mathbb{E}_\theta(X)$ ($X \sim \mathcal{F}_\theta$), as a function of θ , has a Continuous Inverse function. Say, if we denote $e(\theta) = \mathbb{E}_\theta(X)$, then we assume that the function $e^{-1}(\cdot)$ is continuous. Then the MoM Estimator for θ will be Consistent.

Indeed, to find the MoM Estimator for θ , we need to solve

$$e(\theta) = \bar{X}_n,$$

the solution of which we denote by $\hat{\theta}_n$. This gives

$$\hat{\theta}_n = e^{-1}(\bar{X}_n).$$

Now, by the WLLN, $\bar{X}_n \xrightarrow{\mathbb{P}} \mathbb{E}(X_1) = e(\theta)$, so

$$\hat{\theta}_n = e^{-1}(\bar{X}_n) \xrightarrow{\mathbb{P}} e^{-1}(e(\theta)) = \theta.$$