

# ASDS Statistics, YSU, Fall 2020

## Lecture 19

Michael Poghosyan

04 Nov 2020

# Contents

- ▶ Limit Theorems

## Note

**Note:** Mostly, in our course, we will deal with the following type of sequences of r.v.s:

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n},$$

## Note

**Note:** Mostly, in our course, we will deal with the following type of sequences of r.v.s:

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n},$$

and to calculate the limit of this sequence  $\bar{X}_1, \bar{X}_2, \dots, \bar{X}_n, \dots$ , we will use our famous Limit Theorems: LLN and CLT.

# Limit Theorems

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s.

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution.

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide.



## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

- ▶  $X_n$ -s are independent.

## Sequence of IID r.v.

Assume  $X_n$  is a sequence of **Independent, Identically Distributed (IID)** r.v.s. This means that:

- ▶ All  $X_n$ -s have the same Distribution. In particular, all numerical partial characteristics of  $X_n$  coincide. In particular,

$$\mathbb{E}(X_1) = \mathbb{E}(X_2) = \dots = \mathbb{E}(X_n) = \dots,$$

$$\text{Var}(X_1) = \text{Var}(X_2) = \dots = \text{Var}(X_n) = \dots$$

We will use this many-many-many-many-... times.

- ▶  $X_n$ -s are independent. Say, in particular,

$$\text{Var}(X_1 + X_2 + \dots + X_n) = \text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_n) = n \cdot \text{Var}(X_1).$$

## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs.

## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$



## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

**Note:** Not an easy task to find the Distribution of  $S_n$  or  $\overline{X}_n$ .

## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

**Note:** Not an easy task to find the Distribution of  $S_n$  or  $\bar{X}_n$ . Even for  $n = 2$ .

## Idea of the LLN and CLT

Assume we have a sequence  $X_n$  of IID rvs. We want to study the behavior of either the sum

$$S_n = X_1 + X_2 + \dots + X_n$$

or the average

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

**Note:** Not an easy task to find the Distribution of  $S_n$  or  $\overline{X}_n$ . Even for  $n = 2$ . We need Convolutions!

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) =$$

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) =$$

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

*so the mean of the means is the mean ☺, and*

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) =$$



## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) =$$

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

The interpretation of  $\mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1)$  and  $\text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}$ :

## What we know about $S_n$ and $\bar{X}_n$

Some important known facts about  $S_n$  and  $\bar{X}_n$  in the general case:

$$\mathbb{E}(S_n) = n \cdot \mathbb{E}(X_1), \quad \mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1);$$

so *the mean of the means is the mean* ☺, and

$$\text{Var}(S_n) = n \cdot \text{Var}(X_1), \quad \text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}.$$

The last property is the mathematical proof of the effectiveness of “7 angam chapir, mek angam ktrir” ☺

The interpretation of  $\mathbb{E}(\bar{X}_n) = \mathbb{E}(X_1)$  and  $\text{Var}(\bar{X}_n) = \frac{\text{Var}(X_1)}{n}$ : the values of  $\bar{X}_n$  are centered at  $\mathbb{E}(X_1)$  and are becoming more and more concentrated around that number as  $n$  increases.

## Some Particular results about $S_n$ and $\overline{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\overline{X}_n$ :

## Some Particular results about $S_n$ and $\bar{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\bar{X}_n$ :

- ▶ If  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

## Some Particular results about $S_n$ and $\bar{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\bar{X}_n$ :

- If  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim$$



## Some Particular results about $S_n$ and $\bar{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\bar{X}_n$ :

- If  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

## Some Particular results about $S_n$ and $\bar{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\bar{X}_n$ :

- If  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

- If  $X_k \sim \text{Pois}(\lambda)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

## Some Particular results about $S_n$ and $\overline{X}_n$

The following are important cases when we can find exactly the Distribution of  $S_n$  and/or  $\overline{X}_n$ :

- ▶ If  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \mathcal{N}(n \cdot \mu, n \cdot \sigma^2)$$

and

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

- ▶ If  $X_k \sim \text{Pois}(\lambda)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Pois}(n \cdot \lambda).$$

## Some Particular results about $S_n$ and $\bar{X}_n$

- ▶ If  $X_k \sim \text{Bernoulli}(p)$ ,  $k = 1, \dots, n$  are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

## Some Particular results about $S_n$ and $\bar{X}_n$

- ▶ If  $X_k \sim \text{Bernoulli}(p)$ ,  $k = 1, \dots, n$  are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

## Some Particular results about $S_n$ and $\overline{X}_n$

- ▶ If  $X_k \sim \text{Bernoulli}(p)$ ,  $k = 1, \dots, n$  are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

- ▶ If  $X_k \sim \text{Binom}(m, p)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim$$

## Some Particular results about $S_n$ and $\overline{X}_n$

- ▶ If  $X_k \sim \text{Bernoulli}(p)$ ,  $k = 1, \dots, n$  are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n, p);$$

- ▶ If  $X_k \sim \text{Binom}(m, p)$ ,  $k = 1, \dots, n$ , are Independent, then

$$S_n = X_1 + \dots + X_n \sim \text{Binom}(n \cdot m, p).$$

## The Weak LLN

Now, what can be said about  $S_n$  and  $\overline{X}_n$  in the general case?



## The Weak LLN

Now, what can be said about  $S_n$  and  $\overline{X}_n$  in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

## The Weak LLN

Now, what can be said about  $S_n$  and  $\bar{X}_n$  in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

### The Weak Law of Large Numbers, WLLN:

If  $X_1, X_2, \dots, X_n$  are IID, with finite  $\mathbb{E}(X_1)$  and Variance  $\text{Var}(X_1)$ , then

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

## The Weak LLN

Now, what can be said about  $S_n$  and  $\overline{X}_n$  in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

### The Weak Law of Large Numbers, WLLN:

If  $X_1, X_2, \dots, X_n$  are IID, with finite  $\mathbb{E}(X_1)$  and Variance  $\text{Var}(X_1)$ , then

$$\overline{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

i.e., for any  $\varepsilon > 0$ ,

$$\mathbb{P} \left( \left| \frac{X_1 + X_2 + \dots + X_n}{n} - \mathbb{E}(X_1) \right| \geq \varepsilon \right) \rightarrow 0, \quad n \rightarrow +\infty.$$

## The Weak LLN

Now, what can be said about  $S_n$  and  $\bar{X}_n$  in the general case? LLN and CLT help us in this matter, they describe the *asymptotic* properties of these guys:

### The Weak Law of Large Numbers, WLLN:

If  $X_1, X_2, \dots, X_n$  are IID, with finite  $\mathbb{E}(X_1)$  and Variance  $\text{Var}(X_1)$ , then

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\mathbb{P}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

i.e., for any  $\varepsilon > 0$ ,

$$\mathbb{P} \left( \left| \frac{X_1 + X_2 + \dots + X_n}{n} - \mathbb{E}(X_1) \right| \geq \varepsilon \right) \rightarrow 0, \quad n \rightarrow +\infty.$$

**Note:** This means that for any  $\varepsilon > 0$ , the chances that  $\bar{X}_n$  is far from  $\mathbb{E}(X_1)$  more than  $\varepsilon$ , is very small, if  $n$  is large.

# The Strong LLN

## The Strong Law of Large Numbers, SLLN, Kolmogorov

If  $X_1, X_2, \dots, X_n$  are IID, with finite  $\mathbb{E}(|X_1|)$ , then

$$\frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{a.s.} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

# The Strong LLN

## The Strong Law of Large Numbers, SLLN, Kolmogorov

If  $X_1, X_2, \dots, X_n$  are IID, with finite  $\mathbb{E}(|X_1|)$ , then


$$\frac{X_1 + X_2 + \dots + X_n}{n} \xrightarrow{\text{a.s.}} \mathbb{E}(X_1), \quad n \rightarrow +\infty,$$

that is,

$$\mathbb{P} \left( \lim_{n \rightarrow +\infty} \frac{X_1 + X_2 + \dots + X_n}{n} = \mathbb{E}(X_1) \right) = 1.$$

# Visualization of the LLN

```
set.seed(111); n <- 1000; expect <- 0.6  
X <- rbinom(n, 1, expect)  
S <- cumsum(X); p <- S/(1:n)  
plot(p, type = "l")  
abline(expect, 0, col = "red", lwd = 2)
```



## Supplements, LLN

Sometimes we are required to calculate limits of the form:

$$\lim_{n \rightarrow +\infty} \frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n}$$

in the *Probability* or *a.s.* sense, for some nice function  $g$ .



## Supplements, LLN

Sometimes we are required to calculate limits of the form:

$$\lim_{n \rightarrow +\infty} \frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n}$$

in the *Probability* or *a.s.* sense, for some nice function  $g$ . Clearly, under the condition that  $\mathbb{E}(g(X_1))$  and  $\text{Var}(g(X_1))$  are finite, or  $\mathbb{E}(|g(X_1)|) < +\infty$ , we will have

$$\frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n} \xrightarrow{\mathbb{P}, a.s.} \mathbb{E}(g(X_1)), \quad n \rightarrow +\infty.$$

## Supplements, LLN

Sometimes we are required to calculate limits of the form:

$$\lim_{n \rightarrow +\infty} \frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n}$$

in the *Probability* or *a.s.* sense, for some nice function  $g$ . Clearly, under the condition that  $\mathbb{E}(g(X_1))$  and  $\text{Var}(g(X_1))$  are finite, or  $\mathbb{E}(|g(X_1)|) < +\infty$ , we will have

$$\frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n} \xrightarrow{\mathbb{P}, a.s.} \mathbb{E}(g(X_1)), \quad n \rightarrow +\infty.$$

Say, for example,

$$\frac{X_1^2 + X_2^2 + \dots + X_n^2}{n} \xrightarrow{\mathbb{P}, a.s.}$$

## Supplements, LLN

Sometimes we are required to calculate limits of the form:

$$\lim_{n \rightarrow +\infty} \frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n}$$

in the *Probability* or *a.s.* sense, for some nice function  $g$ . Clearly, under the condition that  $\mathbb{E}(g(X_1))$  and  $\text{Var}(g(X_1))$  are finite, or  $\mathbb{E}(|g(X_1)|) < +\infty$ , we will have

$$\frac{g(X_1) + g(X_2) + \dots + g(X_n)}{n} \xrightarrow{\mathbb{P}, a.s.} \mathbb{E}(g(X_1)), \quad n \rightarrow +\infty.$$

Say, for example,

$$\frac{X_1^2 + X_2^2 + \dots + X_n^2}{n} \xrightarrow{\mathbb{P}, a.s.} \mathbb{E}(X_1^2), \quad n \rightarrow +\infty.$$

# CLT

The LLN says that the values of  $\bar{X}_n$  are concentrated around  $\mathbb{E}(X_1)$ .

# CLT

The LLN says that the values of  $\bar{X}_n$  are concentrated around  $\mathbb{E}(X_1)$ . But it is not giving us an idea about how the values are distributed around that mean value.

# CLT

The LLN says that the values of  $\bar{X}_n$  are concentrated around  $\mathbb{E}(X_1)$ . But it is not giving us an idea about how the values are distributed around that mean value. CLT helps us in this regard.

# CLT

The LLN says that the values of  $\bar{X}_n$  are concentrated around  $\mathbb{E}(X_1)$ . But it is not giving us an idea about how the values are distributed around that mean value. CLT helps us in this regard.

To give the general idea of the CLT, let us use the following transform: for a r.v.  $X$ , let us denote

$$Z = \text{Standardize}(X) = \frac{X - \mathbb{E}(X)}{\sqrt{\text{Var}(X)}} = \frac{X - \mathbb{E}(X)}{SD(X)},$$

the Standardization (normalization, scaling) of  $X$ .

# CLT

The LLN says that the values of  $\bar{X}_n$  are concentrated around  $\mathbb{E}(X_1)$ . But it is not giving us an idea about how the values are distributed around that mean value. CLT helps us in this regard.

To give the general idea of the CLT, let us use the following transform: for a r.v.  $X$ , let us denote

$$Z = \text{Standardize}(X) = \frac{X - \mathbb{E}(X)}{\sqrt{\text{Var}(X)}} = \frac{X - \mathbb{E}(X)}{SD(X)},$$

the Standardization (normalization, scaling) of  $X$ . Clearly,

$$\mathbb{E}(Z) = 0 \quad \text{and} \quad \text{Var}(Z) = 1.$$



## Basic Idea of the CLT

The basic idea of the CLT is the following: if we have a sequence of IID r.v.  $X_n$ , and we consider their sum  $S_n$  or their average  $\bar{X}_n$ , then

$$\text{Standardize}(S_n) \xrightarrow{D} \mathcal{N}(0, 1)$$

and

$$\text{Standardize}(\bar{X}_n) \xrightarrow{D} \mathcal{N}(0, 1).$$

## Basic Idea of the CLT

The basic idea of the CLT is the following: if we have a sequence of IID r.v.  $X_n$ , and we consider their sum  $S_n$  or their average  $\bar{X}_n$ , then

$$\text{Standardize}(S_n) \xrightarrow{D} \mathcal{N}(0, 1)$$

and

$$\text{Standardize}(\bar{X}_n) \xrightarrow{D} \mathcal{N}(0, 1).$$

Btw, trivially,

$$\text{Standardize}(\bar{X}_n) = \text{Standardize}(S_n),$$

and these two versions of CLT are the same.

## Basic Idea of the CLT

The basic idea of the CLT is the following: if we have a sequence of IID r.v.  $X_n$ , and we consider their sum  $S_n$  or their average  $\bar{X}_n$ , then

$$\text{Standardize}(S_n) \xrightarrow{D} \mathcal{N}(0, 1)$$

and

$$\text{Standardize}(\bar{X}_n) \xrightarrow{D} \mathcal{N}(0, 1).$$

Btw, trivially,

$$\text{Standardize}(\bar{X}_n) = \text{Standardize}(S_n),$$

and these two versions of CLT are the same.

So for large  $n$ , the Distribution of the  $\text{Standardize}(S_n)$  or  $\text{Standardize}(\bar{X}_n)$  is approximately Standard Normal.

## Basic Idea of the CLT

The basic idea of the CLT is the following: if we have a sequence of IID r.v.  $X_n$ , and we consider their sum  $S_n$  or their average  $\bar{X}_n$ , then

$$\text{Standardize}(S_n) \xrightarrow{D} \mathcal{N}(0, 1)$$

and

$$\text{Standardize}(\bar{X}_n) \xrightarrow{D} \mathcal{N}(0, 1).$$

Btw, trivially,

$$\text{Standardize}(\bar{X}_n) = \text{Standardize}(S_n),$$

and these two versions of CLT are the same.

So for large  $n$ , the Distribution of the  $\text{Standardize}(S_n)$  or  $\text{Standardize}(\bar{X}_n)$  is approximately Standard Normal. And this **independent of the initial Distribution of  $X_k$ !**

## Basic Idea of the CLT

The basic idea of the CLT is the following: if we have a sequence of IID r.v.  $X_n$ , and we consider their sum  $S_n$  or their average  $\bar{X}_n$ , then

$$\text{Standardize}(S_n) \xrightarrow{D} \mathcal{N}(0, 1)$$

and

$$\text{Standardize}(\bar{X}_n) \xrightarrow{D} \mathcal{N}(0, 1).$$

Btw, trivially,

$$\text{Standardize}(\bar{X}_n) = \text{Standardize}(S_n),$$

and these two versions of CLT are the same.

So for large  $n$ , the Distribution of the  $\text{Standardize}(S_n)$  or  $\text{Standardize}(\bar{X}_n)$  is approximately Standard Normal. And this **independent of the initial Distribution of  $X_k$ !**

Easy and beautiful, isn't it?

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) =$$



## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

Now we use

$$\mathbb{E}(S_n) =$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

Now we use

$$\mathbb{E}(S_n) = n \cdot \mu, \quad \text{Var}(S_n) =$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

Now we use

$$\mathbb{E}(S_n) = n \cdot \mu, \quad \text{Var}(S_n) = n \cdot \sigma^2.$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

Now we use

$$\mathbb{E}(S_n) = n \cdot \mu, \quad \text{Var}(S_n) = n \cdot \sigma^2.$$

Then,

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma}.$$

## CLT, in the Sums form

Assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

We consider sums

$$S_n = X_1 + X_2 + \dots + X_n.$$

We Standardize  $S_n$ :

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}}.$$

Now we use

$$\mathbb{E}(S_n) = n \cdot \mu, \quad \text{Var}(S_n) = n \cdot \sigma^2.$$

Then,

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma}.$$

The CLT states:

$$\frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma} \xrightarrow{D} \mathcal{N}(0, 1).$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ .

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$



## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) =$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

We use

$$\mathbb{E}(\bar{X}_n) =$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

We use

$$\mathbb{E}(\bar{X}_n) = \mu, \quad \text{Var}(\bar{X}_n) =$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

We use

$$\mathbb{E}(\bar{X}_n) = \mu, \quad \text{Var}(\bar{X}_n) = \frac{\sigma^2}{n}.$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

We use

$$\mathbb{E}(\bar{X}_n) = \mu, \quad \text{Var}(\bar{X}_n) = \frac{\sigma^2}{n}.$$

Then,

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}.$$

## CLT, in the Averages form

Again, assume  $X_n$  be a sequence of IID r.v. with finite expectation  $\mu = \mathbb{E}(X_i)$  and variance  $\sigma^2 = \text{Var}(X_i)$ . We consider the Averages

$$\bar{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}.$$

We Standardize  $\bar{X}_n$ :

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}}.$$

We use

$$\mathbb{E}(\bar{X}_n) = \mu, \quad \text{Var}(\bar{X}_n) = \frac{\sigma^2}{n}.$$

Then,

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}.$$

The CLT states:

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \xrightarrow{D} \mathcal{N}(0, 1).$$

## Two forms of CLT

Of course, these two forms of the CLT are the same: we have

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma}$$

and

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\sigma / \sqrt{n}}.$$



## Two forms of CLT

Of course, these two forms of the CLT are the same: we have

$$\text{Standardize}(S_n) = \frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} = \frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma}$$

and

$$\text{Standardize}(\bar{X}_n) = \frac{\bar{X}_n - \mathbb{E}(\bar{X}_n)}{\sqrt{\text{Var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}.$$

Now,

$$\frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma} = \frac{n \cdot (\frac{S_n}{n} - \mu)}{\sqrt{n} \cdot \sigma} = \frac{\frac{S_n}{n} - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}},$$


so

$$\text{Standardize}(S_n) = \text{Standardize}(\bar{X}_n).$$

Hence, the above two versions of CLT are the same, just one is in terms of  $S_n$ , the other one is in terms of  $\bar{X}_n$ .


# CLT Visually

```
n <- 600 # Sample Size
m <- 1000 # no of Samples
rate <- 0.2
x <- rexp(n*m, rate = rate)
theo.mean <- 1/rate #theoretical mean
theo.sd <- 1/rate #theoretical SD
m <- matrix(x, ncol = m); d <- data.frame(m)
avgs <- sapply(d, mean)
a = theo.mean-3*theo.sd/sqrt(n); b = theo.mean+3*theo.sd/sqrt(n)
hist(avgs, freq = F, xlim = c(a, b), ylim=c(0,3))
par(new = T)
t <- seq(a,b, 0.01)
y <- dnorm(t, mean = theo.mean, sd = theo.sd/sqrt(n))
plot(t,y, type = "l", col="red", lwd = 2, xlim = c(a,b), ylim=c(0,3))
```



## CLT, Visually, v2

```
n <- 600 # Sample Size  
m <- 1000 # no of Samples  
rate <- 0.2  
x <- rexp(n*m, rate = rate)  
m <- matrix(x, ncol = m); d <- data.frame(m)  
avgs <- sapply(d, mean)  
qqnorm(avgs, ylab = "Averages"); qqline(avgs)
```



## CLT, Roughly

In a non-rigorous way, we can write, for large  $n$  (here  $\approx$  means approximately distributed as):

$$\frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma} \approx \mathcal{N}(0, 1) \quad \text{and} \quad \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \approx \mathcal{N}(0, 1).$$

## CLT, Roughly

In a non-rigorous way, we can write, for large  $n$  (here  $\approx$  means approximately distributed as):

$$\frac{S_n - n \cdot \mu}{\sqrt{n} \cdot \sigma} \approx \mathcal{N}(0, 1) \quad \text{and} \quad \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \approx \mathcal{N}(0, 1).$$

or

$$S_n \approx \mathcal{N}(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X}_n \approx \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right).$$

## CLT, Roughly -2

Let us summarize:

## CLT, Roughly -2

Let us summarize:

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_k) = \sigma^2$ , and **are Normally Distributed**, i.e.,  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ , then

## CLT, Roughly -2

Let us summarize:

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_k) = \sigma^2$ , and **are Normally Distributed**, i.e.,  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$S_n \sim \mathcal{N}(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X}_n \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right);$$

so we know the **exact Distributions** of  $S_n$  and  $\bar{X}_n$ .



## CLT, Roughly -2

Let us summarize:

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_K) = \sigma^2$ , and **are Normally Distributed**, i.e.,  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$S_n \sim \mathcal{N}(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X}_n \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right);$$

so we know the **exact Distributions** of  $S_n$  and  $\bar{X}_n$ .

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_K) = \sigma^2$ , and **from any Distribution** (but the same Distribution), then

## CLT, Roughly -2

Let us summarize:

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_K) = \sigma^2$ , and **are Normally Distributed**, i.e.,  $X_k \sim \mathcal{N}(\mu, \sigma^2)$ , then

$$S_n \sim \mathcal{N}(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X}_n \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right);$$

so we know the **exact Distributions** of  $S_n$  and  $\bar{X}_n$ .

- ▶ If  $X_k$ -s are independent, have the Mean  $\mathbb{E}(X_k) = \mu$  and  $\text{Var}(X_K) = \sigma^2$ , and **from any Distribution** (but the same Distribution), then

$$S_n \approx \mathcal{N}(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X}_n \approx \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right);$$

and we know the **asymptotic Distributions** (approximate Distributions for large  $n$ ) of  $S_n$  and  $\bar{X}_n$ .

# CLT, Berry-Eseen Inequality

Now, quickly about the convergence rate of CLT:

**Theorem(18+, Berry-Eseen):** Assume  $X_k$  are IID r.v.s with finite  $\mathbb{E}(X_1) = \mu$ ,  $\text{Var}(X_1) = \sigma^2$  and  $\mathbb{E}(|X_1|^3)$ . Then, for any  $n \in \mathbb{N}$ ,

$$\sup_{x \in \mathbb{R}} |\mathbb{P}(Z_n \leq x) - \Phi(x)| \leq \frac{\mathbb{E}(|X_1 - \mu|^3)}{\sigma^3 \cdot \sqrt{n}},$$

where

$$Z_n = \text{Standardize}(S_n) = \text{Standardize}(\bar{X}_n),$$

and  $\Phi(x)$  is the CDF of  $\mathcal{N}(0, 1)$ .