# Deep Learning

Vazgen Mikayelyan

YSU, Krisp

November 25, 2020

## Outline

Metrics for Classification Problems


## Accuracy

#### Definition 1


Accuracy in classification problems is the percent of correct predictions made by the model over all kinds predictions made:

$$Accuracy = \frac{\textit{Number of correct predictions}}{\textit{Total numbers of predictions made}}$$

## Precision and Recall



## Precision and Recall



How many selected items are relevant?

How many relevant items are selected?

$$Precision = \frac{TP}{TP + FP}, \quad Recall = \frac{TP}{TP + FN}$$

$$Precision = rac{TP}{TP + FP}, \quad Recall = rac{TP}{TP + FN}$$
 
$$F1 = rac{2}{rac{1}{Precision} + rac{1}{Recall}}$$

$$Precision = rac{TP}{TP + FP}, \quad Recall = rac{TP}{TP + FN}$$
  $F1 = rac{2}{rac{1}{Precision} + rac{1}{Recall}}$ 

Why don't to use arithmetic mean instead of harmonic mean?

$$Precision = rac{TP}{TP + FP}, \quad Recall = rac{TP}{TP + FN}$$
  $F1 = rac{2}{rac{1}{Precision} + rac{1}{Recall}}$ 

Why don't to use arithmetic mean instead of harmonic mean? **Answer**: If Precision = 0.1 and Recall = 0.95, then their mean is equal to 0.525 and  $F1 \approx 0.18$ .

• A receiver operating characteristic curve, i.e. ROC curve is a graphical plot that illustrates the diagnostic ability of a binary classifier system as its discrimination threshold is varied.

- A receiver operating characteristic curve, i.e. ROC curve is a graphical plot that illustrates the diagnostic ability of a binary classifier system as its discrimination threshold is varied.
- The diagnostic performance of a test or the accuracy of a test to discriminate diseased cases from normal cases is evaluated using ROC curve analysis.

- A receiver operating characteristic curve, i.e. ROC curve is a graphical plot that illustrates the diagnostic ability of a binary classifier system as its discrimination threshold is varied.
- The diagnostic performance of a test or the accuracy of a test to discriminate diseased cases from normal cases is evaluated using ROC curve analysis.
- A ROC curve is a way to compare diagnostic tests. It is a plot of true positive rate against the false positive rate:

$$TPR = Recall = Sensitivity = \frac{TP}{TP + FN}$$

$$TPR = Recall = Sensitivity = \frac{TP}{TP + FN}$$

$$\mathit{FPR} = 1 - \mathit{Specificity} = 1 - \frac{\mathit{TN}}{\mathit{TN} + \mathit{FP}} = \frac{\mathit{FP}}{\mathit{TN} + \mathit{FP}}$$

$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

• What if *threshold* = 0?

$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

- What if *threshold* = 0?
  - TN = 0, FN = 0

$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

- What if *threshold* = 0?
  - TN = 0, FN = 0
  - TPR = 1, FPR = 1

$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

- What if *threshold* = 0?
  - TN = 0, FN = 0
  - TPR = 1, FPR = 1
- What if *threshold* = 1?


$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

- What if *threshold* = 0?
  - TN = 0, FN = 0
  - TPR = 1, FPR = 1
- What if *threshold* = 1?
  - TP = 0, FP = 0



$$TPR = Recall = Sensitivity = rac{TP}{TP + FN}$$
 
$$FPR = 1 - Specificity = 1 - rac{TN}{TN + FP} = rac{FP}{TN + FP}$$

- What if *threshold* = 0?
  - TN = 0, FN = 0
  - TPR = 1, FPR = 1
- What if *threshold* = 1?
  - TP = 0, FP = 0
  - TPR = 0, FPR = 0



## Definition 2

#### Definition 2

AUC (area under the curve) is the area under the ROC curve.

• What if  $AUC \approx 0.5$ ?

#### Definition 2

- What if  $AUC \approx 0.5$ ?
- What if  $AUC \approx 1$ ?

#### Definition 2

- What if  $AUC \approx 0.5$ ?
- What if  $AUC \approx 1$ ?
- How to choose threshold?

#### Definition 2

- What if  $AUC \approx 0.5$ ?
- What if  $AUC \approx 1$ ?
- How to choose threshold?
- How to use ROC curve for multi-class model?

# Balanced Cross Entropy

Simple Cross Entropy

$$L(w) = \frac{1}{n} \sum_{i=1}^{n} \left( -y_i^T \log f_w(x_i) \right).$$

# Balanced Cross Entropy

Simple Cross Entropy

$$L(w) = \frac{1}{n} \sum_{i=1}^{n} \left( -y_i^T \log f_w(x_i) \right).$$

Balanced Cross Entropy

$$L(w) = \frac{1}{n} \sum_{i=1}^{n} \left( -\left(\alpha * y_{i}\right)^{T} \log f_{w}\left(x_{i}\right) \right).$$

Where 
$$\alpha^T = [\alpha_1, \dots, \alpha_k]$$
,  $\alpha_i \in (0, 1)$  and  $\sum_{i=1}^k \alpha_i = 1$ .