

Pabna University Of Science And Technology



Department of Information And Communication Engineering

Assignment

Course Name: Engineering Statistics
Course Code: STAT-2201

Submitted By:

M.Q.A Haniya Khanam

ID:220632

Session:2021-22

2nd Year 2nd Semester

Department of ICE, PUST

Submitted To:

Dr. Md. Sarwar Hosain

Associate Professor

Department of ICE, PUST

Date of Submission:23-04-2025

Name of the Topic: Comparison of two sample Means.

The **comparison of two sample means** is a statistical technique used to determine whether there is a significant difference between the average values (means) of two independent groups or samples. This is a core concept in inferential statistics, commonly used in experiments, surveys, and observational studies.

Sample Mean:

The **sample mean** is the average value of a sample — a subset of a population. It's a point estimate of the population mean and is calculated as:

$$\bar{x} = \frac{\sum x_i}{n}$$

Where:

- \bar{x} = sample mean
- x_i = each value in the sample
- n = number of observations in the sample

Types of Two-Sample Mean Comparisons

There are two major cases:

1. Independent Samples

When the two samples are unrelated (e.g., group A and group B are made up of different individuals).

Test Used: Two-Sample (Independent) t-Test

This tests the null hypothesis:

$$H_0 : \mu_1 = \mu_2$$

Versus an alternative hypothesis (one-sided or two-sided):

$$H_1 : u_1 \neq u_2 \quad (\text{two-tailed})$$

$$H_1 : u_1 > u_2 \quad \text{or} \quad u_1 < u_2 \quad (\text{one-tailed})$$

Assumptions:

- Data are normally distributed (especially for small samples)
- Samples are independent
- Variances can be equal (pooled t-test) or unequal (Welch's t-test)
- Test Statistic:

If variances are equal:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

Where s_p is the pooled standard deviation:

$$s_p = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}$$

If variances are unequal (Welch's t-test):

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

Degrees of freedom (approximate) are calculated using the **Welch–Satterthwaite equation**.

2. Paired Samples

When the same individuals are measured twice (e.g., before and after treatment) or matched pairs.

Test Used: Paired t-Test

It compares the **mean of the differences** between paired observations.

$$H_0: \mu_d = 0 \quad (\text{no difference})$$

$$t = \frac{\bar{d}}{s_d / \sqrt{n}}$$

Where:

- \bar{d} = mean of the differences
- S_d = standard deviation of the differences
- n = number of pairs

Interpreting Results

- If the **p-value** < significance level (e.g., 0.05), reject the null hypothesis
- This suggests that the difference in means is **statistically significant**.
- A **confidence interval** (typically 95%) can also show the range of possible differences in means.

Example (Independent Samples)

A researcher wants to know if there's a difference in test scores between students in two different schools.

Sample A ($n = 30$): Mean = 75, SD = 10

Sample B ($n = 30$): Mean = 80, SD = 12

Using a two-sample t-test, the researcher can determine if the difference in means ($80 - 75 = 5$) is statistically significant or due to random variation.