



# IMA101 – A Short Introduction To Medical Imaging Modalities


Loïc Le Folgoc

LTCI, Télécom Paris, IP Paris


[loic.lefolgoc@telecom-paris.fr](mailto:loic.lefolgoc@telecom-paris.fr)

# Conventional radiography or X-ray


Source: Wikipedia




One of the 1<sup>st</sup> radiographs  
by Wilhelm Röntgen



Chest radiograph





- Uses X-rays, a high-energy electromagnetic radiation
- Ionizing radiation, harmful to living tissue
- Wavelength: gamma-ray  $\leq$  X-ray  $\leq$  UVs




- 2D image (single projection)
- Primary beam attenuated when passing through the body
- Depends on density and structural composition of anatomy (e.g., bone more attenuated than air and soft tissue)
- Applications:
  - Bone fracture detection
  - Chest X-rays: pneumonia, heart failure, etc.
  - Mammograms: breast cancer
  - ...

# Computed Tomography or CT scan



- Same principle as for X-rays but...
- ... X-ray source is motorized & rotates around the body
- After a full rotation, signal processed to digitally form a 2D slice (cf. inverse Radon transform)
- 3D image constructed progressively as a stack of 2D slices, as the motorized bed is moved forward in the gantry
- Intensity signal measured in Hounsfield Units (HU):  
water = 0 HU, air = -1000 HU, dense bone ~2000 HU
- View bone fractures, eroded joints
- View tumors or lesions (abdomen, lungs, head, bones...)
- Head scan: locate clots leading to stroke, hemorrhage
- Lungs: pulmonary embolism, excess fluid, emphysema...
- Contrast agents can be added to enhance the visibility of soft tissues or blood vessels
- e.g., CT angiography to examine the circulatory system, to look for possible obstructions

# Computed Tomography or CT scan





- Same principle as for X-rays but...
- ... X-ray source is motorized & rotates around the body
- After a full rotation, signal processed to digitally form a 2D slice (cf. inverse Radon transform)
- 3D image constructed progressively as a stack of 2D slices, as the motorized bed is moved forward in the gantry
- Intensity signal measured in Hounsfield Units (HU):  
water = 0 HU, air = -1000 HU, bone = 500 HU +
- View bone fractures, eroded joints
- View tumors or lesions (abdomen, lungs, head, bones...)
- Head scan: locate clots leading to stroke, hemorrhage
- Lungs: pulmonary embolism, excess fluid, emphysema...
- Contrast agents can be added to enhance the visibility of soft tissues or blood vessels
- e.g., CT angiography to examine the circulatory system, to look for possible obstructions




# Ultrasound imaging or (ultra)sonography (US)


Ultrasound transducer  
(probe)



- Uses high-frequency sound waves
- Waves bounce off body tissue and make echoes
- The echoes are received by the transducer and transformed in a digital image
- Safe for health
- 2D or 3D




3D fetal imaging  
(3 orthogonal views & volume reconstruction)



3D+t cardiac imaging  
(overlaid left ventricle contour)

- Effective for imaging soft tissues
- Many applications for screening & diagnosis
  - Cardiology (echocardiography)
  - Prenatal ultrasound
  - Point-of-care ultrasound
  - Abdominal US (liver tumors...)
  - Hemodynamics (blood flow)
  - Urology (bladder...)

# Magnetic Resonance Imaging (MRI)



$T_1$ -weighted  
(TR=600, TE=11)

Density-weighted  
(TR=3000, TE=17)

$T_2$ -weighted  
(TR=3800, TE=102)




- Perfect to image soft tissues
- Safe except contraindications (metal implants, etc.)
- Based on Nuclear Magnetic Resonance (NMR)
- Magnetization of hydrogen nuclei in an imaged sample
- 1.5T, 3T, up to 14T MRI depending on the strength of the large uniform magnetic field  $B_0$  applied to align the magnetic dipole moments of the  $H$  nuclei
- Many parameters resulting in many sequences with different appearances and contrast
  - Change repetition time  $TR$  and echo time  $TE$  (ms) to control the sensitivity of the signal to local tissue relaxation times  $T_1$  and  $T_2$  (depending on the proton density)
  - Flip angle
  - Acquisition plane (3D as stack of 2D)
  - ...

# Magnetic Resonance Imaging (MRI)

## MRI with contrast agent



Magnetic Resonance Angiography of the brain to image blood vessels (also common for neck, thoracic and abdominal aorta, renal arteries, legs)




Gadolinium-enhanced T1-w MRI to image the glioblastoma (tumour)


# Magnetic Resonance Imaging (MRI)

MRI can reveal kinematics

Source: M. Markl. Velocity encoding..., 2006.



**Phase-contrast MRI** reveals the 3D blood flow in arteries  
(e.g., shown here, heart and aorta in a sagittal slice)





**Cine MRI** showing the beating heart  
(left and right ventricles, 2 orthogonal planes shown)



# Magnetic Resonance Imaging (MRI)


MRI can reveal functional information → **functional MRI**




- Blood-oxygen-level dependent (BOLD) contrast
- fMRI measures brain activity via hemodynamic responses
  - Blood releases oxygen to active neurons at a greater rate than to inactive neurons
  - The difference between oxygenated and deoxygenated blood is measurable

- Used in research
- Detect signal changes in response to different stimuli

# Positron Emission Tomography (PET)



← Whole-body PET scan  
using a glucose-based  
radiotracer



↓ Brain PET scan

- Functional imaging technique
- Can be combined with CT and MRI to overlay an anatomical reference under the functional data (PET-CT, PET-MRI)
- Exposes to ionizing radiation
- A radioactive substance (radiotracer) is used to measure metabolic processes or physiological activity
  - The radiotracer undergoes positron emission decay in the tissue
  - Positrons are indirectly located as they annihilate with an electron
- Applications:
  - Oncology: detect tumours (glucose-based radiotracer)
  - Neuroimaging: measure the blood flow (oxygen-based radiotracer)
  - ...