


```
n = 20
slope = -0.000267    (se = 0.002413)
t(slope) = -0.111    p(slope) = 0.913014
intercept = 0.014101    (se = 0.028912)
t(intercept) = 0.488    p(intercept) = 0.631631
R² = 0.0007

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept not significant → no consistent baseline difference detected.
```




```
n = 20
slope = 0.002811    (se = 0.004028)
t(slope) = 0.698    p(slope) = 0.494179
intercept = -0.009835    (se = 0.048252)
t(intercept) = -0.204    p(intercept) = 0.840777
R² = 0.0263

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept not significant → no consistent baseline difference detected.
```




```
n = 20
slope = 0.000969    (se = 0.003635)
t(slope) = 0.266    p(slope) = 0.792890
intercept = -0.008670    (se = 0.043545)
t(intercept) = -0.199    p(intercept) = 0.844414
R² = 0.0039

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept not significant → no consistent baseline difference detected.
```




```
n = 20
slope = 0.000286    (se = 0.002502)
t(slope) = 0.114    p(slope) = 0.910337
intercept = 0.012200    (se = 0.029976)
t(intercept) = 0.407    p(intercept) = 0.688809
R² = 0.0007

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept not significant → no consistent baseline difference detected.
```




```
n = 20
slope = -0.828173    (se = 0.235761)
t(slope) = -3.513    p(slope) = 0.002485
intercept = 128.025842    (se = 2.421573)
t(intercept) = 52.869    p(intercept) = 0.000000
R² = 0.4067

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -2.405782    (se = 1.205152)
t(slope) = -1.996    p(slope) = 0.061262
intercept = 175.728515    (se = 12.378505)
t(intercept) = 14.196    p(intercept) = 0.000000
R² = 0.1813

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -1.081795    (se = 0.346918)
t(slope) = -3.118    p(slope) = 0.005937
intercept = 136.761121    (se = 3.563304)
t(intercept) = 38.380    p(intercept) = 0.000000
R² = 0.3507

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.023877    (se = 0.006945)
t(slope) = 3.438    p(slope) = 0.002934
intercept = -5.309021    (se = 0.071335)
t(intercept) = -74.424    p(intercept) = 0.000000
R² = 0.3964

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.030222    (se = 0.007942)
t(slope) = 3.805    p(slope) = 0.001296
intercept = -5.415879    (se = 0.081580)
t(intercept) = -66.388    p(intercept) = 0.000000
R² = 0.4458

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.000000    (se = 0.000000)
t(slope) = 0.000    p(slope) = 1.000000
intercept = 20.352461    (se = 0.000000)
t(intercept) = 13644655226018606.000    p(intercept) = 0.000000
R² = 0.0000

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.034467    (se = 0.003839)
t(slope) = -8.977    p(slope) = 0.000000
intercept = 0.778761    (se = 0.045991)
t(intercept) = 16.933    p(intercept) = 0.000000
R² = 0.8174

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.024543    (se = 0.005417)
t(slope) = -4.531    p(slope) = 0.000259
intercept = 0.588957    (se = 0.064893)
t(intercept) = 9.076    p(intercept) = 0.000000
R² = 0.5328

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.037202    (se = 0.004351)
t(slope) = -8.550    p(slope) = 0.000000
intercept = 0.746122    (se = 0.052121)
t(intercept) = 14.315    p(intercept) = 0.000000
R² = 0.8024

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.011629    (se = 0.002575)
t(slope) = -4.517    p(slope) = 0.000267
intercept = 0.465428    (se = 0.030841)
t(intercept) = 15.091    p(intercept) = 0.000000
R² = 0.5313

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.819451    (se = 0.198823)
t(slope) = -4.122    p(slope) = 0.000641
intercept = 15.203584    (se = 2.042177)
t(intercept) = 7.445    p(intercept) = 0.000001
R² = 0.4855

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -3.024099    (se = 1.333166)
t(slope) = -2.268    p(slope) = 0.035847
intercept = 47.010913    (se = 13.693371)
t(intercept) = 3.433    p(intercept) = 0.002966
R² = 0.2223

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -1.257305    (se = 0.294669)
t(slope) = -4.267    p(slope) = 0.000464
intercept = 21.778704    (se = 3.026642)
t(intercept) = 7.196    p(intercept) = 0.000001
R² = 0.5028

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.016602    (se = 0.004066)
t(slope) = 4.083    p(slope) = 0.000699
intercept = -0.306996    (se = 0.041768)
t(intercept) = -7.350    p(intercept) = 0.000001
R² = 0.4808

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.024119    (se = 0.005541)
t(slope) = 4.353    p(slope) = 0.000383
intercept = -0.414001    (se = 0.056911)
t(intercept) = -7.274    p(intercept) = 0.000001
R² = 0.5128

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.000000    (se = 0.000000)
t(slope) = N/A    p(slope) = N/A
intercept = 0.000000    (se = 0.000000)
t(intercept) = N/A    p(intercept) = N/A
R² = 0.0000

Slope p-value unavailable; cannot determine significance here.
Intercept p-value unavailable; cannot determine significance here.
```




```
n = 20
slope = -0.034199    (se = 0.003259)
t(slope) = -10.493    p(slope) = 0.000000
intercept = 0.764660    (se = 0.039042)
t(intercept) = 19.586    p(intercept) = 0.000000
R² = 0.8595

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.027354    (se = 0.004527)
t(slope) = -6.042    p(slope) = 0.000010
intercept = 0.598792    (se = 0.054231)
t(intercept) = 11.041    p(intercept) = 0.000000
R² = 0.6698

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```



n = 20
slope = -0.038170 (se = 0.003272)
t(slope) = -11.667 p(slope) = 0.000000
intercept = 0.754792 (se = 0.039191)
t(intercept) = 19.259 p(intercept) = 0.000000
$R^2 = 0.8832$

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.




```
n = 20
slope = -0.011915    (se = 0.001619)
t(slope) = -7.361    p(slope) = 0.000001
intercept = 0.453228    (se = 0.019389)
t(intercept) = 23.375    p(intercept) = 0.000000
R² = 0.7507

Slope ≠ 0 (statistically significant) → metric difference changes with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.008722    (se = 0.209481)
t(slope) = 0.042    p(slope) = 0.967247
intercept = -112.822257    (se = 2.151647)
t(intercept) = -52.435    p(intercept) = 0.000000
R² = 0.0001

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.618316    (se = 0.942217)
t(slope) = -0.656    p(slope) = 0.519974
intercept = -128.717602    (se = 9.677809)
t(intercept) = -13.300    p(intercept) = 0.000000
R² = 0.0234

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.175510    (se = 0.251145)
t(slope) = -0.699    p(slope) = 0.493585
intercept = -114.982417    (se = 2.579595)
t(intercept) = -44.574    p(intercept) = 0.000000
R² = 0.0264

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.007275    (se = 0.006492)
t(slope) = -1.121    p(slope) = 0.277176
intercept = 5.002025    (se = 0.066680)
t(intercept) = 75.016    p(intercept) = 0.000000
R² = 0.0652

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = -0.006103    (se = 0.007834)
t(slope) = -0.779    p(slope) = 0.446042
intercept = 5.001878    (se = 0.080461)
t(intercept) = 62.165    p(intercept) = 0.000000
R² = 0.0326

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```




```
n = 20
slope = 0.000000    (se = 0.000000)
t(slope) = 0.000    p(slope) = 1.000000
intercept = -20.352461    (se = 0.000000)
t(intercept) = -13644655226018606.000    p(intercept) = 0.000000
R² = 0.0000

Slope ≈ 0 (not significant) → metric difference not correlated with epoch.
Intercept is significant → persistent baseline difference between models.
```