

Problem 1.1. (30 points total)

- (a) (20 points) Clearly documenting your extremely careful work, compute the reduced row-echelon form of the matrix

$$A = \begin{bmatrix} 1 & 2 & 1 & 3 & 0 \\ 2 & 4 & 4 & 10 & -2 \\ 1 & 2 & 5 & 11 & -4 \end{bmatrix}$$

- (b) (1 points) Which columns of A have pivots? 1, 3 (list column numbers)

$$\begin{array}{c} \bullet \\ 2 \\ 1 \end{array} \begin{bmatrix} 1 & 2 & 1 & 3 & 0 \\ 2 & 4 & 4 & 10 & -2 \\ 1 & 2 & 5 & 11 & -4 \end{bmatrix} \sim \begin{array}{c} \bullet \\ 2 \end{array} \begin{bmatrix} 1 & 2 & 1 & 3 & 0 \\ 0 & 0 & 2 & 4 & -2 \\ 0 & 0 & 4 & 8 & -4 \end{bmatrix} \sim \times_{1/2} \begin{bmatrix} 1 & 2 & 1 & 3 & 0 \\ 0 & 0 & 2 & 4 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{c} 1 \\ \bullet \end{array} \begin{bmatrix} 1 & 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \sim \boxed{\begin{bmatrix} 1 & 2 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}}$$

- (c) (9 points) If possible, find the solution of the linear system below in parametric vector form.

$$\begin{cases} x_1 + 2x_2 + x_3 + 3x_4 = 0 \\ 2x_1 + 4x_2 + 4x_3 + 10x_4 = -2 \\ x_1 + 2x_2 + 5x_3 + 11x_4 = -4 \end{cases}$$

Since the augmented matrix for this system is the matrix given in part (a), the rref computed in (a) gives the information needed for the parametric vector form:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -1 \\ 0 \\ -2 \\ 1 \end{bmatrix}$$

Problem 1.2. (6 points) Carefully simplify each valid expression. For any invalid expression, explain why it is invalid.

$$(a) \quad \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \quad \boxed{\text{is undefined}}$$

Here there is a 5×4 matrix times a 5×1 matrix.

$$(b) \quad \begin{bmatrix} 1 & 3 & 1 \\ 4 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} - \begin{bmatrix} 3 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 5 \\ 9 & 8 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 4 & 0 \end{bmatrix} - \begin{bmatrix} 3 & 3 \\ 2 & 2 \end{bmatrix} = \boxed{\begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}}$$

Alternatively, one may recognize the given expression as a product minus the first two terms of its outer product expansion. Hence, the expression is equal to the third term in the outer product expansion $\begin{bmatrix} 1 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix}$.

$$(c) \quad \begin{bmatrix} 1 & 3 & 1 \\ 4 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 3 & 1 \\ 4 & 2 & 3 \end{bmatrix}^T - \begin{bmatrix} 1 & 3 & 1 \\ 4 & 2 & 3 \end{bmatrix}^T \begin{bmatrix} 1 & 3 & 1 \\ 4 & 2 & 3 \end{bmatrix} \quad \text{is } \boxed{\text{undefined}}.$$

The first product in the expression is a 2×3 matrix times a 3×2 matrix giving a 2×2 matrix.

The second product in the expression is a 3×2 matrix times a 2×3 matrix giving a 3×3 matrix.

Since these two products have different dimensions their difference is undefined.

$$(d) \quad [1 \quad 1 \quad 1 \quad 1 \quad 1] \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \boxed{\text{is undefined}}$$

The product given is a 1×5 matrix times a 5×4 matrix by a 5×1 matrix.

$$(e) \quad [0 \quad 0] \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix}^T = \boxed{[0 \quad 0 \quad 0 \quad 0 \quad 0]}$$

The given product of four matrices is a 1×2 matrix times a 2×2 matrix times a 2×1 matrix times a 1×5 matrix; hence, the product is defined and is a 1×5 matrix. Since one of the factors is a zero matrix, the product is also a zero matrix.

$$(f) \quad \begin{bmatrix} \sqrt{3} & -1 \\ 3 & -\sqrt{3} \end{bmatrix}^2 = \begin{bmatrix} \sqrt{3} & -1 \\ 3 & -\sqrt{3} \end{bmatrix} \begin{bmatrix} \sqrt{3} & -1 \\ 3 & -\sqrt{3} \end{bmatrix} = \begin{bmatrix} 3-3 & -\sqrt{3}+\sqrt{3} \\ 3\sqrt{3}-3\sqrt{3} & -3+3 \end{bmatrix} = \boxed{\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}}$$

Hence, we find an unexpected square root of the 2×2 zero matrix.

Problem 1.3. (6 points) For each matrix below, compute the matrix inverse, if it exists. If it fails to exist explain why.

$$A_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

$$\begin{matrix} 1 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 & -1 \\ 0 & 1 & 0 & 1 \end{bmatrix} \Rightarrow A_1^{-1} = \boxed{\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}}$$

$$A_2 = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{matrix} 1 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{matrix} 1 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{matrix} 1 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & -1 & 1 & -1 \\ 0 & 1 & 0 & 0 & 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow A_2^{-1} = \boxed{\begin{bmatrix} 1 & -1 & 1 & -1 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix}}$$

$$A_3 = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

$$\bullet_1 \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Hence the second column has no pivot and the inverse does not exist.

$$A_4 = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$\bullet_4 \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{bmatrix} \sim \bullet_2 \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{bmatrix}$$

Hence the third column has no pivot and the inverse does not exist.

$$A_5 = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 10 \end{bmatrix}$$

$$\begin{aligned} & \bullet \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 10 & 0 & 0 & 1 \end{bmatrix} \sim \bullet_2 \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -11 & -7 & 0 & 1 \end{bmatrix} \sim \times -1/3 \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{bmatrix} \sim \\ & \begin{matrix} 3 \\ 2 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & 2 & 4/3 & -1/3 & 0 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{bmatrix} \sim \begin{matrix} 2 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 2 & 0 & -2 & 6 & -3 \\ 0 & 1 & 0 & -2/3 & 11/3 & -2 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & -2/3 & -4/3 & 1 \\ 0 & 1 & 0 & -2/3 & 11/3 & -2 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{bmatrix} \end{aligned}$$

$$A_5^{-1} = \boxed{\begin{bmatrix} -2/3 & -4/3 & 1 \\ -2/3 & 11/3 & -2 \\ 1 & -2 & 1 \end{bmatrix}}$$

$$A_6 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 4 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{matrix} \bullet \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 4 & 0 & 0 & 1 & 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -2 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & -4 & 0 & 0 & 1 \end{bmatrix}$$

$$A_6^{-1} = \boxed{\begin{bmatrix} 1 & 0 & 0 & 0 \\ -2 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ -4 & 0 & 0 & 1 \end{bmatrix}}$$

Problem 1.4. (4 points) If possible, find the LU factorization of each matrix below:

$$A_1 = \begin{bmatrix} 0 & 1 & 2 & 3 & 0 \\ 1 & 2 & 3 & 4 & 0 \end{bmatrix}$$

not possible as row exchange is necessary in the forward phase.

$$A_2 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \end{bmatrix}$$

$$\begin{matrix} \bullet \\ 2 \\ 3 \end{matrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \text{pref}(A_2)$$

$$A_2 = \left[\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \right]$$

$$A_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

not possible as row exchange is necessary in the forward phase.

$$A_4 = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 4 & 6 & 8 & 10 & 12 & 14 & 16 \end{bmatrix}$$

$$\bullet \begin{matrix} 1 \\ 2 \end{matrix} \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 4 & 6 & 8 & 10 & 12 & 14 & 16 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} = \text{pref}(A_4)$$

$$A_4 = \left[\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \right]$$

Problem 1.5. (4 points) If possible, find the CR factorization of each matrix below:

$$A_1 = \begin{bmatrix} 0 & 1 & 2 & 3 & 0 \\ 1 & 2 & 3 & 4 & 0 \end{bmatrix}$$

$$\begin{matrix} \downarrow \\ \uparrow \end{matrix} \begin{bmatrix} 0 & 1 & 2 & 3 & 0 \\ 1 & 2 & 3 & 4 & 0 \end{bmatrix} \sim \begin{matrix} 2 \\ \bullet \end{matrix} \begin{bmatrix} 1 & 2 & 3 & 4 & 0 \\ 0 & 1 & 2 & 3 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1 & -2 & 0 \\ 0 & 1 & 2 & 3 & 0 \end{bmatrix} = \text{rref}(A_1)$$

$$A_1 = \boxed{\begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 & -2 & 0 \\ 0 & 1 & 2 & 3 & 0 \end{bmatrix}}$$

$$A_2 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \end{bmatrix}$$

$$\begin{matrix} \bullet \\ 2 \\ 3 \end{matrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 2 & 2 \\ 3 & 3 & 3 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \text{rref}(A_2)$$

$$A_2 = \boxed{\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}}$$

$$A_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{array}{c} \downarrow \\ \uparrow \end{array} \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \sim \begin{array}{c} \downarrow \\ \uparrow \end{array} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \text{rref}(A_3)$$

$$A_3 = \left[\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right]$$

$$A_4 = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 4 & 6 & 8 & 10 & 12 & 14 & 16 \end{bmatrix}$$

$$\begin{array}{c} \bullet \\ 2 \end{array} \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 2 & 4 & 6 & 8 & 10 & 12 & 14 & 16 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} = \text{rref}(A_4)$$

$$A_4 = \left[\begin{bmatrix} 1 \\ 2 \end{bmatrix} [1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7 \ 8] \right]$$

Problem 2. (10 points) Evaluate the truth of each statement below. If the statement is true write T in the box preceeding the statment. Otherwise, write F .

- (a) F A homogeneous linear system must have infinitely many solutions.

For example, $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ has $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ as its unique solution.

- (b) F If A is 3×3 and B is 3×3 then it must be that $AB \neq BA$.

For example, $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

- (c) T A sum of two symmetric 3×3 matrices is symmetric.

- (d) T The product of two invertible 3×3 matrices is invertible.

$$(AB)^{-1} = B^{-1}A^{-1}$$

- (e) F Every matrix may be factored as LU where L is lower unitriangular and U is proto-row-echelon.

Matrices requiring row exchange have no LU factorization.

- (a) F A linear system with one or more free variables has infinitely many solutions.

The linear system with augmented matrix $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ has a free variable but no solutions.

- (b) F Two different matrices cannot have the same reduced row-echelon form.

$$\text{rref}\left(\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}\right) = \text{rref}\left(\begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}\right) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

- (c) F An inconsistent system must have more equations than unknowns.

The linear system with the augmented matrix $[0 \ 0 \ 1]$ has one equation with two unknowns but is inconsistent.

- (d) T The linear combination of two solutions of a homogeneous linear system is also a solution.

If $A\mathbf{x} = \mathbf{0}$ and $A\mathbf{y} = \mathbf{0}$, then $A(\alpha\mathbf{x} + \beta\mathbf{y}) = \alpha A\mathbf{x} + \beta A\mathbf{y} = \alpha\mathbf{0} + \beta\mathbf{0} = \mathbf{0}$.

- (e) F If A and B are 2×2 matrices and $AB = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, then either A or B must be equal to $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Problem 3. (10 points) [Sample Problem A] What is the result of the MATLAB commands below.

```
>> A = [1 2 3; 2 3 4];
>> rref(A)
```

ans =

$$\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

```
>> rref([2 2; 0 2]*A)
```

ans =

$$\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

The long way.

$$\begin{bmatrix} 2 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \end{bmatrix} = \begin{bmatrix} \bullet & & \\ & 2/3 & \end{bmatrix} \begin{bmatrix} 6 & 10 & 14 \\ 4 & 6 & 8 \end{bmatrix} \sim \begin{bmatrix} \times 1/6 & & \\ & \times -3/2 & \end{bmatrix} \begin{bmatrix} 6 & 10 & 14 \\ 0 & -2/3 & -4/3 \end{bmatrix} \sim \begin{bmatrix} 5/3 & & \\ & \bullet & \end{bmatrix} \begin{bmatrix} 1 & 5/3 & 7/3 \\ 0 & 1 & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$$

The short way. Premultiplying by an invertible matrix does not change the rref.

Problem 3. (10 points) [Sample Problem B] Suppose that the MATLAB function `initialzeros` is defined by

```
function zerocount=initialzeros(A)
    [m n]=size(A);
    zerocount=zeros(m,1);    % column vector of m zeros
    for r=1:m
        for c=1:n
            if A(r,c) ~= 0
                break;
            end
            zerocount(r)=zerocount(r)+1;
        end
    end
end
```

Assuming `initialzeros` is in the current MATLAB search path, what is the output obtained by entering the following MATLAB commands:

```
>> A = [ 0 0 2 3 4; 0 0 0 0 0; 1 2 3 4 5; 0 0 2 0 0; 0 1 2 3 2; 0 0 0 0 0];
>> initialzeros(A)
```

```
ans =
     2
     5
     0
     2
     1
     5
```

`Zerocount` counts the number of zeros at the beginning of each row.

Problem 4. (15 points) [Sample Problem A] The matrices listed below constitute all possible 3×3 reduced row-echelon matrices. Note that some of these matrices contain parameters which are arbitrary scalar values.

$$A_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad A_2 = \begin{bmatrix} 1 & a & b \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad A_3 = \begin{bmatrix} 0 & 1 & a \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad A_4 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$A_5 = \begin{bmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 0 \end{bmatrix} \quad A_6 = \begin{bmatrix} 1 & a & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad A_7 = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad A_8 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(a) Determine the rank of each matrix above.

$$\text{rank}(A_1) = 0, \quad \text{rank}(A_2) = \text{rank}(A_3) = \text{rank}(A_4) = 1,$$

$$\text{rank}(A_5) = \text{rank}(A_6) = \text{rank}(A_7) = 2, \quad \text{rank}(A_8) = 3.$$

(b) For each matrix above, give the inverse, if possible.

Every matrix, except A_8 , has a column without a pivot and hence has no inverse.

$$A_8^{-1} = A_8 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(c) Which of the matrices above, considered as augmented matrices, represent an inconsistent system of linear equations?

$\boxed{A_4, A_6, A_7, A_8}$ represent inconsistent systems because each has a pivot in the last column.

(d) For each matrix above that represents a linear system having multiple solutions, provide the solution set in parametric vector form.

$$A_1 : \quad \mathbf{x} = \boxed{\alpha \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 0 \\ 1 \end{bmatrix}} \quad \text{the whole plane}$$

$$A_2 : \quad \mathbf{x} = \boxed{\begin{bmatrix} b \\ 0 \end{bmatrix} + \beta \begin{bmatrix} -a \\ 1 \end{bmatrix}} \quad \text{nonhorizontal lines}$$

$$A_3 : \quad \mathbf{x} = \boxed{\begin{bmatrix} 0 \\ a \end{bmatrix} + \alpha \begin{bmatrix} 1 \\ 0 \end{bmatrix}} \quad \text{horizontal lines}$$

Problem 4. (15 points) [Sample Problem B] Suppose

$$A = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 3 & 2 & 1 & 1 \\ 1 & 3 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} 1 & 1 & 0 & -3 \\ -2 & -1 & 0 & 5 \\ 3 & 2 & 1 & -10 \\ -2 & -2 & -1 & 9 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}, \quad B^{-1} = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix}.$$

(a) Solve the linear system

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 3 & 2 & 1 & 1 \\ 1 & 3 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

$$\mathbf{x} = A^{-1} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & -3 \\ -2 & -1 & 0 & 5 \\ 3 & 2 & 1 & -10 \\ -2 & -2 & -1 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \boxed{\begin{bmatrix} -1 \\ 2 \\ -4 \\ 4 \end{bmatrix}}$$

(b) Solve the linear system

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 3 & 2 & 1 & 1 \\ 1 & 3 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

$$\begin{aligned} \mathbf{x} &= B^{-1}A^{-1} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & -3 \\ -2 & -1 & 0 & 5 \\ 3 & 2 & 1 & -10 \\ -2 & -2 & -1 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \\ &= \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \\ -4 \\ 4 \end{bmatrix} = \begin{bmatrix} -2-2 \\ 1+4+4 \\ -2-8-4 \\ 4+4 \end{bmatrix} = \boxed{\begin{bmatrix} -4 \\ 9 \\ -14 \\ 8 \end{bmatrix}} \end{aligned}$$

(c) Simplify

$$\begin{aligned} &\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}^{-1} \left(\begin{bmatrix} 1 & 1 & 2 & 2 \\ 3 & 2 & 1 & 1 \\ 1 & 3 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}^{-1} \right)^{-1} \begin{bmatrix} 1 & 1 & 2 & 2 \\ 3 & 2 & 1 & 1 \\ 1 & 3 & 3 & 2 \\ 1 & 1 & 1 & 1 \end{bmatrix}^{-1} \\ &= B^{-1}(A^{-1}B^{-1})^{-1}A^{-1} = B^{-1}(BA)A^{-1} = (B^{-1}B)(AA^{-1}) = \boxed{\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}} \end{aligned}$$

Problem 4. (15 points) Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \\ 7 & 8 \end{bmatrix}$. Fully explain your responses.

(a) Determine the rank (number of pivots) of the matrix A .

$$\begin{array}{c} \bullet \\ 3 \\ 5 \\ 7 \end{array} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \\ 7 & 8 \end{bmatrix} \sim \begin{array}{c} \bullet \\ 2 \\ 3 \end{array} \begin{bmatrix} 1 & 2 \\ 0 & -2 \\ 0 & -4 \\ 0 & -6 \end{bmatrix} \sim \times -1/2 \begin{bmatrix} 1 & 2 \\ 0 & -2 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \sim \begin{array}{c} 2 \\ \bullet \end{array} \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

There are two pivots; hence, $\boxed{\text{rank}(A) = 2}$.

(b) Determine the rank of the matrix $A^T A$.

$$A^T A = \begin{bmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 1+9+25+49 & 2+12+30+56 \\ 2+12+30+56 & 4+16+36+64 \end{bmatrix} = \begin{bmatrix} 84 & 100 \\ 100 & 120 \end{bmatrix}$$

$$84 \cdot 120 - 100 \cdot 100 \neq 0$$

So $A^T A$ is invertible and must therefore have two pivots. $\boxed{\text{rank}(A^T A) = 2}$.

(c) Determine the rank of the matrix AA^T .

The long way.

$$AA^T = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{bmatrix} = \begin{array}{c} \bullet \\ 11/5 \\ 17/5 \\ 23/5 \end{array} \begin{bmatrix} 5 & 11 & 17 & 23 \\ 11 & 25 & 39 & 53 \\ 17 & 39 & 61 & 83 \\ 23 & 53 & 83 & 113 \end{bmatrix} \sim \begin{array}{c} \bullet \\ 2 \\ 3 \end{array} \begin{bmatrix} 5 & 11 & 17 & 23 \\ 0 & 4/5 & 8/5 & 12/5 \\ 0 & 8/5 & 16/5 & 24/5 \\ 0 & 12/5 & 24/5 & 36/5 \end{bmatrix} \sim$$

$$\begin{bmatrix} 5 & 11 & 17 & 23 \\ 0 & 4/5 & 8/5 & 12/5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \Rightarrow \boxed{\text{rank}(AA^T) = 2}$$

The short way. Premultiplying by the same elementary matrices that would bring A to $\text{rref}(A)$ shows

$$AA^T \sim \text{rref}(A)A^T = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{bmatrix} = \begin{array}{c} \bullet \\ 2 \end{array} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & 5 & 7 \\ 0 & -2 & -4 & -6 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

So $A^T A$ also has two pivots; hence, $\boxed{\text{rank}(A^T A) = 2}$.

Some morals to the story:

- (1) A row operation on a product may be performed on the first factor before multiplying. This follows from the fact that a row operation may be represented by premultiplication by an elementary matrix E . $E(AB) = (EA)B$
- (2) A , A^T , $A^T A$, and AA^T all share the same rank.

Problem 5. (15 points) [Sample Problem A] An $n \times n$ matrix A is congruent to an $n \times n$ matrix B if and only if there is a nonsingular $n \times n$ matrix S such that

$$SAS^T = B.$$

- (a) Explain why every $n \times n$ matrix is congruent to itself.

Taking $S = I$ shows $SAS^T = IAI = A$.

- (b) Explain why if A is congruent to B then B is congruent to A .

Since A is congruent to B there is a nonsingular matrix S such that $SAS^T = B$. To show that B is congruent to A we must find a nonsingular matrix \tilde{S} such that $\tilde{S}B\tilde{S}^T = A$. We multiply the known equality $SAS^T = B$ on the left by S^{-1} and on the right by S^{-T} :

$$\begin{aligned} S^{-1}SAS^TS^{-T} &= S^{-1}BS^{-T} \\ \Rightarrow IAI &= S^{-1}BS^{-T} \\ \Rightarrow \tilde{S}B\tilde{S}^T &= A \quad \text{where } \tilde{S} = S^{-1} \end{aligned}$$

- (c) Explain why if A is congruent to B and B is congruent to C then A is congruent to C .

Since A is congruent to B there is $S \in \text{GL}(n)$ such that $SAS^T = B$ and since B is congruent to C there is $\tilde{S} \in \text{GL}(n)$ such that $\tilde{S}B\tilde{S}^T = C$. To show that A is congruent to C we must find $\check{S} \in \text{GL}(n)$ such that $\check{S}A\check{S}^T = C$. Multiplying the equality $SAS^T = B$ on the left by \tilde{S} and on the right by \tilde{S}^T gives

$$\tilde{S}SAS^T\tilde{S}^T = \tilde{S}B\tilde{S}^T = C.$$

Hence, $\check{S}A\check{S}^T = C$ where $\check{S} = \tilde{S}S$ (since $\check{S}^T = (\tilde{S}S)^T = S^T\tilde{S}^T$).

Problem 5. (15 points) [Sample Problem B]

- (a) Suppose that
- $\mathbf{a} \in \mathbb{R}^m$
- and
- $\mathbf{b} \in \mathbb{R}^n$
- what is the rank of
- \mathbf{ab}^T
- ? Explain why.

The rank is either 0 or 1. If \mathbf{a} or \mathbf{b} is a zero vector then \mathbf{ab}^T is a zero matrix and has a rank of zero. If neither $\mathbf{a} = \mathbf{0}$ nor $\mathbf{b} = \mathbf{0}$ then every row of \mathbf{ab}^T is a multiple of \mathbf{b}^T . Hence, \mathbf{ab}^T has only one pivot.

- (b) Consider the matrix $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 8 & 12 \\ -3 & -6 & -9 \end{bmatrix}$. Find $\mathbf{a} \in \mathbb{R}^3$ and $\mathbf{b} \in \mathbb{R}^3$ such that $A = \mathbf{ab}^T$.

$$\boxed{\mathbf{a} = \begin{bmatrix} 1 \\ 4 \\ -3 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}} \quad \text{alternatively} \quad \mathbf{a} = \begin{bmatrix} 10 \\ 40 \\ -30 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 1/10 \\ 2/10 \\ 3/10 \end{bmatrix}$$

- (c) Consider the matrix $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$. Find vectors $\mathbf{a}_1, \mathbf{a}_2, \mathbf{b}_1, \mathbf{b}_2$ such that

$$A = \mathbf{a}_1 \mathbf{b}_1^T + \mathbf{a}_2 \mathbf{b}_2^T.$$

An idea: Select \mathbf{a}_1 and \mathbf{b}_1 to make $A - \mathbf{a}_1 \mathbf{b}_1^T$ have rank 1. Eliminating the first row and column with $\mathbf{a}_1 = [1 \ 4 \ 7]^T$ and $\mathbf{b}_1 = [1 \ 2 \ 3]^T$ would seem to have some promise. Indeed,

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 4 & 8 & 12 \\ 7 & 14 & 21 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} \begin{bmatrix} 0 & -3 & -6 \end{bmatrix}$$

Hence,

$$\boxed{\mathbf{a}_1 = \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix}, \quad \mathbf{b}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \mathbf{a}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 0 \\ -3 \\ -6 \end{bmatrix}}$$

Alternatively, because the matrix has rank 2, the outer product expansion of the CR factorization contains two terms:

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \end{bmatrix} + \begin{bmatrix} 2 \\ 5 \\ 8 \end{bmatrix} \begin{bmatrix} 0 & 1 & 2 \end{bmatrix}$$

Problem 5. (15 points) [Sample Problem C] Carefully explain your answers to each part of this question.

(a) How many 2×2 real symmetric matrices are square roots of $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$?

$$\begin{bmatrix} a & b \\ b & c \end{bmatrix}^2 = \begin{bmatrix} a^2 + b^2 & b(a+c) \\ b(a+c) & b^2 + c^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \Rightarrow a^2 + b^2 = 1, b(a+c) = 0, \text{ and } b^2 + c^2 = 1.$$

Since $b(a+c) = 0$ either $b = 0$ or $a+c = 0$. If $b = 0$ then $a^2 = 1$ and $c^2 = 1$. Hence, there are four square roots when $b = 0$:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

If $a+c = 0$ then $a = \pm\sqrt{1-b^2}$ so the square roots are

$$\begin{bmatrix} \pm\sqrt{1-b^2} & b \\ b & \mp\sqrt{1-b^2} \end{bmatrix} \quad -1 \leq b \leq 1.$$

There are infinitely many symmetric square roots of the 2×2 identity matrix.

(b) How many 2×2 real antisymmetric matrices are square roots of $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$?

$$\begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix}^2 = \begin{bmatrix} -a^2 & 0 \\ 0 & -a^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Hence, there are no real antisymmetric square roots of I .

(c) How many 2×2 real antisymmetric matrices are fourth roots of $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$?

$$\begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix}^4 = \begin{bmatrix} a^4 & 0 \\ 0 & a^4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Hence, $a = \pm 1$ (assuming $a \in \mathbb{R}$). The two real, antisymmetric fourth roots of I are

$$\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Problem 5. (15 points) [Sample Problem D] Consider Gauss-Jordan applied to $A =$

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}:$$

$$\begin{array}{ccccccc} & A & & \text{pref}(A) & & \text{nref}(A) & & \text{rref}(A) \\ \bullet & \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} & \sim & \bullet & \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{bmatrix} & \sim & \times -1/3 & \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 0 \end{bmatrix} & \sim & \bullet & \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} & \sim & \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \end{array}$$

- (a) Find a lower unitriangular matrix L , a diagonal matrix D , and an upper unitriangular matrix U such that

$$A = L \text{pref}(A), \quad \text{pref}(A) = D \text{nref}(A), \quad \text{and} \quad \text{nref}(A) = U \text{rref}(A).$$

$$L = \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -7 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right)^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 7 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{bmatrix}$$

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1/3 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1}$$

$$U = \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1}$$

$$L = \boxed{\begin{bmatrix} 1 & 0 & 0 \\ 4 & 1 & 0 \\ 7 & 2 & 1 \end{bmatrix}} \quad D = \boxed{\begin{bmatrix} 1 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 1 \end{bmatrix}} \quad U = \boxed{\begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}}$$

Because that matrix is not of full rank (there is a row of zeros in the row-echelon forms), there are multiple correct answers for D and U :

$$D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & a \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} 1 & 2 & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{bmatrix}$$

Note that D is required to be diagonal and U , upper unitriangular.

- (b) Find a matrix C such that $A = C \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$.

From the CR factorization, we may select C having the pivot columns of A as its columns (this solution is unique).

$$C = \boxed{\begin{bmatrix} 1 & 2 \\ 4 & 5 \\ 7 & 8 \end{bmatrix}}$$

(c) Find all matrices S such that $A = S \operatorname{rref}(A)$.

$$\text{We seek } S = \begin{bmatrix} a & x & u \\ b & y & v \\ c & z & w \end{bmatrix}:$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{bmatrix} a & x & u \\ b & y & v \\ c & z & w \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} a & x & 2x - a \\ b & y & 2y - b \\ c & z & 2z - c \end{bmatrix}$$



Hence, we obtain a system of 9 linear equations for the components of S : $a = 1$, $b = 4$, $c = 7$, $x = 2$, $y = 5$, $z = 8$, $2x - a = 3$, $2y - b = 6$, $2z - c = 9$. The first 6 equations determine the values of a, b, c, x, y, z . The last 3 equations are consistent with the determinations. The last 3 variables u, v, w are unconstrained (i.e. free variables). Thus,

$$S = \begin{bmatrix} 1 & 2 & u \\ 4 & 5 & v \\ 7 & 8 & w \end{bmatrix} \quad \text{where } u, v, w \text{ are free parameters.}$$

Check:

$$\begin{bmatrix} 1 & 2 & u \\ 4 & 5 & v \\ 7 & 8 & w \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Extra Credit. (5 points) [Sample Problem B] Showing your work, find the smallest positive integers x_1 , x_2 , x_3 and x_4 that balances the chemical equation for the combustion of ethanol:



$$\begin{array}{c} \text{C} \\ \text{H} \\ \text{O} \end{array} \quad x_1 \begin{bmatrix} 2 \\ 6 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix} = x_3 \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} + x_4 \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$$

$$\bullet \begin{array}{c} 3 \\ \frac{1}{2} \end{array} \begin{bmatrix} 2 & 0 & -1 & 0 & 0 \\ 6 & 0 & 0 & -2 & 0 \\ 1 & 2 & -2 & -1 & 0 \end{bmatrix} \sim \begin{array}{c} \downarrow \\ \uparrow \end{array} \begin{bmatrix} 2 & 0 & -1 & 0 & 0 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 2 & -\frac{3}{2} & -1 & 0 \end{bmatrix} \sim \begin{array}{c} \times \frac{1}{2} \\ \times \frac{1}{2} \\ \times \frac{1}{3} \end{array} \begin{bmatrix} 2 & 0 & -1 & 0 & 0 \\ 0 & 2 & -\frac{3}{2} & -1 & 0 \\ 0 & 0 & 3 & -2 & 0 \end{bmatrix} \sim$$

$$\begin{array}{c} -\frac{1}{2} \\ -\frac{3}{4} \\ \bullet \end{array} \begin{bmatrix} 1 & 0 & -\frac{1}{2} & 0 & 0 \\ 0 & 1 & -\frac{3}{4} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 & -\frac{2}{3} & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & -\frac{1}{3} & 0 \\ 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & -\frac{2}{3} & 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_4 \begin{bmatrix} \frac{1}{3} \\ 1 \\ \frac{2}{3} \\ 1 \end{bmatrix}$$

$$x_4 = 3 \quad \text{efficiently clears denominators}$$

$$\Rightarrow \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \boxed{\begin{bmatrix} 1 \\ 3 \\ 2 \\ 3 \end{bmatrix}}$$