## EE24BTECH11007 - Arnav Makarand Yadnopavit

## Question:

The area of the region bounded by parabola  $y^2 = x$  and the straight line 2y = x is **Solution:** 

Parameter	Description	Value
V	$x^{\top}Vx + 2u^{\top}x + f = 0$	$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$
и	$x^{T}Vx + 2u^{T}x + f = 0$	$\begin{pmatrix} \frac{-1}{2} \\ 0 \end{pmatrix}$
f	$x^{T}Vx + 2u^{T}x + f = 0$	0
h	Point on the given line	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$
m	Direction vector of given line	$\begin{pmatrix} 2 \\ 1 \end{pmatrix}$

TABLE 0: Values

The point of intersection of the line with the parabola is  $x_i = h + k_i m$ , where,  $k_i$  is a constant and is calculated as follows:-

$$k_{i} = \frac{1}{m^{\top}Vm} \left( -m^{\top} (Vh + u) \pm \sqrt{[m^{\top} (Vh + u)]^{2} - g(h)(m^{\top}Vm)} \right)$$
(0.1)

Substituting the input parameters in  $k_i$ , We get,

$$k_i = 0, 2 \tag{0.2}$$

From Table 0 and Equation 0.1, we get,

$$x_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + (0) \begin{pmatrix} 2 \\ 1 \end{pmatrix} \tag{0.3}$$

$$\implies x_1 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{0.4}$$

$$x_2 = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + (2) \begin{pmatrix} 2 \\ 1 \end{pmatrix} \tag{0.5}$$

$$\implies x_2 = \begin{pmatrix} 4\\2 \end{pmatrix} \tag{0.6}$$

The area bounded by the curve  $y^2 = x$  and line 2y = x is given by

$$\int_0^2 2y - y^2 dy = \left(y^2 - \frac{y^3}{3}\right)_0^2 \tag{0.7}$$

$$= \left(4 - \frac{8}{3}\right) - 0\tag{0.8}$$

$$=\frac{4}{3}\tag{0.9}$$




Fig. 0.1: Plot of Parabola  $y^2 = x$  along with the line 2y = x