

Maxi Brandstetter, Felix Kirschner, Arne Heimendahl

University of Cologne

September 7, 2018

# Outline

## 1 Local and quantum correlation matrices

- Local correlation matrices
- Quantum correlation matrices
- The relations between quantum correlation and local correlation matrices

Nice slide to draw the connection between the games an LC

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

- Set of all local correlation matrices:  $\text{LC}_{m,n}$

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

- Set of all local correlation matrices:  $\text{LC}_{m,n}$

## Lemma

$$\text{LC}_{m,n} = \text{conv}\{\xi \eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

- Set of all local correlation matrices:  $\text{LC}_{m,n}$

## Lemma

$$\text{LC}_{m,n} = \text{conv}\{\xi \eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

- No matter which probabilistic strategy there is a deterministic one which is at least as good as the one one chooses

$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$ .

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )





$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )
- Suffices to show that  $LC_{m,n}$  is convex.



$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )
- Suffices to show that  $LC_{m,n}$  is convex.
- Let  $a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}]$  for  $k \in \{0, 1\}$



$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$ .

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )
- Suffices to show that  $LC_{m,n}$  is convex.
- Let  $a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}]$  for  $k \in \{0, 1\}$
- Find  $(X_i), (Y_j)$  with  $|X_i|, |Y_j| \leq 1$  almost surely such that

$$\beta a_{ij}^{(0)} + (1 - \beta) a_{ij}^{(1)} = \mathbb{E}[X_i Y_j]$$

for  $\beta \in [0, 1]$



$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )
- Suffices to show that  $LC_{m,n}$  is convex.
- Let  $a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}]$  for  $k \in \{0, 1\}$
- Find  $(X_i), (Y_j)$  with  $|X_i|, |Y_j| \leq 1$  almost surely such that

$$\beta a_{ij}^{(0)} + (1 - \beta) a_{ij}^{(1)} = \mathbb{E}[X_i Y_j]$$

for  $\beta \in [0, 1]$

- Define a Bernoulli random variable  $\alpha$  such that  $\mathbb{P}(\alpha = 0) = \beta$ ,  $\mathbb{P}(\alpha = 1) = 1 - \beta$  and set  $X_i = X_i^{(\alpha)}, Y_j = Y_j^{(\alpha)}$



$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- $\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )
- Suffices to show that  $LC_{m,n}$  is convex.
- Let  $a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}]$  for  $k \in \{0, 1\}$
- Find  $(X_i), (Y_j)$  with  $|X_i|, |Y_j| \leq 1$  almost surely such that

$$\beta a_{ij}^{(0)} + (1 - \beta) a_{ij}^{(1)} = \mathbb{E}[X_i Y_j]$$

for  $\beta \in [0, 1]$

- Define a Bernoulli random variable  $\alpha$  such that  $\mathbb{P}(\alpha = 0) = \beta$ ,  $\mathbb{P}(\alpha = 1) = 1 - \beta$  and set  $X_i = X_i^{(\alpha)}, Y_j = Y_j^{(\alpha)}$
- Then

$$\begin{aligned} \mathbb{E}[X_i Y_j] &= \mathbb{E}[X_i^{(\alpha)} Y_j^{(\alpha)} \mathbb{1}_{\{\alpha=0\}}] + \mathbb{E}[X_i^{(\alpha)} Y_j^{(\alpha)} \mathbb{1}_{\{\alpha=1\}}] \\ &= \beta \mathbb{E}[X_i^{(0)} Y_j^{(0)}] + (1 - \beta) \mathbb{E}[X_i^{(1)} Y_j^{(1)}] \end{aligned}$$



$$\text{LC}_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

- Let  $a_{ij} = \mathbb{E}[X_i Y_j]$  for  $\mathbb{R}$ -valued random variables  $(X_i), (Y_j)$  defined on a common probability space  $\Omega$  with  $|X_i|, |Y_j| \leq 1$  almost surely.

$$\text{LC}_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

- Let  $a_{ij} = \mathbb{E}[X_i Y_j]$  for  $\mathbb{R}$ -valued random variables  $(X_i), (Y_j)$  defined on a common probability space  $\Omega$  with  $|X_i|, |Y_j| \leq 1$  almost surely.
- Set  $X = (X_1, \dots, X_m)$  and  $Y = (Y_1, \dots, Y_n)$ , then  $X \in [-1, 1]^m$ ,  $Y \in [-1, 1]^n$  almost surely.

$$LC_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

- Let  $a_{ij} = \mathbb{E}[X_i Y_j]$  for  $\mathbb{R}$ -valued random variables  $(X_i), (Y_j)$  defined on a common probability space  $\Omega$  with  $|X_i|, |Y_j| \leq 1$  almost surely.
- Set  $X = (X_1, \dots, X_m)$  and  $Y = (Y_1, \dots, Y_n)$ , then  $X \in [-1, 1]^m$ ,  $Y \in [-1, 1]^n$  almost surely.
- Hypercube description by its vertices:  $[-1, 1]^d = \text{conv}\{\xi \mid \xi \in \{-1, 1\}^d\}$



$$LC_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

- Let  $a_{ij} = \mathbb{E}[X_i Y_j]$  for  $\mathbb{R}$ -valued random variables  $(X_i), (Y_j)$  defined on a common probability space  $\Omega$  with  $|X_i|, |Y_j| \leq 1$  almost surely.
- Set  $X = (X_1, \dots, X_m)$  and  $Y = (Y_1, \dots, Y_n)$ , then  $X \in [-1, 1]^m$ ,  $Y \in [-1, 1]^n$  almost surely.
- Hypercube description by its vertices:  $[-1, 1]^d = \text{conv}\{\xi \mid \xi \in \{-1, 1\}^d\}$
- Define random variables  $\lambda_\xi^{(X)} : \Omega^m \rightarrow [0, 1]$  such that

$$X(\omega) = \sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) \xi$$

almost surely and  $\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) = 1$

$$LC_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- Define random variables  $\lambda_\xi^{(X)} : \Omega^m \rightarrow [0, 1]$  such that

$$X(\omega) = \sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) \xi$$

almost surely and  $\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) = 1$



$$LC_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- Define random variables  $\lambda_\xi^{(X)} : \Omega^m \rightarrow [0, 1]$  such that

$$X(\omega) = \sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) \xi$$

almost surely and  $\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) = 1$

- Using the same decomposition for  $Y$  we obtain

$$\begin{aligned} a_{ij} &= \mathbb{E}[X_i Y_j] = \mathbb{E}\left[\left(\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)} \xi_i\right) \left(\sum_{\eta \in \{-1, 1\}^n} \lambda_\eta^{(Y)} \eta_j\right)\right] \\ &= \sum_{\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n} \mathbb{E}[\lambda_\xi^{(X)} \lambda_\eta^{(Y)}] \xi_i \eta_j \\ &= \left(\sum_{\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n} \mathbb{E}[\lambda_\xi^{(X)}] \mathbb{E}[\lambda_\eta^{(Y)}]\right) \xi_i \eta_j \end{aligned}$$



$$LC_{m,n} \subset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

- Define random variables  $\lambda_\xi^{(X)} : \Omega^m \rightarrow [0, 1]$  such that

$$X(\omega) = \sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) \xi$$

almost surely and  $\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)}(\omega) = 1$

- Using the same decomposition for  $Y$  we obtain

$$\begin{aligned} a_{ij} &= \mathbb{E}[X_i Y_j] = \mathbb{E}\left[\left(\sum_{\xi \in \{-1, 1\}^m} \lambda_\xi^{(X)} \xi_i\right) \left(\sum_{\eta \in \{-1, 1\}^n} \lambda_\eta^{(Y)} \eta_j\right)\right] \\ &= \sum_{\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n} \mathbb{E}[\lambda_\xi^{(X)} \lambda_\eta^{(Y)}] \xi_i \eta_j \\ &= \left(\sum_{\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n} \mathbb{E}[\lambda_\xi^{(X)}] \mathbb{E}[\lambda_\eta^{(Y)}]\right) \xi_i \eta_j \end{aligned}$$

- $\sum_{\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n} \mathbb{E}[\lambda_\xi^{(X)}] \mathbb{E}[\lambda_\eta^{(Y)}] = 1$  the matrix  $(a_{ij})$  is a convex combination of  $\xi\eta^T$ ,  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$



Some nice frame to connect QCs to the games

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be self-adjoint operators on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  for some positive integers  $d_1, d_2$ , satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ .  $A = (a_{ij})$  is called *quantum correlation matrix* if there exists a state **Introduce a symbol to define operators from one space to another**  $\rho \in D(\mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2})$  such that

$$a_{ij} = \text{Tr } \rho(X_i \otimes Y_j).$$

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be self-adjoint operators on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  for some positive integers  $d_1, d_2$ , satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ .  $A = (a_{ij})$  is called *quantum correlation matrix* if there exists a state **Introduce a symbol to define operators from one space to another**  $\rho \in D(\mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2})$  such that

$$a_{ij} = \text{Tr } \rho(X_i \otimes Y_j).$$

- Set of all quantum correlation matrices denoted by  $\text{QC}_{m,n}$

## Lemma

$$\text{QC}_{m,n} = \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\},$$

$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- $a_{ij} = \text{Tr } \rho X_i \otimes Y_j$ , state  $\rho$  on a Hilbert space  $\mathcal{H} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$  and Hermitian operators  $(X_i)_{1 \leq i \leq m}$ ,  $(Y_j)_{1 \leq j \leq n}$  on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- $a_{ij} = \text{Tr } \rho X_i \otimes Y_j$ , state  $\rho$  on a Hilbert space  $\mathcal{H} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$  and Hermitian operators  $(X_i)_{1 \leq i \leq m}$ ,  $(Y_j)_{1 \leq j \leq n}$  on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$
- Define a positive semidefinite symmetric bilinear form on the space of Hermitian operators on  $\mathcal{H}$  by  $\beta : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$  where  $\beta(S, T) = \text{Re}(\text{Tr } \rho ST)$ .

$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- $a_{ij} = \text{Tr } \rho X_i \otimes Y_j$ , state  $\rho$  on a Hilbert space  $\mathcal{H} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$  and Hermitian operators  $(X_i)_{1 \leq i \leq m}$ ,  $(Y_j)_{1 \leq j \leq n}$  on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$
- Define a positive semidefinite symmetric bilinear form on the space of Hermitian operators on  $\mathcal{H}$  by  $\beta : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$  where  $\beta(S, T) = \text{Re}(\text{Tr } \rho ST)$ .
- perhaps verification of at least some of these properties

$$QC_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- $a_{ij} = \text{Tr } \rho X_i \otimes Y_j$ , state  $\rho$  on a Hilbert space  $\mathcal{H} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$  and Hermitian operators  $(X_i)_{1 \leq i \leq m}$ ,  $(Y_j)_{1 \leq j \leq n}$  on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$
- Define a positive semidefinite symmetric bilinear form on the space of Hermitian operators on  $\mathcal{H}$  by  $\beta : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$  where  $\beta(S, T) = \text{Re}(\text{Tr } \rho ST)$ .
- perhaps verification of at least some of these properties
- Obtain an inner product space  $U := B^{sa}(\mathcal{H}) / \ker \beta$  equipped with the inner product

$$\tilde{\beta}([S], [T]) = \beta(S, T).$$

$$QC_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- $a_{ij} = \text{Tr } \rho X_i \otimes Y_j$ , take  $\rho$  on a Hilbert space  $\mathcal{H} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2}$  and Hermitian operators  $(X_i)_{1 \leq i \leq m}$ ,  $(Y_j)_{1 \leq j \leq n}$  on  $\mathbb{C}^{d_1}$ , respectively  $\mathbb{C}^{d_2}$  satisfying  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$
- Define a positive semidefinite symmetric bilinear form on the space of Hermitian operators on  $\mathcal{H}$  by  $\beta : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$  where  $\beta(S, T) = \text{Re}(\text{Tr } \rho ST)$ .
- perhaps verification of at least some of these properties
- Obtain an inner product space  $U := B^{sa}(\mathcal{H}) / \ker \beta$  equipped with the inner product

$$\tilde{\beta}([S], [T]) = \beta(S, T).$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )
- Project the  $y_j$ 's orthogonally onto  $\text{span}\{x_1, \dots, x_m\}$  (wlog  $m \leq n$ )



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )
- Project the  $y_j$ 's orthogonally onto  $\text{span}\{x_1, \dots, x_m\}$  (wlog  $m \leq n$ )
- $\pi(y_j)$  the projection of  $y_j$  then  $\tilde{\beta}(x_i, \pi(y_j)) = \tilde{\beta}(x_i, y_j)$





$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )
- Project the  $y_j$ 's orthogonally onto  $\text{span}\{x_1, \dots, x_m\}$  (wlog  $m \leq n$ )
- $\pi(y_j)$  the projection of  $y_j$  then  $\tilde{\beta}(x_i, \pi(y_j)) = \tilde{\beta}(x_i, y_j)$
- Let  $\{a_1, \dots, a_r\}$  be an orthonormal basis of  $\text{span}\{x_1, \dots, x_m\}$  with respect to  $\beta$  and  $x_i = \sum_{k=1}^r \alpha_k^{(i)} a_k$  and  $\pi(y_j) = \sum_{k=1}^r \gamma_k^{(j)} a_k$  for  $\alpha^{(i)}, \gamma^{(j)} \in \mathbb{R}^r$



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )
- Project the  $y_j$ 's orthogonally onto  $\text{span}\{x_1, \dots, x_m\}$  (wlog  $m \leq n$ )
- $\pi(y_j)$  the projection of  $y_j$  then  $\tilde{\beta}(x_i, \pi(y_j)) = \tilde{\beta}(x_i, y_j)$
- Let  $\{a_1, \dots, a_r\}$  be an orthonormal basis of  $\text{span}\{x_1, \dots, x_m\}$  with respect to  $\beta$  and  $x_i = \sum_{k=1}^r \alpha_k^{(i)} a_k$  and  $\pi(y_j) = \sum_{k=1}^r \gamma_k^{(j)} a_k$  for  $\alpha^{(i)}, \gamma^{(j)} \in \mathbb{R}^r$
- Then  $a_{ij} = \tilde{\beta}(x_i, y_j) = \tilde{\beta}(x, \pi(y)) = \sum_{1 \leq k, l \leq r} \alpha_k^{(i)} \gamma_l^{(j)} \tilde{\beta}(a_k, a_l) = \sum_{k=1}^r \alpha_k^{(i)} \gamma_k^{(j)} = \langle \alpha^{(i)}, \gamma^{(j)} \rangle.$



$$\text{QC}_{m,n} \subset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Identify  $X_i \otimes I, I \otimes Y_j$  with vectors  $x_i, y_j$  in  $U$ , then

$$\tilde{\beta}(x_i, y_j) = \beta(X_i, Y_j) = \text{ReTr}(\rho X_i \otimes Y_j) = a_{ij}$$

- $\beta(X \otimes I, X \otimes I), \beta(I \otimes Y, I \otimes Y) \leq 1$   
(this can be shown by using a *Schmidt-decomposition* of  $\rho$  and using  $\|X_i\|_\infty, \|Y_j\|_\infty \leq 1$ )
- Project the  $y_j$ 's orthogonally onto  $\text{span}\{x_1, \dots, x_m\}$  (wlog  $m \leq n$ )
- $\pi(y_j)$  the projection of  $y_j$  then  $\tilde{\beta}(x_i, \pi(y_j)) = \tilde{\beta}(x_i, y_j)$
- Let  $\{a_1, \dots, a_r\}$  be an orthonormal basis of  $\text{span}\{x_1, \dots, x_m\}$  with respect to  $\beta$  and  $x_i = \sum_{k=1}^r \alpha_k^{(i)} a_k$  and  $\pi(y_j) = \sum_{k=1}^r \gamma_k^{(j)} a_k$  for  $\alpha^{(i)}, \gamma^{(j)} \in \mathbb{R}^r$
- Then  $a_{ij} = \tilde{\beta}(x_i, y_j) = \tilde{\beta}(x, \pi(y)) = \sum_{1 \leq k, l \leq r} \alpha_k^{(i)} \gamma_l^{(j)} \tilde{\beta}(a_k, a_l) = \sum_{k=1}^r \alpha_k^{(i)} \gamma_k^{(j)} = \langle \alpha^{(i)}, \gamma^{(j)} \rangle.$
- $|\alpha^{(i)}|, |\gamma^{(j)}| \leq 1$  due to  $\tilde{\beta}(x_i), \tilde{\beta}(y_j) \leq 1$



- In order to show

$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

we will use the following

- In order to show

$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

we will use the following

### Proposition

For all  $n \geq 1$  there is a subspace of the  $2^n \times 2^n$  Hermitian matrices where every vector is the multiple of a unitary matrix.

- In order to show

$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

we will use the following

### Proposition

For all  $n \geq 1$  there is a subspace of the  $2^n \times 2^n$  Hermitian matrices where every vector is the multiple of a unitary matrix.

- The proof is based on  $n$ -fold tensor products of the Pauli matrices which are the three matrices

$$X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

- $X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ ,  $Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ ,  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

## Proof.

- Define

$$U_i = I^{\otimes(i-1)} \otimes X \otimes Y^{\otimes(n-i)},$$

$$U_{n+i} = I^{\otimes(i-1)} \otimes Z \otimes Y^{\otimes(n-i)}, \quad i = 1, \dots, n$$

- $U_i$ 's are anti-commuting traceless Hermitian unitaries, i.e.  $U_i U_j = -U_j U_i$  for  $i \neq j$  and  $U_i^2 = I$
- For  $X = \sum_{i=1}^{2n} \xi_i U_i$ ,  $Y = \sum_{i=1}^{2n} \eta_i U_i$  we can calculate

$$\begin{aligned} XY &= \sum_{i=1}^{2n} \xi_i \eta_i I + \sum_{1 \leq i, j \leq 2n} \xi_i \eta_j U_i U_j \\ &= \sum_{i=1}^{2n} \xi_i \eta_i I + \sum_{1 \leq i < j \leq 2n} \xi_i \eta_j U_i U_j - \sum_{1 \leq i < j \leq 2n} U_i U_j = \sum_{i=1}^{2n} \xi_i \eta_i I \\ &= \langle \xi, \eta \rangle I. \end{aligned}$$

- The result follows by setting  $X = Y$ .

$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- Let  $(x_i)_{1 \leq i \leq m}, (y_j)_{1 \leq j \leq n} \subset \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j| \leq 1$ .



$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- Let  $(x_i)_{1 \leq i \leq m}, (y_j)_{1 \leq j \leq n} \subset \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j| \leq 1$ .
- Let  $X_i = \sum_{k=1}^{\min\{m,n\}} x_i(k) U_i$  and  $Y_j^T = \sum_{k=1}^{\min\{m,n\}} y_j(k) U_k$  where the  $U_i$ 's are  $d \times d$  matrices with  $d = 2^{\lceil \min\{m,n\}/2 \rceil}$

$$\mathcal{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- Let  $(x_i)_{1 \leq i \leq m}, (y_j)_{1 \leq j \leq n} \subset \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j| \leq 1$ .
- Let  $X_i = \sum_{k=1}^{\min\{m,n\}} x_i(k) U_k$  and  $Y_j^T = \sum_{k=1}^{\min\{m,n\}} y_j(k) U_k$  where the  $U_i$ 's are  $d \times d$  matrices with  $d = 2^{\lceil \min\{m,n\}/2 \rceil}$
- $\text{Tr}(X_i Y_j^T) = d \cdot \langle x_i, y_j \rangle$  and  $\|X_i\|_\infty \leq 1$  since  $X_i X_i^* = |x_i|^2 I$  and  $|x_i|^2 \leq 1$  (the same holds for  $Y_j$ )

$$\mathcal{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}$$

- Let  $(x_i)_{1 \leq i \leq m}, (y_j)_{1 \leq j \leq n} \subset \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j| \leq 1$ .
- Let  $X_i = \sum_{k=1}^{\min\{m,n\}} x_i(k) U_k$  and  $Y_j^T = \sum_{k=1}^{\min\{m,n\}} y_j(k) U_k$  where the  $U_i$ 's are  $d \times d$  matrices with  $d = 2^{\lceil \min\{m,n\}/2 \rceil}$
- $\text{Tr}(X_i Y_j^T) = d \cdot \langle x_i, y_j \rangle$  and  $\|X_i\|_\infty \leq 1$  since  $X_i X_i^* = |x_i|^2 I$  and  $|x_i|^2 \leq 1$  (the same holds for  $Y_j$ )
- Let  $|\phi\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle$  and  $\rho = |\phi\rangle \langle \phi|$ . Note that we can write  $\rho$  as

$$\rho = |\phi\rangle \langle \phi| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |kk\rangle \langle ll| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |k\rangle \langle l| \otimes |k\rangle \langle l|$$

$$\text{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Let  $|\phi\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle$  and  $\rho = |\phi\rangle \langle \rho|$ . Note that we can write  $\rho$  as

$$\rho = |\phi\rangle \langle \phi| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |kk\rangle \langle ll| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |k\rangle \langle l| \otimes |k\rangle \langle l|$$



$$\mathcal{QC}_{m,n} \supset \{(\langle x_i, y_j \rangle)_{1 \leq i \leq m, 1 \leq j \leq n} \mid x_i, y_j \in \mathbb{R}^{\min\{m,n\}}, |x_i| \leq 1, |y_j| \leq 1\}.$$

- Let  $|\phi\rangle = \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle$  and  $\rho = |\phi\rangle \langle \phi|$ . Note that we can write  $\rho$  as

$$\rho = |\phi\rangle \langle \phi| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |kk\rangle \langle ll| = \frac{1}{d} \sum_{1 \leq k, l \leq d} |k\rangle \langle l| \otimes |k\rangle \langle l|$$

- Then

$$\begin{aligned} \text{Tr}(\rho X_i \otimes Y_j) &= \frac{1}{d} \sum_{1 \leq k, l \leq d} \text{Tr}(|k\rangle \langle l| X_i \otimes |k\rangle \langle l| Y_j) = \frac{1}{d} \sum_{1 \leq k, l \leq d} \text{Tr}(|k\rangle \langle l| X_i) \text{Tr}(|k\rangle \langle l| Y_j) \\ &= \frac{1}{d} \text{Tr} X_i Y_j^T = \langle x_i, y_j \rangle. \end{aligned}$$



- $QC_{m,n}$  is convex

- $\text{QC}_{m,n}$  is convex
- Let  $a_{ij} = \langle x_i, y_j \rangle$  and  $\bar{a}_{ij} = \langle \bar{x}_i, \bar{y}_j \rangle$  for  $x_i, y_j, \bar{x}_i, \bar{y}_j \in \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j|, |\bar{x}_i|, |\bar{y}_j| \leq 1$ .

- $QC_{m,n}$  is convex
- Let  $a_{ij} = \langle x_i, y_j \rangle$  and  $\bar{a}_{ij} = \langle \bar{x}_i, \bar{y}_j \rangle$  for  $x_i, y_j, \bar{x}_i, \bar{y}_j \in \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j|, |\bar{x}_i|, |\bar{y}_j| \leq 1$ .
- define vectors  $\tilde{x}_i := (\sqrt{\lambda}x_i, \sqrt{1-\lambda}\bar{x}_i)$ ,  $\tilde{y}_j := (\sqrt{\lambda}y_j, \sqrt{1-\lambda}\bar{y}_j)$  for  $\lambda \in [0, 1]$



- $\text{QC}_{m,n}$  is convex
- Let  $a_{ij} = \langle x_i, y_j \rangle$  and  $\bar{a}_{ij} = \langle \bar{x}_i, \bar{y}_j \rangle$  for  $x_i, y_j, \bar{x}_i, \bar{y}_j \in \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j|, |\bar{x}_i|, |\bar{y}_j| \leq 1$ .
- define vectors  $\tilde{x}_i := (\sqrt{\lambda}x_i, \sqrt{1-\lambda}\bar{x}_i)$ ,  $\tilde{y}_j := (\sqrt{\lambda}y_j, \sqrt{1-\lambda}\bar{y}_j)$  for  $\lambda \in [0, 1]$
- it holds  $|\tilde{x}_i| \leq \lambda|(x_i, 0)| + (1-\lambda)|(0, \bar{x}_i)| \leq 1$  and  $\langle \tilde{x}_i, \tilde{y}_j \rangle = \lambda \langle x_i, y_j \rangle + (1-\lambda) \langle \bar{x}_i, \bar{y}_j \rangle$ .

- $\text{QC}_{m,n}$  is convex
- Let  $a_{ij} = \langle x_i, y_j \rangle$  and  $\bar{a}_{ij} = \langle \bar{x}_i, \bar{y}_j \rangle$  for  $x_i, y_j, \bar{x}_i, \bar{y}_j \in \mathbb{R}^{\min\{m,n\}}$  such that  $|x_i|, |y_j|, |\bar{x}_i|, |\bar{y}_j| \leq 1$ .
- define vectors  $\tilde{x}_i := (\sqrt{\lambda}x_i, \sqrt{1-\lambda}\bar{x}_i)$ ,  $\tilde{y}_j := (\sqrt{\lambda}y_j, \sqrt{1-\lambda}\bar{y}_j)$  for  $\lambda \in [0, 1]$
- it holds  $|\tilde{x}_i| \leq \lambda|(x_i, 0)| + (1-\lambda)|(0, \bar{x}_i)| \leq 1$  and  $\langle \tilde{x}_i, \tilde{y}_j \rangle = \lambda \langle x_i, y_j \rangle + (1-\lambda) \langle \bar{x}_i, \bar{y}_j \rangle$ .
- Right dimension is obtained by projecting  $(\tilde{x}_i)_{1 \leq i \leq m}$ ,  $(\tilde{y}_j)_{1 \leq j \leq n}$  on  $\text{span}\{x_1, \dots, x_m\}$  or  $\text{span}\{y_1, \dots, y_n\}$ , as in the proof before.

- $\text{LC}_{m,n} \subset \text{QC}_{m,n}$

- $LC_{m,n} \subset QC_{m,n}$
- Set  $x_i = \xi_i |0\rangle$  and  $y_j = \eta_j |0\rangle$  it immediately follows  $\xi_i \eta_j = \langle x_i, y_j \rangle$ . Hence,  $\xi \eta^T \in QC_{m,n}$  (rest follows with the convexity of  $QC_{m,n}$ )

- $LC_{m,n} \subset QC_{m,n}$
- Set  $x_i = \xi_i |0\rangle$  and  $y_j = \eta_j |0\rangle$  it immediately follows  $\xi_i \eta_j = \langle x_i, y_j \rangle$ . Hence,  $\xi \eta^T \in QC_{m,n}$  (rest follows with the convexity of  $QC_{m,n}$ )
- Inclusion is strict in general

- Let us consider the case  $n = m = 2$ .

- Let us consider the case  $n = m = 2$ .
- It holds

$$\text{LC}_{2,2} = \text{conv}\left\{\pm \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \pm \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix}, \pm \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}, \pm \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}\right\}.$$

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (1)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .



## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

- $\text{LC}_{2,2}$  affinely isomorphic to the cross polytope scaled by 2, i.e.  
 $\text{LC}_{2,2} \cong 2CP_4 := \text{conv}\{\pm e_i \mid i = 1, \dots, 4\}$

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

- $\text{LC}_{2,2}$  affinely isomorphic to the cross polytope scaled by 2, i.e.  
 $\text{LC}_{2,2} \cong 2CP_4 := \text{conv}\{\pm e_i \mid i = 1, \dots, 4\}$
- $(CP_n)^o = [-1, 1]^n$

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

- $\text{LC}_{2,2}$  affinely isomorphic to the cross polytope scaled by 2, i.e.  
 $\text{LC}_{2,2} \cong 2CP_4 := \text{conv}\{\pm e_i \mid i = 1, \dots, 4\}$
- $(CP_n)^{\circ} = [-1, 1]^n$
- Both points implies that the face lattice of  $\text{LC}_{2,2}$  is isomorphic to the opposite lattice of  $[-1, 1]^4$

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

- $\text{LC}_{2,2}$  affinely isomorphic to the cross polytope scaled by 2, i.e.  
 $\text{LC}_{2,2} \cong 2CP_4 := \text{conv}\{\pm e_i \mid i = 1, \dots, 4\}$
- $(CP_n)^{\circ} = [-1, 1]^n$
- Both points implies that the face lattice of  $\text{LC}_{2,2}$  is isomorphic to the opposite lattice of  $[-1, 1]^4$
- $\text{LC}_{2,2}$  has as many facets as  $[-1, 1]^4$  vertices, hence  $2^4$

## Claim

$$\text{LC}_{2,2} = \{A \in \mathbb{R}^{2 \times 2} \mid -1 \leq \text{Tr } AM \leq 1 \text{ for all } M \in \mathcal{K}\}, \quad (2)$$

where  $\mathcal{K} = \{\frac{1}{2}\sigma(\begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix}), \sigma(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}) \mid \sigma \in \{\text{id}(1\ 2), (1\ 3), (1\ 4)\}\}$ .

- $\text{LC}_{2,2}$  affinely isomorphic to the cross polytope scaled by 2, i.e.  
 $\text{LC}_{2,2} \cong 2CP_4 := \text{conv}\{\pm e_i \mid i = 1, \dots, 4\}$
- $(CP_n)^{\circ} = [-1, 1]^n$
- Both points implies that the face lattice of  $\text{LC}_{2,2}$  is isomorphic to the opposite lattice of  $[-1, 1]^4$
- $\text{LC}_{2,2}$  has as many facets as  $[-1, 1]^4$  vertices, hence  $2^4$
- Equation 2 is a non-redundant hyperplane description of  $\text{LC}_{2,2}$ , hence all linear constraints define facets.

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
- $\max\{\text{Tr}(AM) \mid A \in \text{LC}_{2,2}\} = 2$



- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
- $\max\{\text{Tr}(AM) \mid A \in \text{LC}_{2,2}\} = 2$
- $A \in \text{QC}_{2,2}$  we obtain, by Cauchy-Schwarz and  $|y_i| \leq 1$ ,

$$\begin{aligned} \text{Tr}(AM) &= \langle x_1, y_1 \rangle + \langle x_1, y_2 \rangle + \langle x_2, y_1 \rangle - \langle x_2, y_2 \rangle \\ &= \langle x_1 + x_2, y_1 \rangle + \langle x_1 - x_2, y_2 \rangle \leq |x_1 + x_2| |y_1| + |x_1 - x_2| |y_2| \\ &\leq |x_1 + x_2| + |x_1 - x_2|. \end{aligned}$$

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
- $\max\{\text{Tr}(AM) \mid A \in \text{LC}_{2,2}\} = 2$
- $A \in \text{QC}_{2,2}$  we obtain, by Cauchy-Schwarz and  $|y_i| \leq 1$ ,

$$\begin{aligned} \text{Tr}(AM) &= \langle x_1, y_1 \rangle + \langle x_1, y_2 \rangle + \langle x_2, y_1 \rangle - \langle x_2, y_2 \rangle \\ &= \langle x_1 + x_2, y_1 \rangle + \langle x_1 - x_2, y_2 \rangle \leq |x_1 + x_2||y_1| + |x_1 - x_2||y_2| \\ &\leq |x_1 + x_2| + |x_1 - x_2|. \end{aligned}$$

- $(|x_1 + x_2| + |x_1 - x_2|)^2 \leq 4(|x_1|^2 + |x_2|^2)$

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
- $\max\{\text{Tr}(AM) \mid A \in \text{LC}_{2,2}\} = 2$
- $A \in \text{QC}_{2,2}$  we obtain, by Cauchy-Schwarz and  $|y_i| \leq 1$ ,

$$\begin{aligned} \text{Tr}(AM) &= \langle x_1, y_1 \rangle + \langle x_1, y_2 \rangle + \langle x_2, y_1 \rangle - \langle x_2, y_2 \rangle \\ &= \langle x_1 + x_2, y_1 \rangle + \langle x_1 - x_2, y_2 \rangle \leq |x_1 + x_2||y_1| + |x_1 - x_2||y_2| \\ &\leq |x_1 + x_2| + |x_1 - x_2|. \end{aligned}$$

- $(|x_1 + x_2| + |x_1 - x_2|)^2 \leq 4(|x_1|^2 + |x_2|^2)$
- $\text{Tr}(AM) \leq |x_1 + x_2| + |x_1 - x_2| \leq 2\sqrt{|x_1|^2 + |x_2|^2} \leq 2\sqrt{2}.$

- We will show that the inclusion is strict by showing that both sets yield different values if we maximize in a certain direction
- Let  $M = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$
- $\max\{\text{Tr}(AM) \mid A \in \text{LC}_{2,2}\} = 2$
- $A \in \text{QC}_{2,2}$  we obtain, by Cauchy-Schwarz and  $|y_i| \leq 1$ ,

$$\begin{aligned} \text{Tr}(AM) &= \langle x_1, y_1 \rangle + \langle x_1, y_2 \rangle + \langle x_2, y_1 \rangle - \langle x_2, y_2 \rangle \\ &= \langle x_1 + x_2, y_1 \rangle + \langle x_1 - x_2, y_2 \rangle \leq |x_1 + x_2||y_1| + |x_1 - x_2||y_2| \\ &\leq |x_1 + x_2| + |x_1 - x_2|. \end{aligned}$$

- $(|x_1 + x_2| + |x_1 - x_2|)^2 \leq 4(|x_1|^2 + |x_2|^2)$
- $\text{Tr}(AM) \leq |x_1 + x_2| + |x_1 - x_2| \leq 2\sqrt{|x_1|^2 + |x_2|^2} \leq 2\sqrt{2}$ .
- Bound is achieved by  $A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ , induced by the vectors  $x_1 = x_2 = \frac{1}{\sqrt{2}}(1, 1)$  and  $y_1 = y_2 = (1, 0)$