

Maxi Brandstetter, Felix Kirschner, Arne Heimendahl

University of Cologne

September 6, 2018

# Outline

- 1 Local and quantum correlation matrices
  - Local correlation matrices

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

Set of all local correlation matrices:  $\text{LC}_{m,n}$

## Definition

Let  $(X_i)_{1 \leq i \leq m}$  and  $(Y_j)_{1 \leq j \leq n}$  be families of random variables on a common probability space such that  $|X_i|, |Y_j| \leq 1$  almost surely. Then  $A = (a_{ij})$  is the corresponding *classical (or local) correlation matrix* if

$$a_{ij} = \mathbb{E}[X_i Y_j]$$

for all  $1 \leq i \leq m, 1 \leq j \leq n$ .

Set of all local correlation matrices:  $\text{LC}_{m,n}$

## Lemma

$$\text{LC}_{m,n} = \text{conv}\{\xi \eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

No matter which probabilistic strategy there is a deterministic one which is at least as good as the one one chooses

## Lemma

$$\text{LC}_{m,n} = \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

$$\text{LC}_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

## Lemma

$$LC_{m,n} = \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

$\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )

## Lemma

$$LC_{m,n} = \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

$\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )

Suffices to show that  $LC_{m,n}$  is convex.

$$a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}] \text{ for } k \in \{0, 1\}$$

Find  $(X_i), (Y_j)$  with  $|X_i|, |Y_j| \leq 1$  almost surely such that

$$\beta a_{ij}^{(0)} + (1 - \beta) a_{ij}^{(1)} = \mathbb{E}[X_i Y_j]$$

for  $\beta \in [0, 1]$



## Lemma

$$LC_{m,n} = \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}$$

$$LC_{m,n} \supset \text{conv}\{\xi\eta^T \mid \xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n\}.$$

$\xi\eta^T \in LC_{m,n}$  for all  $\xi \in \{-1, 1\}^m, \eta \in \{-1, 1\}^n$  (Choose  $X_i \equiv \xi_i, Y_j \equiv \eta_j$ )

Suffices to show that  $LC_{m,n}$  is convex.

$$a_{ij}^{(k)} = \mathbb{E}[X_i^{(k)} Y_j^{(k)}] \text{ for } k \in \{0, 1\}$$

Find  $(X_i), (Y_j)$  with  $|X_i|, |Y_j| \leq 1$  almost surely such that

$$\beta a_{ij}^{(0)} + (1 - \beta) a_{ij}^{(1)} = \mathbb{E}[X_i Y_j]$$

for  $\beta \in [0, 1]$

Define a Bernoulli random variable  $\alpha$  such that  $\mathbb{P}(\alpha = 0) = \beta$ ,  $\mathbb{P}(\alpha = 1) = 1 - \beta$  and set  $X_i = X_i^{(\alpha)}, Y_j = Y_j^{(\alpha)}$