ABC formula for the HP-11C (and HP-15C)

A.J.M. Jacobs

March 5, 2013

Contents

1	The ABC formula HP-11C code	2
2	How to use the ABC formula	4

1 The ABC formula HP-11C code

key strokes	step	display code(s)	remark
ON g P/R f CLEAR PRGM			Put the calculator ON Enter program mode Clear program space is needed
f LBL A	01	42,21,11	Start position ABC formula
STO 3	02	44 3	Save 'c'
$R\!\!\downarrow$	03	33	
CHS	04	16	
STO 2	05	44 2	Save '-b'
R↓	06	33	
2	07	2	
×	80	20	
STO 1	09	44 1	Save '2a'
2	10	2	
×	11	20	'4a'
RCL 3	12	45 3	
×	13	20	'4ac'
RCL 2	14	45 2	
$\lceil g \rceil \ x^2$	15	43 11	'b*b'
-	16	30	
CHS	17	16	'd = b*b - 4ac'
g X<0	18	43 10	If $d < 0$ then NO solutions
GTO 1	19	22 1	

key strokes	step	<pre>display code(s)</pre>	remark
\sqrt{x}	20	11	Square root of 'd'
STO 0	21	44 0	Save square root of 'd'
RCL 2	22	45 2	
+	23	40	
RCL 1	24	45 1	
÷	25	10	+ ABC formula
STO 4	26	44 4	Save x1
RCL 2	27	45 2	
RCL 0	28	45 0	
-	29	30	
RCL 1	30	45 1	
÷	31	10	- ABC formula
STO 5	32	44 5	Save x2
RCL 4	33	45 4	Read x1
g RTN	34	43 32	Return with x1 in X and x2 in Y
f LBL 1	35	42,21, 1	The are no real value solutions
0	36	0	
STO 4	37	44 4	Save 0 in x1
STO 5	38	44 5	Save 0 in x2
ENTER	39	36	
g RTN	40	43 32	Return with 0 in X and 0 in Y, no real value solutions
			,

2 How to use the ABC formula

A quadratic equation with real or complex coefficients has two solutions, called roots. These two solutions may or may not be distinct, and they may or may not be real. With this program only real solutions are calculated.

Having:

$$ax^2 + bx + c = 0$$

The RPN coded ABC formula starts with Label "A". If Label "A" is the first step the program can be started with the R/S (Run/Stop) button. Enter the values a, b and c and execute the function by pressing the R/S key.

Example with a = 1, b = -5 and c = 4:

Keystrokes:

1

ENTER

5

CHS

ENTER

4

R/S

running

Result: $4.0000 x \leq y 1.0000$

So the solution is: $x_1 = 1$ and $x_2 = 4$