(a) If $u = \tan^{-1} \left(\frac{x^3 + y^3}{x - y} \right)$ prove that ۲.

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \sin 2u$$

$$x^{2} \frac{\partial^{2} u}{\partial x^{2}} + y^{2} \frac{\partial^{2} u}{\partial y^{2}} + 2xy \frac{\partial^{2} u}{\partial x \partial y} = 2 \cos 3u \sin u.$$

have a volume of 32 c.c. Find the dimensions of the box requiring least material for its A rectangular box open at the top is to construction. **(**P)

- Find the area lying between the parabola $y = 4x - x^2$ and the line y = x. (a
 - Find, by double integration, the volume of the solid

generated by revolving the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

about the y-axis.

Find the volume of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

Prove that **(**e)

$$\beta(m, n) = \frac{\Gamma m \Gamma n}{\Gamma m + n}$$
, hence evaluate Γ 1/2.

B. Tech. 1st Semester F-Scheme Examination,

December-2016

MATHEMATICS-I

Paper-MATH-101-F

[Maximum marks : 100 Time allowed: 3 hours]

five questions with selecting one question from each unit. All questions carry equal Note: Question No. I is compulsory. Attempt total

(a) Discuss the behavior of series

$$\frac{\sqrt{2}-1}{3^3-1} + \frac{\sqrt{3}-1}{4^3-1} + \frac{\sqrt{4}-1}{5^3-1} + \frac{\sqrt{5}-1}{6^3-1} + \dots$$

Prove that **((2**)

$$\log (1 + x) = x - \frac{x^2}{2} - \frac{x^3}{3} + \frac{x^4}{4}$$

(c) Find asymptotes parallel to co-ordinate axis of the

$$x^2 y^2 - x^2 y - xy^2 + x + y + 1 = 0$$

(d) Define rank of matrix. Find rank of matrix

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ -2 & 0 & 5 & 7 \end{bmatrix}$$

(e) Using Cayley-Hamilton theorem, find A⁶

- Write the relationship between Cartesian coords and cylindrical polar coords. Also write relationship between Cartesian coords and Spherical polar coords. 9
 - Define Beta and Gamma function. (g)
- State Leibnitz theorem for the nth derivative of the product of two functions.

- Examine the convergence of the series whose nth term is $\sqrt{n^4 + 1} - \sqrt{n^4 - 1}$. (a) 7
 - Discuss the convergence of the series (q)

$$\frac{2}{1}x + \frac{9}{8}x^2 + \frac{64}{81}x^3 + \dots$$
 to ∞

Test for absolute/conditionally convergence of the (a) 3

series
$$2 - \frac{4}{2!} + \frac{8}{3!} - \frac{16}{4!} + \frac{32}{5!}$$

For what values of x are the following series

convergent: $\frac{1}{1-x} + \frac{1}{2(1-x)^2} + \frac{1}{3(1-x)^3} + \dots \infty$

Find non-singular matrices P and Q such that PAQ is in the normal form for the

$$A = \begin{bmatrix} 0 & 2 & 3 & 4 \\ 2 & 3 & 5 & 4 \\ 4 & 8 & 13 & 12 \end{bmatrix}$$

Find the value of λ such that the following equations have unique solution:

$$\lambda x + 2y - 2z - 1 = 0$$
, $4x + 2\lambda y - z - 2 = 0$, $6x + 6y + \lambda z - 3 = 0$

and use matrix method to solve these equations when $\lambda = 2$

- If λ is an eigen value of a non-singular matrix A, show that (a)
- (i) $\lambda 1$ is an eigen value of A 1
- (ii) $\frac{|A|}{2}$ is an eigen value of adj. A
 - (b) Show that the matrix

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

satisfies its own characteristic equation and $\begin{bmatrix} 0 & 0 & -1 \end{bmatrix}$ or otherwise find A⁻¹.

If log y = tan-1 x, show that (a) 9

 $(1+x^2)y_{n+2} + \{2(n+1)x-1\}y_{n+1} + n(n+1)y_n = 0$ and hence find y_3 , y_4 and y_5 at x = 0.

on an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is given by $p = UD^3$ where CD is the semi-diameter conjust Show that the radius of curvature p 9

Expand $e^x \log (1 + y)$ in powers of \times and y upto erms of third degree. (3)