

Lax-Wendroff Flux Reconstruction for hyperbolic conservation laws

March 17, 2023

Arpit Babbar

`arpit@tifrbng.res.in`



TIFR-CAM, Bangalore

Department of Mathematics

IISER Trivandrum


- Review of Finite Volume Method (FVM) and Flux Reconstruction (FR)
- Introduce Lax-Wendroff Flux Reconstruction (LWFR) where numerical fluxes are carefully constructed to improve accuracy and stability
- Introduce a second order variant of blending limiter of Henneman Et Al [8] in context of LW schemes which will be used to create a provably admissibility preserving LW scheme
- Extend LWFR to unstructured, curved meshes with free stream preservation

- Review of Finite Volume Method (FVM) and Flux Reconstruction (FR)
- Introduce Lax-Wendroff Flux Reconstruction (LWFR) where numerical fluxes are carefully constructed to improve accuracy and stability
- Introduce a second order variant of blending limiter of Henneman Et Al [8] in context of LW schemes which will be used to create a provably admissibility preserving LW scheme
- Extend LWFR to unstructured, curved meshes with free stream preservation

- Review of Finite Volume Method (FVM) and Flux Reconstruction (FR)
- Introduce Lax–Wendroff Flux Reconstruction (LWFR) where numerical fluxes are carefully constructed to improve accuracy and stability
- Introduce a second order variant of blending limiter of Henneman Et Al [8] in context of LW schemes which will be used to create a provably admissibility preserving LW scheme
- Extend LWFR to unstructured, curved meshes with free stream preservation


- Review of Finite Volume Method (FVM) and Flux Reconstruction (FR)
- Introduce Lax–Wendroff Flux Reconstruction (LWFR) where numerical fluxes are carefully constructed to improve accuracy and stability
- Introduce a second order variant of blending limiter of Henneman Et Al [8] in context of LW schemes which will be used to create a provably admissibility preserving LW scheme
- Extend LWFR to unstructured, curved meshes with free stream preservation

$$\begin{aligned} u_t + f(u)_x &= 0, \\ u(x, 0) &= u_0(x). \end{aligned}$$




Piecewise constant states

$$\int_{I_e} u_t + \int_{I_e} f(u)_x = 0$$



Piecewise constant states


$$\frac{d\bar{u}_e^n}{dt} + \frac{f_{e+\frac{1}{2}} - f_{e-\frac{1}{2}}}{\Delta x_e} = 0$$



Piecewise constant states

$$\frac{d\bar{u}_e^n}{dt} + \frac{f_{e+\frac{1}{2}} - f_{e-\frac{1}{2}}}{\Delta x_e} = 0$$


$$f_{e+\frac{1}{2}} = \frac{f(\bar{u}_e) + f(\bar{u}_{e+1})}{2} + \max_{u \in I[\bar{u}_e, \bar{u}_{e+1}]} |f'(u)| \frac{\bar{u}_e - \bar{u}_{e+1}}{2}$$



Piecewise constant states

$$\frac{d\bar{u}_e^n}{dt} + \frac{f_{e+\frac{1}{2}} - f_{e-\frac{1}{2}}}{\Delta x_e} = 0$$

$$f_{e+\frac{1}{2}} = \frac{f(\bar{u}_e) + f(\bar{u}_{e+1})}{2} + \max_{u \in I[\bar{u}_e, \bar{u}_{e+1}]} |f'(u)| \frac{\bar{u}_e - \bar{u}_{e+1}}{2}$$



Piecewise constant states

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$


Degree N approximation

$$\Omega = \bigcup_e I_e$$

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$

Degree N approximation


$$\Omega = \bigcup_e I_e$$



$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$



Degree N approximation

$$\Omega = \bigcup_e I_e$$




We use Gauss-Legendre solution points.

$$\frac{d}{dt} u_{e,i} = -\frac{\partial f_h}{\partial x}(x_{e,i}), \quad 1 \leq i \leq N+1.$$

Degree N approximate solution u_h Degree $N + 1$ Continuous Flux Approximation f_h 

Flux Reconstruction (FR) by Huynh [9]

6/75



$$u^{n+1} = u^n - \Delta t F_x^n,$$

where $F = f(u) + \frac{\Delta t}{2}(f(u))_t + \frac{\Delta t^2}{3!}f(u)_{tt} + \dots + \frac{\Delta t^N}{(N+1)!}\frac{\partial^N}{\partial t^N}f(u) \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u) dt$

$$u^{n+1} = u^n - \Delta t F_x^n,$$

where $F = f(u) + \frac{\Delta t}{2}(f(u))_t + \frac{\Delta t^2}{3!}f(u)_{tt} + \dots + \frac{\Delta t^N}{(N+1)!}\frac{\partial^N}{\partial t^N}f(u) \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u) dt$

Taylor series method

$$f(u)_t = f'(u) u_t, \quad u_t = -f(u)_x.$$

$$u^{n+1} = u^n - \Delta t F_x^n,$$

where $F = f(u) + \frac{\Delta t}{2}(f(u))_t + \frac{\Delta t^2}{3!}f(u)_{tt} + \dots + \frac{\Delta t^N}{(N+1)!}\frac{\partial^N}{\partial t^N}f(u) \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u) dt$

Taylor series method

$$f(u)_t = f'(u) u_t, \quad u_t = -f(u)_x.$$

Approximate Lax-Wendroff procedure (Zorio Et Al. [21])

$$\begin{aligned} f(u)_t &\approx \frac{f(u(x, t + \Delta t)) - f(u(x, t - \Delta t))}{2\Delta t} + O(\Delta t^2) \\ &\approx \frac{f(u + \Delta t \textcolor{blue}{u}_t) - f(u - \Delta t \textcolor{blue}{u}_t)}{2\Delta t} + O(\Delta t^2), \end{aligned}$$

and $\textcolor{blue}{u}_t = -f(u)_x$.

$$u^{n+1} = u^n - \Delta t F_x^n,$$

where $F = f(u) + \frac{\Delta t}{2}(f(u))_t + \frac{\Delta t^2}{3!}f(u)_{tt} + \dots + \frac{\Delta t^N}{(N+1)!}\frac{\partial^N}{\partial t^N}f(u) \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} f(u) dt$

Taylor series method

$$f(u)_t = f'(u) u_t, \quad u_t = -f(u)_x.$$

Approximate Lax-Wendroff procedure (Zorio Et Al. [21])

$$\begin{aligned} f(u)_t &\approx \frac{f(u(x, t + \Delta t)) - f(u(x, t - \Delta t))}{2\Delta t} + O(\Delta t^2) \\ &\approx \frac{f(u + \Delta t \textcolor{blue}{u}_t) - f(u - \Delta t \textcolor{blue}{u}_t)}{2\Delta t} + O(\Delta t^2), \end{aligned}$$

and $\textcolor{blue}{u}_t = -f(u)_x$.

F_h^δ corrected using $F_{e+\frac{1}{2}}$.

Past works : **Dissipation 1** (D1)

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{u}_{e+\frac{1}{2}}^+ - \textcolor{red}{u}_{e+\frac{1}{2}}^-].$$

Past works : **Dissipation 1** (D1)

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{u}_{e+\frac{1}{2}}^+ - \textcolor{red}{u}_{e+\frac{1}{2}}^-].$$

Dissipation 2 (D2) flux

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{U}_{e+\frac{1}{2}}^+ - \textcolor{red}{U}_{e+\frac{1}{2}}^-].$$

where

$$U = u + \frac{\Delta t}{2} u_t + \frac{\Delta t^2}{3!} u_{tt} + \cdots + \frac{\Delta t^N}{(N+1)!} \frac{\partial^N u}{\partial t^N} \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} u_h dt.$$

Past works : **Dissipation 1** (D1)

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{u}_{e+\frac{1}{2}}^+ - \textcolor{red}{u}_{e+\frac{1}{2}}^-].$$

Dissipation 2 (D2) flux

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{U}_{e+\frac{1}{2}}^+ - \textcolor{red}{U}_{e+\frac{1}{2}}^-].$$

where

$$U = u + \frac{\Delta t}{2} u_t + \frac{\Delta t^2}{3!} u_{tt} + \cdots + \frac{\Delta t^N}{(N+1)!} \frac{\partial^N u}{\partial t^N} \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} u_h dt.$$

Upwind flux for $u_t + a u_x = 0$

$$F_{e+\frac{1}{2}} = \begin{cases} F_{e+\frac{1}{2}}^- & a > 0, \\ F_{e+\frac{1}{2}}^- & a \leq 0. \end{cases}$$

Past works : **Dissipation 1** (D1)

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{u}_{e+\frac{1}{2}}^+ - \textcolor{red}{u}_{e+\frac{1}{2}}^-].$$

Dissipation 2 (D2) flux

$$F_{e+\frac{1}{2}} = \frac{1}{2} [F_{e+\frac{1}{2}}^- + F_{e+\frac{1}{2}}^+] - \frac{1}{2} \lambda_{e+\frac{1}{2}} [\textcolor{red}{U}_{e+\frac{1}{2}}^+ - \textcolor{red}{U}_{e+\frac{1}{2}}^-].$$


where

$$U = u + \frac{\Delta t}{2} u_t + \frac{\Delta t^2}{3!} u_{tt} + \cdots + \frac{\Delta t^N}{(N+1)!} \frac{\partial^N u}{\partial t^N} \approx \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} u_h dt.$$

Upwind flux for $u_t + a u_x = 0$


$$F_{e+\frac{1}{2}} = \begin{cases} F_{e+\frac{1}{2}}^- & a > 0, \\ F_{e+\frac{1}{2}}^- & a \leq 0. \end{cases}$$

Equivalent to ADER-DG for linear problems.




$$F_{e+\frac{1}{2}}^{\pm} = F_h^{\delta}(x_{e+\frac{1}{2}}^{\pm})$$

Sub-optimal accuracy for some non-linear problems,
robustness issues



Lax-Wendroff time averaging performed at faces,
only $\partial_t^{(k)} \mathbf{u}$ terms extrapolated



LWFR-D2 with EA scheme




RKFR

Performance gains measured


11/75

GL, Radau



Performance gain of D2 over D1

GL, Radau



Performance gain of LW over RK

For more results comparing LW-D2/LW-D1 and LW/RK, see [2].


Godunov's order barrier theorem

Linear schemes which do not add oscillations can be at most first order accurate.

Godunov's order barrier theorem

Linear schemes which do not add oscillations can be at most first order accurate.


TVD/TVB Limiter



Godunov's order barrier theorem

Linear schemes which do not add oscillations can be at most first order accurate.

TVD/TVB Limiter



$$\alpha_e = \alpha(\mathbf{u}_e)$$

Blending limiter
(Hennemann Et Al. [8])

Solution points and subcells



High order LWFR update

$$\mathbf{u}_e^{H,n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e^H.$$

High order LWFR update

$$\mathbf{u}_e^{H,n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e^H.$$

Lower order subcell update (FO FVM or MUSCL-Hancock)

$$\mathbf{u}_e^{L,n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e^L.$$

High order LWFR update

$$\mathbf{u}_e^{H,n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e^H.$$

Lower order subcell update (FO FVM or MUSCL-Hancock)

$$\mathbf{u}_e^{L,n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e^L.$$

Blend residual with $\alpha_e \in [0, 1]$

$$\mathbf{R}_e = (1 - \alpha_e) \mathbf{R}_e^H + \alpha_e \mathbf{R}_e^L,$$

Limited update

$$\mathbf{u}_e^{n+1} = \mathbf{u}_e^n - \frac{\Delta t}{\Delta x_e} \mathbf{R}_e.$$

Choice of α_e : Smoothness indicator [8]

Legendre expansion of degree N polynomial $\epsilon = \epsilon(\xi)$

$$\epsilon = \sum_{j=1}^{N+1} m_j L_j, \quad m_j = \langle \epsilon, L_j \rangle_{L^2},$$

Energy content (Persson and Peraire [12])

$$\mathbb{E} := \max \left(\frac{m_{N+1}^2}{\beta_1 m_1^2 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1^2 + \sum_{j=2}^N m_j^2} \right), \quad 0 \leq \beta_i \leq 1.$$

Choice of α_e : Smoothness indicator [8]

Legendre expansion of degree N polynomial $\epsilon = \epsilon(\xi)$

$$\epsilon = \sum_{j=1}^{N+1} m_j L_j, \quad m_j = \langle \epsilon, L_j \rangle_{L^2},$$

Energy content (Persson and Peraire [12])

$$\mathbb{E} := \max \left(\frac{m_{N+1}^2}{\beta_1 m_1^2 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1^2 + \sum_{j=2}^N m_j^2} \right), \quad 0 \leq \beta_i \leq 1.$$

$$\epsilon = \rho p$$

Choice of α_e : Smoothness indicator [8]


Legendre expansion of degree N polynomial $\epsilon = \epsilon(\xi)$

$$\epsilon = \sum_{j=1}^{N+1} m_j L_j, \quad m_j = \langle \epsilon, L_j \rangle_{L^2},$$

Energy content (Persson and Peraire [12])

$$\mathbb{E} := \max \left(\frac{m_{N+1}^2}{\beta_1 m_1^2 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1^2 + \sum_{j=2}^N m_j^2} \right), \quad 0 \leq \beta_i \leq 1.$$

$$\epsilon = \rho p$$



$$\alpha(\mathbb{E}) = \frac{1}{1 + \exp(-\frac{s}{\mathbb{T}}(\mathbb{E} - \mathbb{T}))}$$

$$\mathbb{T}(N) = 0.5 \cdot 10^{-1.8(N+1)^{1/4}}, \quad \alpha(\mathbb{E}=0) = 0.0001$$

$$\tilde{\alpha} = \begin{cases} 0, & \text{if } \alpha < \alpha_{\min} \\ \alpha, & \text{if } \alpha_{\min} \leq \alpha \leq 1 - \alpha_{\min} \\ 1, & \text{if } 1 - \alpha_{\min} < \alpha \end{cases}$$

$$\alpha^{\text{final}} = \max_{e \in V_e} \{\alpha, 0.5 \alpha_e\}$$

Solution points and subcells



Subcell $[x_{j-\frac{1}{2}}^e, x_{j+\frac{1}{2}}^e]$

$$x_{j+\frac{1}{2}}^e - x_{j-\frac{1}{2}}^e = \Delta x_e w_j, \quad 1 \leq j \leq N+1,$$

where $\{w_j\}_{j:1}^{N+1}$ are the Gauss-Legendre quadrature weights.

FVM



$$\bar{u}_e^{n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} (F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}}).$$

Solution points and subcells



Subcell $[x_{j-\frac{1}{2}}^e, x_{j+\frac{1}{2}}^e]$

$$x_{j+\frac{1}{2}}^e - x_{j-\frac{1}{2}}^e = \Delta x_e w_j, \quad 1 \leq j \leq N+1,$$

where $\{w_j\}_{j:1}^{N+1}$ are the Gauss-Legendre quadrature weights.

FVM



$$\bar{u}_e^{n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} (F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}}).$$

Solution points and subcells




Subcell $[x_{j-\frac{1}{2}}^e, x_{j+\frac{1}{2}}^e]$

$$x_{j+\frac{1}{2}}^e - x_{j-\frac{1}{2}}^e = \Delta x_e w_j, \quad 1 \leq j \leq N + 1,$$

where $\{w_j\}_{j:1}^{N+1}$ are the Gauss-Legendre quadrature weights.

FVM



$$\bar{u}_e^{n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} (F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}}).$$

Solution points and subcells




Subcell $[x_{j-\frac{1}{2}}^e, x_{j+\frac{1}{2}}^e]$

$$x_{j+\frac{1}{2}}^e - x_{j-\frac{1}{2}}^e = \Delta x_e w_j, \quad 1 \leq j \leq N + 1,$$

where $\{w_j\}_{j:1}^{N+1}$ are the Gauss-Legendre quadrature weights.

FVM



$$\bar{u}_e^{n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} (F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}}).$$

Interface numerical flux

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Interface numerical flux

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Interface numerical flux

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Interface numerical flux

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Initial candidate

$$\tilde{\mathbf{F}}_{e+\frac{1}{2}} = \left(1 - \alpha_{e+\frac{1}{2}}\right) \mathbf{F}_{e+\frac{1}{2}}^{\text{LW}} + \alpha_{e+\frac{1}{2}} \mathbf{f}_{e,N+3/2}, \quad \alpha_{e+\frac{1}{2}} = \frac{1}{2}(\alpha_e + \alpha_{e+1}).$$

Lower order update of last solution point of cell e

$$\tilde{\mathbf{u}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\tilde{\mathbf{F}}_{e+\frac{1}{2}} - \mathbf{f}_{e,N+1/2}).$$

Assume **concave** p such that

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}.$$

For purely low order

$$\tilde{\mathbf{u}}_{\text{low}}^{n+1} = \mathbf{u}_{e,N+1}^n - \frac{\Delta t}{\Delta x_e w_{N+1}} (\mathbf{f}_{e,N+3/2} - \mathbf{f}_{e,N+1/2}) \in \mathcal{U}_{\text{ad}}.$$

Thus, for

$$\theta = \min \left(\left| \frac{\epsilon - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})}{p(\tilde{\mathbf{u}}^{n+1}) - p(\tilde{\mathbf{u}}_{\text{low}}^{n+1})} \right|, 1 \right),$$

we will have

$$p(\theta \tilde{\mathbf{u}}^{n+1} + (1 - \theta) \mathbf{u}_{\text{low}}^{n+1}) \geq \theta p(\tilde{\mathbf{u}}^{n+1}) + (1 - \theta) p(\mathbf{u}_{\text{low}}^{n+1}) > \epsilon.$$

Final choice: $\mathbf{F}_{e+\frac{1}{2}} = \theta \tilde{\mathbf{F}}_{e+\frac{1}{2}} + (1 - \theta) \mathbf{f}_{e,N+3/2}$.

Low order residual

$$\begin{aligned}(\tilde{\mathbf{u}}_1^e)^{n+1} &= (\mathbf{u}_1^e)^n - \frac{\Delta t}{w_1 \Delta x_e} \left[f_{\frac{3}{2}} - F_{e-\frac{1}{2}} \right], \\(\tilde{\mathbf{u}}_j^e)^{n+1} &= (\mathbf{u}_j^e)^n - \frac{\Delta t}{w_j \Delta x_e} \left[f_{j+\frac{1}{2}} - f_{j-\frac{1}{2}} \right], \quad 2 \leq j \leq N, \\(\tilde{\mathbf{u}}_N^e)^{n+1} &= (\mathbf{u}_N^e)^n - \frac{\Delta t}{w_{N+1} \Delta x_e} \left[F_{e+\frac{1}{2}} - f_{N-\frac{1}{2}} \right].\end{aligned}$$

By appropriate choice of $F_{e \pm \frac{1}{2}}$,

$$(\tilde{\mathbf{u}}_j^e)^{n+1} \in \mathcal{U}_{\text{ad}}, \quad 1 \leq j \leq N+1.$$

$$\Rightarrow \bar{\mathbf{u}}_e^{n+1} = \bar{\mathbf{u}}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right) = \sum_{j=1}^{N+1} w_j \tilde{\mathbf{u}}_j^{n+1} \in \mathcal{U}_{\text{ad}}.$$

Zhang-Shu's scaling limiter applies.

Low order residual

$$\begin{aligned}(\tilde{\mathbf{u}}_1^e)^{n+1} &= (\mathbf{u}_1^e)^n - \frac{\Delta t}{w_1 \Delta x_e} \left[f_{\frac{3}{2}} - F_{e-\frac{1}{2}} \right], \\(\tilde{\mathbf{u}}_j^e)^{n+1} &= (\mathbf{u}_j^e)^n - \frac{\Delta t}{w_j \Delta x_e} \left[f_{j+\frac{1}{2}} - f_{j-\frac{1}{2}} \right], \quad 2 \leq j \leq N, \\(\tilde{\mathbf{u}}_N^e)^{n+1} &= (\mathbf{u}_N^e)^n - \frac{\Delta t}{w_{N+1} \Delta x_e} \left[F_{e+\frac{1}{2}} - f_{N-\frac{1}{2}} \right].\end{aligned}$$

By appropriate choice of $F_{e \pm \frac{1}{2}}$,

$$(\tilde{\mathbf{u}}_j^e)^{n+1} \in \mathcal{U}_{\text{ad}}, \quad 1 \leq j \leq N+1.$$

$$\Rightarrow \bar{\mathbf{u}}_e^{n+1} = \bar{\mathbf{u}}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right) = \sum_{j=1}^{N+1} w_j \tilde{\mathbf{u}}_j^{n+1} \in \mathcal{U}_{\text{ad}}.$$

Zhang-Shu's scaling limiter applies.

Low order residual

$$\begin{aligned}(\tilde{\mathbf{u}}_1^e)^{n+1} &= (\mathbf{u}_1^e)^n - \frac{\Delta t}{w_1 \Delta x_e} \left[f_{\frac{3}{2}} - F_{e-\frac{1}{2}} \right], \\(\tilde{\mathbf{u}}_j^e)^{n+1} &= (\mathbf{u}_j^e)^n - \frac{\Delta t}{w_j \Delta x_e} \left[f_{j+\frac{1}{2}} - f_{j-\frac{1}{2}} \right], \quad 2 \leq j \leq N, \\(\tilde{\mathbf{u}}_N^e)^{n+1} &= (\mathbf{u}_N^e)^n - \frac{\Delta t}{w_{N+1} \Delta x_e} \left[F_{e+\frac{1}{2}} - f_{N-\frac{1}{2}} \right].\end{aligned}$$

By appropriate choice of $F_{e \pm \frac{1}{2}}$,

$$(\tilde{\mathbf{u}}_j^e)^{n+1} \in \mathcal{U}_{\text{ad}}, \quad 1 \leq j \leq N+1.$$

$$\Rightarrow \bar{\mathbf{u}}_e^{n+1} = \bar{\mathbf{u}}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right) = \sum_{j=1}^{N+1} w_j \tilde{\mathbf{u}}_j^{n+1} \in \mathcal{U}_{\text{ad}}.$$

Zhang-Shu's scaling limiter applies.

Low order residual

$$\begin{aligned}(\tilde{\mathbf{u}}_1^e)^{n+1} &= (\mathbf{u}_1^e)^n - \frac{\Delta t}{w_1 \Delta x_e} \left[f_{\frac{3}{2}} - F_{e-\frac{1}{2}} \right], \\(\tilde{\mathbf{u}}_j^e)^{n+1} &= (\mathbf{u}_j^e)^n - \frac{\Delta t}{w_j \Delta x_e} \left[f_{j+\frac{1}{2}} - f_{j-\frac{1}{2}} \right], \quad 2 \leq j \leq N, \\(\tilde{\mathbf{u}}_N^e)^{n+1} &= (\mathbf{u}_N^e)^n - \frac{\Delta t}{w_{N+1} \Delta x_e} \left[F_{e+\frac{1}{2}} - f_{N-\frac{1}{2}} \right].\end{aligned}$$


By appropriate choice of $F_{e \pm \frac{1}{2}}$,

$$(\tilde{\mathbf{u}}_j^e)^{n+1} \in \mathcal{U}_{\text{ad}}, \quad 1 \leq j \leq N+1.$$


$$\Rightarrow \bar{\mathbf{u}}_e^{n+1} = \bar{\mathbf{u}}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right) = \sum_{j=1}^{N+1} w_j \tilde{\mathbf{u}}_j^{n+1} \in \mathcal{U}_{\text{ad}}.$$

Zhang-Shu's scaling limiter applies.

$$\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$$




$$\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$$



$$\begin{aligned}\mathbf{u}_{j-\frac{1}{2}+} &= \mathbf{u}_j(x_{j-\frac{1}{2}}), & \mathbf{u}_{j+\frac{1}{2}-} &= \mathbf{u}_j(x_{j+\frac{1}{2}}) \\ \mathbf{u}_j(x) &= \mathbf{u}_j^n + \boldsymbol{\sigma}_j (x - x_j)\end{aligned}$$


$$\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$$



$$\begin{aligned}\mathbf{u}_{j-\frac{1}{2}+} &= \mathbf{u}_j(x_{j-\frac{1}{2}}), & \mathbf{u}_{j+\frac{1}{2}-} &= \mathbf{u}_j(x_{j+\frac{1}{2}}) \\ \mathbf{u}_j(x) &= \mathbf{u}_j^n + \boldsymbol{\sigma}_j (x - x_j)\end{aligned}$$

$$\begin{aligned}\boldsymbol{\sigma}_j &= \text{minmod}\left(\beta_e \frac{\mathbf{u}_{j+1} - \mathbf{u}_j}{x_{j+1} - x_j}, D_{\text{cent}}(\mathbf{u})_j, \beta_e \frac{\mathbf{u}_j^n - \mathbf{u}_{j-1}^n}{x_j - x_{j-1}}\right) \\ \beta_e &= 2 - \alpha_e\end{aligned}$$

Admissibility of low order method



Theorem. (Extension of Berthon [3]) Consider the hyperbolic conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ which preserves the convex set Ω . Let $\{\mathbf{u}_i^n\}_{i \in \mathbb{Z}}$ be the approximate solution at time level n and assume that $\mathbf{u}_i^n \in \Omega$ for all $i \in \mathbb{Z}$. Consider **conservative reconstructions**

$$\mathbf{u}_i^{n,+} = \mathbf{u}_i + (x_{i+1/2} - x_i) \boldsymbol{\sigma}_i, \quad \mathbf{u}_i^{n,-} = \mathbf{u}_i + (x_{i-1/2} - x_i) \boldsymbol{\sigma}_i.$$

Define $\mathbf{u}_i^{*,\pm}$ satisfying

$$\mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \mu^+ \mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm},$$

where

$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}}.$$

Assume that the slope $\boldsymbol{\sigma}_i$ is chosen so that

$$\mathbf{u}_i^{*,\pm} \in \mathcal{U}_{\text{ad}}.$$

Then, under **appropriate** time step restrictions, the updated solution \mathbf{u}_i^{n+1} , defined by the MUSCL-Hancock procedure is in \mathcal{U}_{ad} .

Berthon defined $\mathbf{u}_i^{*,\pm}$

$$\frac{1}{2}\mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \frac{1}{2}\mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm}.$$

Generalizing Berthon's proof

Berthon defined $\mathbf{u}_i^{*,\pm}$

$$\frac{1}{2}\mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \frac{1}{2}\mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm}.$$

Our generalization

$$\mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \mu^+ \mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm},$$

where

$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}}.$$

Generalizing Berthon's proof

Berthon defined $\mathbf{u}_i^{*,\pm}$

$$\frac{1}{2}\mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \frac{1}{2}\mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm}.$$

Our generalization

$$\mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,\pm} + \mu^+ \mathbf{u}_i^{n,+} = 2\mathbf{u}_i^{n,\pm},$$

where

$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}}.$$


For **conservative reconstruction**

$$\mathbf{u}_i^{*,\pm} = \mathbf{u}_i^n + 2(x_{i\pm 1/2} - x_i)\boldsymbol{\sigma}_i,$$


noting that

$$\mathbf{u}_i^{n,+} = \mathbf{u}_i + (x_{i+1/2} - x_i)\boldsymbol{\sigma}_i, \quad \mathbf{u}_i^{n,-} = \mathbf{u}_i + (x_{i-1/2} - x_i)\boldsymbol{\sigma}_i.$$

Idea of proof




Idea of proof



$$\frac{1}{\Delta x_i} \int_{x_{i-1/2}}^{x_{i+1/2}} \mathbf{u}^h(x, \Delta t / 2) dx = \mathbf{u}_i^{n+\frac{1}{2}, +}$$


Idea of proof



$$\frac{1}{\Delta x_i} \int_{x_{i-1/2}}^{x_{i+1/2}} \mathbf{u}^h(x, \Delta t / 2) dx = \mathbf{u}_i^{n+\frac{1}{2}, +}$$


$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n$$

Idea of proof



$$\frac{1}{\Delta x_i} \int_{x_{i-1/2}}^{x_{i+1/2}} \mathbf{u}^h(x, \Delta t / 2) dx = \mathbf{u}_i^{n+\frac{1}{2}, +}$$

$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n$$



Enforcing slope restriction

Given candidate slope σ_i ,

$$\mathbf{u}_i^{*,\pm} := \mathbf{u}_i^n + 2(x_{i\pm 1/2} - x_i) \sigma_i.$$



Find $\theta \in [0, 1]$

$$\mathbf{u}_i^n + 2(x_{i\pm 1/2} - x_i) \theta \sigma_i \in \mathcal{U}_{\text{ad}}$$

by Zhang-Shu type procedure.

Unstructured, curvilinear grids


23/75



$$(\xi, \eta) \mapsto \mathbf{x}^e(\xi, \eta)$$

\mathbf{x}^e is a **degree k** polynomial in ξ, η

$$\begin{aligned}\mathbf{x} &= (x_1, x_2, x_3) = (x, y, z) \\ \boldsymbol{\xi} &= (\xi_1, \xi_2, \xi_3) = (\xi, \eta, \zeta)\end{aligned}$$



$$\begin{aligned}\mathbf{a}_i &= \frac{\partial}{\partial \xi_i} \mathbf{x} \\ \mathbf{a}^i &= \nabla_{\mathbf{x}} \xi^i = J^{-1} (\mathbf{a}_j \times \mathbf{a}_k) \\ J &= \mathbf{a}_1 \cdot (\mathbf{a}_2 \times \mathbf{a}_3)\end{aligned}$$

Covariant and contravariant coordinate vectors in relation to the coordinate lines

$$\mathbf{u}_t + \nabla_{\mathbf{x}} \cdot \mathbf{f} = 0 \longrightarrow \tilde{\mathbf{u}}_t + \nabla_{\boldsymbol{\xi}} \cdot \tilde{\mathbf{f}} = 0,$$

where

$$\begin{aligned}\tilde{\mathbf{u}} &= J \vec{Q}, \\ \tilde{\mathbf{f}}^i &= J \mathbf{a}^i \cdot \mathbf{f} = \sum_{n=1}^3 J a_n^i f_n.\end{aligned}$$

$$\begin{aligned} & \frac{d\boldsymbol{u}_{e,\boldsymbol{p}}^\delta}{dt} + \frac{1}{J} \nabla_{\boldsymbol{\xi}}^N \cdot \tilde{\boldsymbol{f}}_e(\boldsymbol{\xi}_{\boldsymbol{p}}) \\ & + \frac{1}{J} \sum_{i=1}^3 ((\tilde{\boldsymbol{f}}_e \cdot \boldsymbol{n}_{s,i})^* - \tilde{\boldsymbol{f}}_e \cdot \boldsymbol{n}_{s,i})(\boldsymbol{\xi}_i^R) g'_R(\xi_{p_i}) \\ & + \frac{1}{J} \sum_{i=1}^3 ((\tilde{\boldsymbol{f}}_e \cdot \boldsymbol{n}_{s,i})^* - \tilde{\boldsymbol{f}}_e \cdot \boldsymbol{n}_{s,i})(\boldsymbol{\xi}_i^L) g'_L(\xi_{p_i}) = \mathbf{0}, \end{aligned}$$

$$\begin{aligned} \mathbf{u}_t + \frac{1}{J} \nabla_{\boldsymbol{\xi}} \cdot \tilde{\mathbf{f}} &= 0, \\ \tilde{\mathbf{f}}^i &= J \mathbf{a}^i \cdot \mathbf{f} = \sum_{n=1}^3 J a_n^i \mathbf{f}_n. \end{aligned}$$

$$\begin{aligned} \mathbf{u}_t + \frac{1}{J} \nabla_{\boldsymbol{\xi}} \cdot \tilde{\mathbf{f}} &= 0, \\ \tilde{\mathbf{f}}^i &= J \mathbf{a}^i \cdot \mathbf{f} = \sum_{n=1}^3 J a_n^i \mathbf{f}_n. \end{aligned}$$

$$\begin{aligned} \mathbf{u}^{n+1}(\boldsymbol{\xi}) &= \mathbf{u}^n(\boldsymbol{\xi}) - \frac{1}{J} \Delta t \nabla_{\boldsymbol{\xi}} \cdot \tilde{\mathbf{F}} \\ \tilde{\mathbf{F}} &= \sum_{k=0}^N \frac{\Delta t^k}{(k+1)!} \partial_t^k \tilde{\mathbf{f}} \end{aligned}$$

$$\begin{aligned} \boldsymbol{u}_t + \frac{1}{J} \nabla_{\boldsymbol{\xi}} \cdot \tilde{\boldsymbol{f}} &= 0, \\ \tilde{\boldsymbol{f}}^i &= J \boldsymbol{a}^i \cdot \boldsymbol{f} = \sum_{n=1}^3 J a_n^i \boldsymbol{f}_n. \end{aligned}$$

$$\begin{aligned} \boldsymbol{u}^{n+1}(\boldsymbol{\xi}) &= \boldsymbol{u}^n(\boldsymbol{\xi}) - \frac{1}{J} \Delta t \nabla_{\boldsymbol{\xi}} \cdot \tilde{\boldsymbol{F}} \\ \tilde{\boldsymbol{F}} &= \sum_{k=0}^N \frac{\Delta t^k}{(k+1)!} \partial_t^k \tilde{\boldsymbol{f}} \end{aligned}$$

$$\begin{aligned} \boldsymbol{u}_{e,\boldsymbol{p}}^{n+1} - \boldsymbol{u}_{e,\boldsymbol{p}}^n &+ \frac{1}{J} \Delta t \nabla_{\boldsymbol{\xi}} \cdot \tilde{\boldsymbol{F}}_e^\delta(\boldsymbol{\xi}_{\boldsymbol{p}}) \\ &+ \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial\Omega_{s,i}} ((\tilde{\boldsymbol{F}}_e^\delta \cdot \boldsymbol{n}_{s,i})^* - \tilde{\boldsymbol{F}}_e^\delta \cdot \boldsymbol{n}_{s,i})(\boldsymbol{\xi}_i^R) g'_R(\xi_{p_i}) dS_{\boldsymbol{\xi}} \\ &+ \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial\Omega_{s,i}} ((\tilde{\boldsymbol{F}}_e^\delta \cdot \boldsymbol{n}_{s,i})^* - \tilde{\boldsymbol{F}}_e^\delta \cdot \boldsymbol{n}_{s,i})(\boldsymbol{\xi}_i^L) g'_L(\xi_{p_i}) dS_{\boldsymbol{\xi}} = \mathbf{0}. \end{aligned}$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi^i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi^i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Assume a free stream

$$\mathbf{u}^n = \underline{\mathbf{c}} \text{ and } \mathbf{f}(\underline{\mathbf{c}}) = \mathbf{c}$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi^i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Assume a free stream

$$\mathbf{u}^n = \underline{\mathbf{c}} \text{ and } \mathbf{f}(\underline{\mathbf{c}}) = \mathbf{c}$$

$$\begin{aligned} \mathbf{u}_t &= -\frac{1}{J} \nabla_{\xi}^N \cdot \tilde{\mathbf{f}}_e^\delta \\ &= -\sum_{n=1}^3 \left(\sum_{i=1}^3 \partial_{\xi^i}^N (J a_n^i) \right) \mathbf{c}_n = 0. \end{aligned}$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi_i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Assume a free stream

$$\mathbf{u}^n = \underline{\mathbf{c}} \text{ and } \mathbf{f}(\underline{\mathbf{c}}) = \mathbf{c}$$

$$\begin{aligned} \mathbf{u}_t &= -\frac{1}{J} \nabla_{\xi}^N \cdot \tilde{\mathbf{f}}_e^\delta \\ &= -\sum_{n=1}^3 \left(\sum_{i=1}^3 \partial_{\xi_i}^N (J a_n^i) \right) \mathbf{c}_n = 0. \end{aligned}$$

$$N=1: \quad \partial_t \tilde{\mathbf{f}}^\delta = \frac{\tilde{\mathbf{f}}(\mathbf{u} + \Delta t \mathbf{u}_t) - \tilde{\mathbf{f}}(\mathbf{u} - \Delta t \mathbf{u}_t)}{2 \Delta t} = \mathbf{0}$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi_i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Assume a free stream

$$\mathbf{u}^n = \underline{\mathbf{c}} \text{ and } \mathbf{f}(\underline{\mathbf{c}}) = \mathbf{c}$$

$$\begin{aligned} \mathbf{u}_t &= -\frac{1}{J} \nabla_{\xi}^N \cdot \tilde{\mathbf{f}}_e^\delta \\ &= -\sum_{n=1}^3 \left(\sum_{i=1}^3 \partial_{\xi_i}^N (J a_n^i) \right) \mathbf{c}_n = 0. \end{aligned}$$

$$N=1: \quad \partial_t \tilde{\mathbf{f}}^\delta = \frac{\tilde{\mathbf{f}}(\mathbf{u} + \Delta t \mathbf{u}_t) - \tilde{\mathbf{f}}(\mathbf{u} - \Delta t \mathbf{u}_t)}{2 \Delta t} = \mathbf{0}$$

$$\Rightarrow \tilde{\mathbf{F}}^\delta = \tilde{\mathbf{f}}^\delta + \frac{\Delta t}{2} \partial_t \tilde{\mathbf{f}}^\delta = \tilde{\mathbf{f}}^\delta,$$

Claimed free stream preservation conditions

$$\sum_{i=1}^3 \partial_{\xi_i}^N (J \mathbf{a}^i) = \mathbf{0}$$

Assume a free stream

$$\mathbf{u}^n = \underline{\mathbf{c}} \text{ and } \mathbf{f}(\underline{\mathbf{c}}) = \mathbf{c}$$

$$\begin{aligned} \mathbf{u}_t &= -\frac{1}{J} \nabla_{\xi}^N \cdot \tilde{\mathbf{f}}_e^\delta \\ &= -\sum_{n=1}^3 \left(\sum_{i=1}^3 \partial_{\xi_i}^N (J a_n^i) \right) \mathbf{c}_n = 0. \end{aligned}$$

$$N=1: \quad \partial_t \tilde{\mathbf{f}}^\delta = \frac{\tilde{\mathbf{f}}(\mathbf{u} + \Delta t \mathbf{u}_t) - \tilde{\mathbf{f}}(\mathbf{u} - \Delta t \mathbf{u}_t)}{2 \Delta t} = \mathbf{0}$$

$$\Rightarrow \tilde{\mathbf{F}}^\delta = \tilde{\mathbf{f}}^\delta + \frac{\Delta t}{2} \partial_t \tilde{\mathbf{f}}^\delta = \tilde{\mathbf{f}}^\delta,$$

where

$$\tilde{\mathbf{f}}_i^\delta = J \mathbf{a}^i \cdot \mathbf{c} = J \mathbf{a}^i \cdot \mathbf{c}.$$

$$\begin{aligned} \mathbf{u}^{n+1} - \mathbf{u}^n + \frac{1}{J} \Delta t \left(\sum_{i=1}^3 \partial_{\xi^i}^N (J \mathbf{a}^i) \right) \cdot \mathbf{c} \\ + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((J \mathbf{a}^i \cdot \mathbf{c})^* - J \mathbf{a}^i \cdot \mathbf{c})(\xi_i^R) g'_R(\xi_{p_i}) dS_{\xi} \\ + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((J \mathbf{a}^i \cdot \mathbf{c})^* - J \mathbf{a}^i \cdot \mathbf{c})(\xi_i^L) g'_L(\xi_{p_i}) dS_{\xi} = \mathbf{0}. \end{aligned}$$

$$\begin{aligned}
& \boldsymbol{u}^{n+1} - \boldsymbol{u}^n + \frac{1}{J} \Delta t \left(\sum_{i=1}^3 \partial_{\xi^i}^N (\boldsymbol{J} \boldsymbol{a}^i) \right) \cdot \boldsymbol{c} \\
& + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((\boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})^* - \boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})(\xi_i^R) g'_R(\xi_{p_i}) dS_{\xi} \\
& + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((\boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})^* - \boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})(\xi_i^L) g'_L(\xi_{p_i}) dS_{\xi} = \mathbf{0}.
\end{aligned}$$

Metric identities

$$\sum_{i=1}^3 \partial_{\xi^i}^N (\boldsymbol{J} \boldsymbol{a}^i) = \mathbf{0}$$

$$\begin{aligned}
& \boldsymbol{u}^{n+1} - \boldsymbol{u}^n + \frac{1}{J} \Delta t \left(\sum_{i=1}^3 \partial_{\xi^i}^N (\boldsymbol{J} \boldsymbol{a}^i) \right) \cdot \boldsymbol{c} \\
& + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((\boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})^* - \boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})(\xi_i^R) g'_R(\xi_{p_i}) dS_{\xi} \\
& + \frac{1}{J} \Delta t \sum_{i=1}^3 \int_{\partial \Omega_{s,i}} ((\boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})^* - \boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})(\xi_i^L) g'_L(\xi_{p_i}) dS_{\xi} = \mathbf{0}.
\end{aligned}$$

Metric identities

$$\sum_{i=1}^3 \partial_{\xi^i}^N (\boldsymbol{J} \boldsymbol{a}^i) = \mathbf{0}$$

Conformality

$$(\boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c})^* - \boldsymbol{J} \boldsymbol{a}^i \cdot \boldsymbol{c} = \mathbf{0}$$

$$J\mathbf{a}^1 = (y_\eta, -x_\eta), \quad J\mathbf{a}^2 = (-y_\xi, x_\xi).$$

$$\partial_\xi^N J\mathbf{a}^1 + \partial_\eta^N J\mathbf{a}^2 = \mathbf{0}$$

2-D

$$\begin{aligned}\partial_\xi^N (y_\eta) - \partial_\eta^N (x_\xi) &= 0 \\ -\partial_\xi^N (x_\eta) + \partial_\eta^N (x_\xi) &= 0\end{aligned}$$

Condition: Degree of mesh $\leq N$

$$J\mathbf{a}^1 = (y_\eta, -x_\eta), \quad J\mathbf{a}^2 = (-y_\xi, x_\xi).$$

$$\partial_\xi^N J\mathbf{a}^1 + \partial_\eta^N J\mathbf{a}^2 = \mathbf{0}$$

2-D

$$\begin{aligned}\partial_\xi^N (y_\eta) - \partial_\eta^N (x_\xi) &= 0 \\ -\partial_\xi^N (x_\eta) + \partial_\eta^N (x_\xi) &= 0\end{aligned}$$

Condition: Degree of mesh $\leq N$

$$J\mathbf{a}^1 = \mathbf{a}^2 \times \mathbf{a}^3, \quad J\mathbf{a}^2 = \mathbf{a}^3 \times \mathbf{a}^1, \quad J\mathbf{a}^3 = \mathbf{a}^1 \times \mathbf{a}^2$$

3-DCondition: Degree of mesh $\leq N/2$.

$$J\mathbf{a}^1 = (y_\eta, -x_\eta), \quad J\mathbf{a}^2 = (-y_\xi, x_\xi).$$

$$\partial_\xi^N J\mathbf{a}^1 + \partial_\eta^N J\mathbf{a}^2 = \mathbf{0}$$

2-D

$$\begin{aligned}\partial_\xi^N (y_\eta) - \partial_\eta^N (x_\xi) &= 0 \\ -\partial_\xi^N (x_\eta) + \partial_\eta^N (x_\xi) &= 0\end{aligned}$$

Condition: Degree of mesh $\leq N$

$$J\mathbf{a}^1 = \mathbf{a}^2 \times \mathbf{a}^3, \quad J\mathbf{a}^2 = \mathbf{a}^3 \times \mathbf{a}^1, \quad J\mathbf{a}^3 = \mathbf{a}^1 \times \mathbf{a}^2$$

3-DCondition: Degree of mesh $\leq N/2$.

Conservative form of metric terms

$$Ja_n^i = -\mathbf{e}_i \cdot \nabla_\xi \times (x_m \nabla_\xi x_l), \quad i = 1, 2, 3, \quad n = 1, 2, 3, \quad (n, m, l) \text{ cyclic.}$$

$$J\mathbf{a}^1 = (y_\eta, -x_\eta), \quad J\mathbf{a}^2 = (-y_\xi, x_\xi).$$

$$\partial_\xi^N J\mathbf{a}^1 + \partial_\eta^N J\mathbf{a}^2 = \mathbf{0}$$

2-D

$$\begin{aligned}\partial_\xi^N (y_\eta) - \partial_\eta^N (x_\xi) &= 0 \\ -\partial_\xi^N (x_\eta) + \partial_\eta^N (x_\xi) &= 0\end{aligned}$$

Condition: Degree of mesh $\leq N$

$$J\mathbf{a}^1 = \mathbf{a}^2 \times \mathbf{a}^3, \quad J\mathbf{a}^2 = \mathbf{a}^3 \times \mathbf{a}^1, \quad J\mathbf{a}^3 = \mathbf{a}^1 \times \mathbf{a}^2$$

3-DCondition: Degree of mesh $\leq N/2$.

Conservative form of metric terms

$$Ja_n^i = -\mathbf{e}_i \cdot \nabla_\xi \times (x_m \nabla_\xi x_l), \quad i = 1, 2, 3, \quad n = 1, 2, 3, \quad (n, m, l) \text{ cyclic.}$$


Free stream always satisfied with **conservative curl form**

$$Ja_n^i = -\mathbf{e}_i \cdot \nabla_\xi^N \times (x_l \nabla_\xi x_m)$$

Mesh degree = 6



Solution polynomial degree 5




Solution polynomial degree 6

Numerical Results


2-D Composite signal [11]

Min = 0, Max = 1




Initial State

Min= -0.01572, Max = 1.01422




TVB - 100

Min = -0.0016, Max = 1.00684



MUSCL-Hancock




Solid wall boundary conditions on $[0, 1]$ with intial condition

$$(\rho, v, p) = \begin{cases} (1, 0, 1000), & \text{if } x < 0.1, \\ (1, 0, 0.01), & 0.1 < x < 0.9, \\ (1, 0, 100), & x > 0.9. \end{cases}$$

Double Mach reflection [19]

34/75




[Animation link](#)

Double Mach Reflection

35/75

$t = 0.2$, NC = 568×142 , Rusanov, Degree $N = 4$



Trixi.jl




LWFR


Double Mach Reflection

36/75

$t = 0.2$, NC= 568×142 , LWFR, Degree $N = 4$




HLLC, $\beta_1 = 0.1, \beta_2 = 1.0$



Rusanov. $\beta_1 = \beta_2 = 1.0$

Kelvin-Helmholtz Instability [18, 15]

37/75




[Animation link](#)

Kelvin-Helmholtz Instability [18, 15]

38/75



Density profile $t = 0.4$, NC = 256^2 , Degree $N = 4$



Trixi.jl

LWFR

$t = 0.001, \quad NC = 160^2, \quad \text{Degree } N = 4$




The initial condition is given by


$$\rho = 1.0, \quad v_1 = v_2 = 0.0, \quad E(x, y) = \begin{cases} \frac{0.244816}{\Delta x \Delta y} & x < \Delta x, y < \Delta y, \\ 10^{-12} & \text{otherwise.} \end{cases}$$

Animation link

Density profile, NC = 450×225 , $t = 0.001$, Degree $N = 4$



Trixi.jl



LWFR

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

- A Jacobian free Lax-Wendroff scheme presented in the collocation based Flux Reconstruction (FR) framework.
- Dissipative part of numerical flux computed with time averaged solution, leading to an increase in CFL number. The obtained scheme is equivalent to ADER-DG for linear problems.
- Central part of numerical flux computed by performing the Lax-Wendroff procedure at the face, leading to improvement in accuracy.
- Sub-cell based blending limiter [8] with MUSCL-Hancock reconstruction introduced for Lax-Wendroff schemes and is found to be more accurate than the first order blending initially proposed in [8].
- Problem independent slope limiting procedure proposed for MUSCL-Hancock schemes on general grids which leads to provable admissibility preservation.
- Using the admissibility preserving MUSCL-Hancock and a flux correction, an admissibility preserving Lax-Wendroff scheme was constructed.
- The scheme was extended to unstructured grids and the free stream condition was proven to be equivalent to that of Flux Reconstruction schemes.

Arpit Babbar, Sudarshan Kumar Kenettinkara, and Praveen Chandrashekar. Lax-wendroff flux reconstruction method for hyperbolic conservation laws. *Journal of Computational Physics*, 2022
<https://doi.org/10.1016/j.jcp.2022.111423>, <https://arxiv.org/abs/2207.02954>

Bibliography

- [1] Semih Akkurt, Freddie Witherden, and Peter Vincent. Cache blocking strategies applied to flux reconstruction. *Computer Physics Communications*, 271:108193, 2022.
- [2] Arpit Babbar, Sudarshan Kumar Kenettinkara, and Praveen Chandrashekhar. Lax-wendroff flux reconstruction method for hyperbolic conservation laws. *Journal of Computational Physics*, page 111423, 2022.
- [3] Christophe Berthon. Why the MUSCL–Hancock scheme is l1-stable. *Numerische Mathematik*, 104(1):27–46, jun 2006.
- [4] Jeff Bezanson, Alan Edelman, Stefan Karpinski, and Viral B Shah. Julia: a fresh approach to numerical computing. *SIAM Review*, 59(1):65–98, 2017.
- [5] Michael Dumbser and Olindo Zanotti. Very high order PNPM schemes on unstructured meshes for the resistive relativistic MHD equations. *Journal of Computational Physics*, 228(18):6991–7006, oct 2009.
- [6] Camille Felton, Mariana Harris, Caleb Logemann, Stefan Nelson, Ian Pelakh, and James A Rossmanith. A positivity-preserving limiting strategy for locally-implicit Lax-Wendroff discontinuous galerkin methods. Jun 2018.
- [7] Youngsoo Ha, Carl Gardner, Anne Gelb, and [\(keepcase|Chi Wang\)](#) Shu. Numerical simulation of high mach number astrophysical jets with radiative cooling. *Journal of Scientific Computing*, 24(1):597–612, jul 2005. Funding Information: We would like to thank Jeff Hester for valuable discussions. The authors would like to acknowledge for the following: (1) Research supported in part by the BK21 project at KAIST (Youngsoo Ha); (2) Research supported in part by the Space Telescope Science Institute under grant HST-GO-09863.06-A (Carl L. Gardner); (3) Research supported in part by the National Science Foundation under grant DMS-0107428. (Anne Gelb); (4) Research supported in part by the Army Research Office under grant DAAD19-00-1-0405 and in part by the National Science Foundation under grant DMS-0207451 (Chi-Wang Shu).
- [8] Sebastian Hennemann, Andrés M. Rueda-Ramírez, Florian J. Hindenlang, and Gregor J. Gassner. A provably entropy stable subcell shock capturing approach for high order split form dg for the compressible euler equations. *Journal of Computational Physics*, 426:109935, 2021.
- [9] H. T. Huynh. A Flux Reconstruction Approach to High-Order Schemes Including Discontinuous Galerkin Methods. Miami, FL, jun 2007. AIAA.

- [10] Haran Jackson. On the eigenvalues of the ADER-WENO galerkin predictor. *Journal of Computational Physics*, 333:409–413, mar 2017.
- [11] Randall J. LeVeque. High-Resolution Conservative Algorithms for Advection in Incompressible Flow. *SIAM Journal on Numerical Analysis*, 33(2):627–665, apr 1996. Bibtex: LeVeque1996.
- [12] Per-Olof Persson and Jaime Peraire. Sub-Cell Shock Capturing for Discontinuous Galerkin Methods. In *44th AIAA Aerospace Sciences Meeting and Exhibit*, Aerospace Sciences Meetings. American Institute of Aeronautics and Astronautics, jan 2006.
- [13] Hendrik Ranocha, Michael Schlottke-Lakemper, Andrew Ross Winters, Erik Faulhaber, Jesse Chan, and Gregor Gassner. Adaptive numerical simulations with Trixi.jl: A case study of Julia for scientific computing. *Proceedings of the JuliaCon Conferences*, 1(1):77, 2022.
- [14] A Rueda-Ramrez and G Gassner. A subcell finite volume positivity-preserving limiter for DGSEM discretizations of the euler equations. In *14th WCCM-ECCOMAS Congress*. CIMNE, 2021.
- [15] Kevin Schaal, Andreas Bauer, Praveen Chandrashekar, Rüdiger Pakmor, Christian Klingenberg, and Volker Springel. Astrophysical hydrodynamics with a high-order discontinuous Galerkin scheme and adaptive mesh refinement. *Monthly Notices of the Royal Astronomical Society*, 453(4):4278–4300, 09 2015.
- [16] L.I. SEDOV. Chapter iv - one-dimensional unsteady motion of a gas. In L.I. SEDOV, editor, *Similarity and Dimensional Methods in Mechanics*, pages 146–304. Academic Press, 1959.
- [17] Chi-Wang Shu and Stanley Osher. Efficient implementation of essentially non-oscillatory shock-capturing schemes, II. *Journal of Computational Physics*, 83(1):32–78, jul 1989.
- [18] Volker Springel. E pur si muove: Galilean-invariant cosmological hydrodynamical simulations on a moving mesh. *Monthly Notices of the Royal Astronomical Society*, 401(2):791–851, 01 2010.
- [19] Paul Woodward and Phillip Colella. The numerical simulation of two-dimensional fluid flow with strong shocks. *Journal of Computational Physics*, 54(1):115–173, apr 1984.
- [20] Xiangxiong Zhang and Chi-Wang Shu. On maximum-principle-satisfying high order schemes for scalar conservation laws. *Journal of Computational Physics*, 229(9):3091–3120, may 2010.
- [21] D. Zorio, A. Baeza, and P. Mulet. An Approximate Lax–Wendroff-Type Procedure for High Order Accurate Schemes for Hyperbolic Conservation Laws. *Journal of Scientific Computing*, 71(1):246–273, apr 2017. Bibtex: Zorio2017.

Thank you

45/75

Joint Work With

Praveen Chandrashekhar,

TIFR-CAM, Bangalore

Sudarshan Kumar Kenettinkara,

IISER-Trivandrum

$$(u_j^e)^{n+1} = (u_j^e)^n - \frac{\Delta t}{\Delta x_e} \frac{\partial F_h}{\partial \xi}(\xi_j), \quad 1 \leq j \leq N+1.$$

Conservation property of LWFR

$$(u_j^e)^{n+1} = (u_j^e)^n - \frac{\Delta t}{\Delta x_e} \frac{\partial F_h}{\partial \xi}(\xi_j), \quad 1 \leq j \leq N+1.$$

For $\{w_j\}_{j=1}^{N+1}$ being the quadrature weights associated with solution points:

$$\sum_{j=1}^{N+1} w_j (u_j^e)^{n+1} = \sum_{j=1}^{N+1} (u_j^e)^n - \frac{\Delta t}{\Delta x_e} \sum_{j=1}^{N+1} w_j \frac{\partial F_h}{\partial \xi}(\xi_j),$$

Conservation property of LWFR

$$(u_j^e)^{n+1} = (u_j^e)^n - \frac{\Delta t}{\Delta x_e} \frac{\partial F_h}{\partial \xi}(\xi_j), \quad 1 \leq j \leq N+1.$$

For $\{w_j\}_{j=1}^{N+1}$ being the quadrature weights associated with solution points:

$$\begin{aligned} \sum_{j=1}^{N+1} w_j (u_j^e)^{n+1} &= \sum_{j=1}^{N+1} (u_j^e)^n - \frac{\Delta t}{\Delta x_e} \sum_{j=1}^{N+1} w_j \frac{\partial F_h}{\partial \xi}(\xi_j), \\ \Rightarrow \bar{u}_e^{n+1} &= \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} \left[F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right]. \end{aligned}$$

High order

$$\bar{u}_e^{H,n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}}^H - F_{e-\frac{1}{2}}^H \right)$$

Low-order

$$\bar{u}_e^{L,n+1} = \bar{u}_e^n - \frac{\Delta t}{\Delta x_e} \left(F_{e+\frac{1}{2}}^L - F_{e-\frac{1}{2}}^L \right).$$

For blended update

$$\bar{u}_e^{n+1} = \bar{u}_e^n - \Delta t \left(F_{e+\frac{1}{2}} - F_{e-\frac{1}{2}} \right),$$

where $F_{e+\frac{1}{2}} = \alpha_e F_{e+\frac{1}{2}}^L + (1 - \alpha_e) F_{e+\frac{1}{2}}^H$.

Conservation requires

$$\alpha_e F_{e+\frac{1}{2}}^L + (1 - \alpha_e) F_{e+\frac{1}{2}}^H = \alpha_{e+1} F_{e+\frac{1}{2}}^L + (1 - \alpha_{e+1}) F_{e+\frac{1}{2}}^H$$

$$\Rightarrow F_{e+\frac{1}{2}}^L = F_{e+\frac{1}{2}}^H$$

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x + \frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x + \frac{1}{2}, e_y}) F_{e_x + \frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x + \frac{1}{2}, e_y} f_{\mathbf{e}, N + \frac{3}{2}, j}, & 1 \leq j \leq N + 1, \\ \tilde{F}_{e_x, e_y + \frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y + \frac{1}{2}}) F_{e_x, e_y + \frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y + \frac{1}{2}} f_{\mathbf{e}, i, N + \frac{3}{2}}, & 1 \leq i \leq N + 1.\end{aligned}$$

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x+\frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x+\frac{1}{2}, e_y}) F_{e_x+\frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x+\frac{1}{2}, e_y} f_{\mathbf{e}, N+\frac{3}{2}, j}, & 1 \leq j \leq N+1, \\ \tilde{F}_{e_x, e_y+\frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y+\frac{1}{2}}) F_{e_x, e_y+\frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y+\frac{1}{2}} f_{\mathbf{e}, i, N+\frac{3}{2}}, & 1 \leq i \leq N+1.\end{aligned}$$

$$\begin{aligned}\tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, j}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, j} - f_{\mathbf{e}, \frac{1}{2}, j} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_j} \left(f_{\mathbf{e}, 1, j+\frac{1}{2}} - f_{\mathbf{e}, 1, j-\frac{1}{2}} \right), \quad 1 < j < N+1, \\ \tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, 1}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, 1} - f_{\mathbf{e}, \frac{1}{2}, 1} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x, e_y+\frac{1}{2}, 1} - f_{\mathbf{e}, 1, \frac{1}{2}} \right), \quad j = 1.\end{aligned}$$

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x+\frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x+\frac{1}{2}, e_y}) F_{e_x+\frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x+\frac{1}{2}, e_y} f_{\mathbf{e}, N+\frac{3}{2}, j}, & 1 \leq j \leq N+1, \\ \tilde{F}_{e_x, e_y+\frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y+\frac{1}{2}}) F_{e_x, e_y+\frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y+\frac{1}{2}} f_{\mathbf{e}, i, N+\frac{3}{2}}, & 1 \leq i \leq N+1.\end{aligned}$$

$$\begin{aligned}\tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, j}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, j} - f_{\mathbf{e}, \frac{1}{2}, j} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_j} \left(f_{\mathbf{e}, 1, j+\frac{1}{2}} - f_{\mathbf{e}, 1, j-\frac{1}{2}} \right), \quad 1 < j < N+1, \\ \tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, 1}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, 1} - f_{\mathbf{e}, \frac{1}{2}, 1} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x, e_y+\frac{1}{2}, 1} - f_{\mathbf{e}, 1, \frac{1}{2}} \right), \quad j = 1.\end{aligned}$$

In the 2-D code, there's two separate face loops for vertical and horizontal faces. This poses a challenge because to ensure $\tilde{\mathbf{u}}^{n+1}$ is admissible, we need to correct both $\tilde{F}_{e_x+\frac{1}{2}, e_y, 1}$ and $\tilde{F}_{e_x, e_y+\frac{1}{2}, 1}$ and these values are never available together.

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x+\frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x+\frac{1}{2}, e_y}) F_{e_x+\frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x+\frac{1}{2}, e_y} f_{\mathbf{e}, N+\frac{3}{2}, j}, & 1 \leq j \leq N+1, \\ \tilde{F}_{e_x, e_y+\frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y+\frac{1}{2}}) F_{e_x, e_y+\frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y+\frac{1}{2}} f_{\mathbf{e}, i, N+\frac{3}{2}}, & 1 \leq i \leq N+1.\end{aligned}$$

$$\begin{aligned}\tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, j}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, j} - f_{\mathbf{e}, \frac{1}{2}, j} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_j} \left(f_{\mathbf{e}, 1, j+\frac{1}{2}} - f_{\mathbf{e}, 1, j-\frac{1}{2}} \right), \quad 1 < j < N+1, \\ \tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, 1}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, 1} - f_{\mathbf{e}, \frac{1}{2}, 1} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x, e_y+\frac{1}{2}, 1} - f_{\mathbf{e}, 1, \frac{1}{2}} \right), \quad j = 1.\end{aligned}$$

In the 2-D code, there's two separate face loops for vertical and horizontal faces. This poses a challenge because to ensure $\tilde{\mathbf{u}}^{n+1}$ is admissible, we need to correct both $\tilde{F}_{e_x+\frac{1}{2}, e_y, 1}$ and $\tilde{F}_{e_x, e_y+\frac{1}{2}, 1}$ and these values are never available together.

To avoid having to store values and doing aposteriori correction, we find appropriate λ_x, λ_y such that

$$\lambda_x + \lambda_y = 1,$$

and then, following the 1-D procedure, construct corrected $F_{e_x+\frac{1}{2}, e_y, 1}$ and $F_{e_x, e_y+\frac{1}{2}, 1}$ such that

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x+\frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x+\frac{1}{2}, e_y}) F_{e_x+\frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x+\frac{1}{2}, e_y} f_{\mathbf{e}, N+\frac{3}{2}, j}, & 1 \leq j \leq N+1, \\ \tilde{F}_{e_x, e_y+\frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y+\frac{1}{2}}) F_{e_x, e_y+\frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y+\frac{1}{2}} f_{\mathbf{e}, i, N+\frac{3}{2}}, & 1 \leq i \leq N+1.\end{aligned}$$


$$\begin{aligned}\tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, j}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, j} - f_{\mathbf{e}, \frac{1}{2}, j} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_j} \left(f_{\mathbf{e}, 1, j+\frac{1}{2}} - f_{\mathbf{e}, 1, j-\frac{1}{2}} \right), \quad 1 < j < N+1, \\ \tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, 1}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, 1} - f_{\mathbf{e}, \frac{1}{2}, 1} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x, e_y+\frac{1}{2}, 1} - f_{\mathbf{e}, 1, \frac{1}{2}} \right), \quad j = 1.\end{aligned}$$

In the 2-D code, there's two separate face loops for vertical and horizontal faces. This poses a challenge because to ensure $\tilde{\mathbf{u}}^{n+1}$ is admissible, we need to correct both $\tilde{F}_{e_x+\frac{1}{2}, e_y, 1}$ and $\tilde{F}_{e_x, e_y+\frac{1}{2}, 1}$ and these values are never available together.

To avoid having to store values and doing aposteriori correction, we find appropriate λ_x, λ_y such that

$$\lambda_x + \lambda_y = 1,$$

and then, following the 1-D procedure, construct corrected $F_{e_x+\frac{1}{2}, e_y, 1}$ and $F_{e_x, e_y+\frac{1}{2}, 1}$ such that



$$\begin{aligned} \boldsymbol{u}_{\boldsymbol{e},1,1}^n - \frac{\Delta t}{\Delta x_{\boldsymbol{e}} \lambda_x w_1} \left(\tilde{F}_{e_x + \frac{1}{2}, e_y, 1} - f_{\boldsymbol{e}, \frac{1}{2}, 1} \right) &\in \Omega, \\ \boldsymbol{u}_{\boldsymbol{e},1,1}^n - \frac{\Delta t}{\Delta y_{\boldsymbol{e}} \lambda_y w_1} \left(\tilde{F}_{e_x, e_y + \frac{1}{2}, 1} - f_{\boldsymbol{e}, 1, \frac{1}{2}} \right) &\in \Omega. \end{aligned}$$

Admissibility preservation in 2-D

Initial candidate:

$$\begin{aligned}\tilde{F}_{e_x+\frac{1}{2}, e_y, j} &= (1 - \alpha_{e_x+\frac{1}{2}, e_y}) F_{e_x+\frac{1}{2}, e_y, j}^{\text{LW}} + \alpha_{e_x+\frac{1}{2}, e_y} f_{\mathbf{e}, N+\frac{3}{2}, j}, & 1 \leq j \leq N+1, \\ \tilde{F}_{e_x, e_y+\frac{1}{2}, i} &= (1 - \alpha_{e_x, e_y+\frac{1}{2}}) F_{e_x, e_y+\frac{1}{2}, i}^{\text{LW}} + \alpha_{e_x, e_y+\frac{1}{2}} f_{\mathbf{e}, i, N+\frac{3}{2}}, & 1 \leq i \leq N+1.\end{aligned}$$


$$\begin{aligned}\tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, j}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, j} - f_{\mathbf{e}, \frac{1}{2}, j} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_j} \left(f_{\mathbf{e}, 1, j+\frac{1}{2}} - f_{\mathbf{e}, 1, j-\frac{1}{2}} \right), & 1 < j < N+1, \\ \tilde{\mathbf{u}}^{n+1} &= \mathbf{u}_{\mathbf{e}, 1, 1}^n - \frac{\Delta t}{\Delta x_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x+\frac{1}{2}, e_y, 1} - f_{\mathbf{e}, \frac{1}{2}, 1} \right) - \frac{\Delta t}{\Delta y_{\mathbf{e}} w_1} \left(\tilde{F}_{e_x, e_y+\frac{1}{2}, 1} - f_{\mathbf{e}, 1, \frac{1}{2}} \right), & j = 1.\end{aligned}$$

In the 2-D code, there's two separate face loops for vertical and horizontal faces. This poses a challenge because to ensure $\tilde{\mathbf{u}}^{n+1}$ is admissible, we need to correct both $\tilde{F}_{e_x+\frac{1}{2}, e_y, 1}$ and $\tilde{F}_{e_x, e_y+\frac{1}{2}, 1}$ and these values are never available together.

To avoid having to store values and doing aposteriori correction, we find appropriate λ_x, λ_y such that

$$\lambda_x + \lambda_y = 1,$$


and then, following the 1-D procedure, construct corrected $F_{e_x+\frac{1}{2}, e_y, 1}$ and $F_{e_x, e_y+\frac{1}{2}, 1}$ such that




$$\begin{aligned} \boldsymbol{u}_{\boldsymbol{e},1,1}^n - \frac{\Delta t}{\Delta x_{\boldsymbol{e}} \lambda_x w_1} \left(\tilde{F}_{e_x + \frac{1}{2}, e_y, 1} - f_{\boldsymbol{e}, \frac{1}{2}, 1} \right) &\in \Omega, \\ \boldsymbol{u}_{\boldsymbol{e},1,1}^n - \frac{\Delta t}{\Delta y_{\boldsymbol{e}} \lambda_y w_1} \left(\tilde{F}_{e_x, e_y + \frac{1}{2}, 1} - f_{\boldsymbol{e}, 1, \frac{1}{2}} \right) &\in \Omega. \end{aligned}$$

$$\lambda_x = \frac{|s_x^e| / \Delta x_{\boldsymbol{e}}}{|s_x^e| / \Delta x_{\boldsymbol{e}} + |s_y^e| / \Delta y_{\boldsymbol{e}}}, \quad \lambda_y = \frac{|s_y^e| / \Delta y_{\boldsymbol{e}}}{|s_x^e| / \Delta x_{\boldsymbol{e}} + |s_y^e| / \Delta y_{\boldsymbol{e}}}$$


$$\mathbb{E} = \max \left(\frac{m_{N+1}^2}{\beta_1 m_1 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1 + \sum_{j=2}^N m_j^2} \right)$$



$$\mathbb{E} = \max \left(\frac{m_{N+1}^2}{\beta_1 m_1 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1 + \sum_{j=2}^N m_j^2} \right)$$



$$\mathbb{E} = \max \left(\frac{m_{N+1}^2}{\beta_1 m_1 + \sum_{j=2}^{N+1} m_j^2}, \frac{m_N^2}{\beta_2 m_1 + \sum_{j=2}^N m_j^2} \right)$$



Lemma. Consider the 1-D Riemann problem

$$\begin{aligned}\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x &= 0, \\ \mathbf{u}(x, 0) &= \begin{cases} \mathbf{u}_l, & x < 0, \\ \mathbf{u}_r, & x > 0, \end{cases}\end{aligned}$$

in $[-h, h] \times [0, \Delta t]$ where

$$\frac{\Delta t}{h} |\sigma_e(\mathbf{u}_l, \mathbf{u}_r)| \leq \frac{1}{2},$$

where $\sigma_e(\mathbf{u}_1, \mathbf{u}_2)$ denotes all eigenvalues of all Jacobian matrices at the states between \mathbf{u}_1 and \mathbf{u}_2 . Then, for all $t \leq \Delta t$, we have

$$\int_{-h}^h \mathbf{u}(x, t) dx = h (\mathbf{u}_l + \mathbf{u}_r) - t (\mathbf{f}(\mathbf{u}_r) - \mathbf{f}(\mathbf{u}_l)).$$

Lemma 1. (*Evolution*) Pick

$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}},$$

so that

$$\frac{\mu^-}{2} \mathbf{u}_i^{n,-} + \frac{1}{2} \mathbf{u}_i^{*,\pm} + \frac{\mu^+}{2} \mathbf{u}_i^{n,+} = \mathbf{u}_i^{n,\pm}.$$

Then, assume that

$$\mathbf{u}_i^{n,\pm} \in \mathcal{U}_{\text{ad}} \quad \text{and} \quad \mathbf{u}_i^{*,\pm} \in \mathcal{U}_{\text{ad}},$$


and the CFL restrictions

$$\begin{aligned} \frac{\Delta t / 2}{\mu^- \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{n,-}, \mathbf{u}_i^{*,\pm})|) &\leq \frac{1}{2}, \\ \frac{\Delta t / 2}{\mu^+ \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{*,\pm}, \mathbf{u}_i^{n,+})|) &\leq \frac{1}{2}, \end{aligned} \tag{1}$$


where $\sigma_e(\mathbf{u}_1, \mathbf{u}_2)$ denotes the maximum spectral radius among all Jacobian matrices at states between \mathbf{u}_1 and \mathbf{u}_2 .

Then, we have invariance of \mathcal{U}_{ad} under the first step of MUSCL-Hancock scheme, i.e.,

$$\mathbf{u}_i^{n+1/2,\pm} \in \mathcal{U}_{\text{ad}}.$$

Proof

Proof



$$\begin{aligned}
 \frac{1}{\Delta x_i} \int_{x_{i-1/2}}^{x_{i+1/2}} \mathbf{u}^h(x, \Delta t/2) dx &= \frac{1}{2} (\mu^- \mathbf{w}_i^{n,-} + \mathbf{u}_i^{*,+} + \mu^+ \mathbf{u}_i^{n,+}) - \frac{\Delta t/2}{\Delta x_i} (f(\mathbf{u}_i^{n,+}) - f(\mathbf{u}_i^{n,-})) \\
 &= \mathbf{u}_i^{n,+} - \frac{\Delta t/2}{\Delta x_i} (f(\mathbf{u}_i^{n,+}) - f(\mathbf{u}_i^{n,-})) = \mathbf{u}_i^{n+\frac{1}{2},+}
 \end{aligned}$$

□

Step 2 : FVM type update

53/75


Define $\mathbf{u}_i^{n+\frac{1}{2}, *}$

$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n.$$

Step 2 : FVM type update

Define $\mathbf{u}_i^{n+\frac{1}{2}, *}$


$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n.$$



Step 2 : FVM type update

Define $\mathbf{u}_i^{n+\frac{1}{2}, *}$

$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n.$$




$$\begin{aligned}
 \mathbf{u}_i^{n+1, -} &:= \mathbf{u}_i^{n+\frac{1}{2}, -} - \frac{\Delta t}{\mu^- \Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *}\right) - f\left(\mathbf{u}_{i-1}^{n+\frac{1}{2}, +}, \mathbf{u}_i^{n+\frac{1}{2}, -}\right) \right) \\
 \mathbf{u}_i^{n+1, *} &:= \mathbf{u}_i^{n+\frac{1}{2}, *} - \frac{\Delta t}{\Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +}\right) - f\left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *}\right) \right) \\
 \mathbf{u}_i^{n+1, +} &:= \mathbf{u}_i^{n+\frac{1}{2}, +} - \frac{\Delta t}{\mu^+ \Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, +}, \mathbf{u}_{i+1}^{n+\frac{1}{2}, -}\right) - f\left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +}\right) \right)
 \end{aligned}$$

Step 2 : FVM type update

Define $\mathbf{u}_i^{n+\frac{1}{2}, *}$

$$\frac{\mu^-}{2} \mathbf{u}_i^{n+\frac{1}{2}, -} + \frac{1}{2} \mathbf{u}_i^{n+\frac{1}{2}, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+\frac{1}{2}, +} = \mathbf{u}_i^n.$$



$$\mathbf{u}_i^{n+1, -} := \mathbf{u}_i^{n+\frac{1}{2}, -} - \frac{\Delta t}{\mu^- \Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *}\right) - f\left(\mathbf{u}_{i-1}^{n+\frac{1}{2}, +}, \mathbf{u}_i^{n+\frac{1}{2}, -}\right) \right)$$

$$\mathbf{u}_i^{n+1, *} := \mathbf{u}_i^{n+\frac{1}{2}, *} - \frac{\Delta t}{\Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +}\right) - f\left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *}\right) \right)$$

$$\mathbf{u}_i^{n+1, +} := \mathbf{u}_i^{n+\frac{1}{2}, +} - \frac{\Delta t}{\mu^+ \Delta x_i / 2} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, +}, \mathbf{u}_{i+1}^{n+\frac{1}{2}, -}\right) - f\left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +}\right) \right)$$

$$\begin{aligned} \mathbf{u}_i^{n+1} &= \mathbf{u}_i^n - \frac{\Delta t}{\Delta x} \left(f\left(\mathbf{u}_i^{n+\frac{1}{2}, +}, \mathbf{u}_{i+1}^{n+\frac{1}{2}, -}\right) - f\left(\mathbf{u}_{i-1}^{n+\frac{1}{2}, +}, \mathbf{u}_i^{n+\frac{1}{2}, -}\right) \right) \\ &= \frac{\mu^-}{2} \mathbf{u}_i^{n+1, -} + \frac{1}{2} \mathbf{u}_i^{n+1, *} + \frac{\mu^+}{2} \mathbf{u}_i^{n+1, +} \end{aligned}$$

Step 2 : FVM type update

Lemma 2. (*Riemann solver*) Assume that the states $(\mathbf{u}_i^{n+\frac{1}{2}, \pm})_{i \in \mathbb{Z}}, (\mathbf{u}_i^{n+\frac{1}{2}, *})_{i \in \mathbb{Z}}$ belong to \mathcal{U}_{ad} , where $\mathbf{u}_i^{n+\frac{1}{2}, *}$ was defined above as

$$\mu^- \mathbf{u}_i^{n+\frac{1}{2}, -} + \mathbf{u}_i^{n+\frac{1}{2}, *} + \mu^+ \mathbf{u}_i^{n+\frac{1}{2}, +} = 2\mathbf{u}_i^n.$$

Then, the updated solution of MUSCL-Hancock scheme is in Ω under the CFL conditions

$$\begin{aligned}
 & \frac{\Delta t}{\mu^- \Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *} \right) \right| \right) \leq \frac{1}{2}, \\
 & \frac{\Delta t}{\mu^- \Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_{i-1}^{n+\frac{1}{2}, +}, \mathbf{u}_i^{n+\frac{1}{2}, -} \right) \right| \right) \leq \frac{1}{2}, \\
 & \frac{\Delta t}{\Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +} \right) \right| \right) \leq \frac{1}{2}, \\
 & \frac{\Delta t}{\Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_i^{n+\frac{1}{2}, -}, \mathbf{u}_i^{n+\frac{1}{2}, *} \right) \right| \right) \leq \frac{1}{2}, \\
 & \frac{\Delta t}{\mu^+ \Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_i^{n+\frac{1}{2}, +}, \mathbf{u}_{i+1}^{n+\frac{1}{2}, -} \right) \right| \right) \leq \frac{1}{2}, \\
 & \frac{\Delta t}{\mu^+ \Delta x_i / 2} \max_{i \in \mathbb{Z}} \left(\left| \sigma_e \left(\mathbf{u}_i^{n+\frac{1}{2}, *}, \mathbf{u}_i^{n+\frac{1}{2}, +} \right) \right| \right) \leq \frac{1}{2}.
 \end{aligned} \tag{2}$$

$$\mu^- \mathbf{u}_i^{n+\frac{1}{2}, -} + \mathbf{u}_i^{n+\frac{1}{2}, *} + \mu^+ \mathbf{u}_i^{n+\frac{1}{2}, +} = 2\mathbf{u}_i^n$$

$$\mathbf{u}_i^{n+\frac{1}{2}, *} = (2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+})) - \frac{\Delta t}{2\Delta x_i} [(f(\mathbf{u}_i^{n,-}) - f(\mathbf{u}_i^{n,+}))]$$

$$\mathbf{u}_i^{n+\frac{1}{2},*} = (2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+})) - \frac{\Delta t}{2\Delta x_i} [(f(\mathbf{u}_i^{n,-}) - f(\mathbf{u}_i^{n,+}))]$$

Lemma 7. (*Link previous lemmas*) Define $\mathbf{u}_i^{*,*}$ to satisfy

$$\mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,*} + \mu^+ \mathbf{u}_i^{n,+} = 2(2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+})), \quad (7)$$

where, as defined before,

$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}}.$$

Assume that $\mathbf{u}_i^{n,\pm}$ and $\mathbf{u}_i^{*,*}$ are in \mathcal{U}_{ad} . Consider the CFL conditions

$$\begin{aligned} \frac{\Delta t / 2}{\mu^- \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{*,*}, \mathbf{u}_i^{n,-})|) &\leq \frac{1}{2}, \\ \frac{\Delta t / 2}{\mu^+ \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{n,+}, \mathbf{u}_i^{*,*})|) &\leq \frac{1}{2}, \end{aligned} \quad (8)$$

then $\mathbf{u}_i^{n+\frac{1}{2},*} \in \mathcal{U}_{\text{ad}}$.

$$\mathbf{u}_i^{n+\frac{1}{2},*} = (2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+})) - \frac{\Delta t}{2\Delta x_i} [(f(\mathbf{u}_i^{n,-}) - f(\mathbf{u}_i^{n,+}))]$$

Lemma 9. (*Link previous lemmas*) Define $\mathbf{u}_i^{*,*}$ to satisfy

$$\mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,*} + \mu^+ \mathbf{u}_i^{n,+} = 2(2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+})), \quad (9)$$

where, as defined before,


$$\mu^- = \frac{x_{i+1/2} - x_i}{x_{i+1/2} - x_{i-1/2}}, \quad \mu^+ = \frac{x_i - x_{i-1/2}}{x_{i+1/2} - x_{i-1/2}}.$$

Assume that $\mathbf{u}_i^{n,\pm}$ and $\mathbf{u}_i^{*,*}$ are in \mathcal{U}_{ad} . Consider the CFL conditions


$$\begin{aligned} \frac{\Delta t / 2}{\mu^- \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{*,*}, \mathbf{u}_i^{n,-})|) &\leq \frac{1}{2}, \\ \frac{\Delta t / 2}{\mu^+ \Delta x / 2} \max_{i \in \mathbb{Z}} (|\sigma_e(\mathbf{u}_i^{n,+}, \mathbf{u}_i^{*,*})|) &\leq \frac{1}{2}, \end{aligned} \quad (10)$$

then $\mathbf{u}_i^{n+\frac{1}{2},*} \in \mathcal{U}_{\text{ad}}$.

Remark 10. For conservative reconstruction, we actually have $\mathbf{u}_i^{*,*} = \mathbf{u}_i$. So, this lemma isn't placing new restrictions.

Proof.

Proof.



$$\begin{aligned}
 \frac{1}{\Delta x_i} \int_{x_{i-1/2}}^{x_{i+1/2}} \mathbf{u}^h \left(x, \frac{\Delta t}{2} \right) dx &= \frac{1}{2} (\mu^+ \mathbf{u}_i^{n,+} + \mathbf{u}_i^{*,*} + \mu^- \mathbf{u}_i^{n,-}) - \frac{\Delta t}{2 \Delta x_i} (f(\mathbf{u}_i^{n,-}) - f(\mathbf{u}_i^{n,+})) \\
 &= \mathbf{u}_i^{n+\frac{1}{2},*}
 \end{aligned}$$

Theorem 11. Consider the conservation law

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$

which preserves a convex set \mathcal{U}_{ad} . Let $\{\mathbf{u}_i^n\}_{i \in \mathbb{Z}}$ be the approximate solution at time level n and assume that $\mathbf{u}_i^n \in \mathcal{U}_{\text{ad}}$ for all $i \in \mathbb{Z}$. Consider **conservative reconstructions**

$$\mathbf{u}_i^{n,+} = \mathbf{u}_i + (x_{i+1/2} - x_i) \boldsymbol{\sigma}_i, \quad \mathbf{u}_i^{n,-} = \mathbf{u}_i + (x_{i-1/2} - x_i) \boldsymbol{\sigma}_i.$$

Define $\mathbf{u}_i^{*,\pm}$ to be

$$\mathbf{u}_i^{*,\pm} = \mathbf{u}_i^n + 2\left(x_{i \pm \frac{1}{2}} - x_i\right) \boldsymbol{\sigma}_i$$

and assume that the slope $\boldsymbol{\sigma}_i$ is chosen so that

$$\mathbf{u}_i^{*,\pm} \in \mathcal{U}_{\text{ad}}.$$

Then, under time step restrictions (1), (2), (10), the updated solution \mathbf{u}_i^{n+1} , defined by the MUSCL-Hancock procedure is in \mathcal{U}_{ad} .

Theorem 12. Consider the conservation law

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = 0$$

which preserves a convex set \mathcal{U}_{ad} . Let $\{\mathbf{u}_i^n\}_{i \in \mathbb{Z}}$ be the approximate solution at time level n and assume that $\mathbf{u}_i^n \in \mathcal{U}_{\text{ad}}$ for all $i \in \mathbb{Z}$. Consider **conservative reconstructions**

$$\mathbf{u}_i^{n,+} = \mathbf{u}_i + (x_{i+1/2} - x_i) \boldsymbol{\sigma}_i, \quad \mathbf{u}_i^{n,-} = \mathbf{u}_i + (x_{i-1/2} - x_i) \boldsymbol{\sigma}_i.$$

Define $\mathbf{u}_i^{*,\pm}$ to be

$$\mathbf{u}_i^{*,\pm} = \mathbf{u}_i^n + 2 \left(x_{i \pm \frac{1}{2}} - x_i \right) \boldsymbol{\sigma}_i$$

and assume that the slope σ_i is chosen so that

$$\mathbf{u}_i^{*,\pm} \in \mathcal{U}_{\text{ad}}.$$

Then, under time step restrictions (1), (2), (10), the updated solution \mathbf{u}_i^{n+1} , defined by the MUSCL-Hancock procedure is in \mathcal{U}_{ad} .

Proof. We only need $\mathbf{u}_i^{n,\pm} \in \mathcal{U}_{\text{ad}}$ to apply the Lemmas 1, 2, 9. To that end, notice

$$\mathbf{u}_i^{n,\pm} = \frac{1}{2} \mathbf{u}_i^{*,\pm} + \frac{1}{2} \mathbf{u}_i^n.$$

□

Enforcing slope restriction

Given candidate slope σ_i ,

$$\mathbf{u}_i^{*,\pm} := \mathbf{u}_i^n + 2(x_{i\pm 1/2} - x_i) \sigma_i.$$

Find $\theta \in [0, 1]$

$$\mathbf{u}_i^n + 2(x_{i\pm 1/2} - x_i) \theta \sigma_i \in \mathcal{U}_{\text{ad}}. \quad (11)$$

For **concave** $p = p(\mathbf{u})$, assume

$$\mathcal{U}_{\text{ad}} = \{\mathbf{u} \in \Omega : p(\mathbf{u}) > 0\}$$

We pick

$$\theta_{\pm} = \min \left(\left| \frac{\epsilon - p(\mathbf{u}_i^n)}{p(\mathbf{u}_i^{*,\pm}) - p(\mathbf{u}_i^n)} \right|, 1 \right)$$

and

$$\theta = \min(\theta_+, \theta_-).$$

By concavity,

$$p(\theta \mathbf{u}_i^{*,\pm} + (1-\theta) \mathbf{u}_i^n) > \theta p(\mathbf{u}_i^{*,\pm}) + (1-\theta) p(\mathbf{u}_i^n) > \epsilon > 0.$$

Thus, this θ will give us (11).

Consider non-conservative variables

$$\mathbf{U}_i^n = \kappa(\mathbf{u}_i^n),$$

so that reconstruction is given by

$$\begin{aligned} \mathbf{U}^n(x) &= \mathbf{U}_i^n + \sigma_i(x - x_i) \\ \mathbf{u}_i^{n,\pm} &:= \kappa^{-1}(\mathbf{U}_i^{n,\pm}) \end{aligned} \tag{12}$$

Theorem 13. Assume that $\mathbf{u}_i^n \in \Omega$ for all $i \in \mathbb{Z}$. Consider $\mathbf{u}_i^{n,\pm}$ defined in (12), $\mathbf{u}_i^{*,\pm}$ defined so that

$$\mu^- \mathbf{u}_i^{n+\frac{1}{2},-} + \mathbf{u}_i^{n+\frac{1}{2},*} + \mu^+ \mathbf{u}_i^{n+\frac{1}{2},+} = 2\mathbf{u}_i^n,$$

and $\mathbf{u}_i^{*,*}$ defined explicitly as

$$\mathbf{u}_i^{*,*} = 4\mathbf{u}_i^n - 3(\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+}).$$

Assume that the slope is chosen so that

$$\mathbf{u}_i^{n,\pm} \in \Omega, \quad \mathbf{u}_i^{*,\pm} \in \Omega \quad \text{and} \quad \mathbf{u}_i^{*,*} \in \Omega.$$

Consider the same CFL conditions (1), (2), (10). Then the updated solution \mathbf{u}_i^{n+1} of MUSCL-Hancock procedure is in Ω .

Remark 14. The definition of $\mathbf{u}_i^{*,*}$ comes from

$$\color{red} \mu^- \mathbf{u}_i^{n,-} + \mathbf{u}_i^{*,*} + \color{red} \mu^+ \mathbf{u}_i^{n,+} = 2(2\mathbf{u}_i^n - (\mu^- \mathbf{u}_i^{n,-} + \mu^+ \mathbf{u}_i^{n,+}))$$

Current solution

$$x \in I_e: \quad u_h^n(\xi) = \sum_{j=1}^{N+1} u_j^n \ell_j(\xi)$$

ADER-DG : Predictor step

Current solution

$$x \in I_e: \quad u_h^n(\xi) = \sum_{j=1}^{N+1} u_j^n \ell_j(\xi)$$

Cell local space-time solution and flux: $\tau = (t - t_n) / \Delta t$

$$\tilde{u}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} \tilde{u}_{rs} \ell_r(\xi) \ell_s(\tau), \quad \tilde{f}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} f(\tilde{u}_{rs}) \ell_r(\xi) \ell_s(\tau).$$

Current solution

$$x \in I_e: \quad u_h^n(\xi) = \sum_{j=1}^{N+1} u_j^n \ell_j(\xi)$$

Cell local space-time solution and flux: $\tau = (t - t_n) / \Delta t$

$$\tilde{u}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} \tilde{u}_{rs} \ell_r(\xi) \ell_s(\tau), \quad \tilde{f}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} f(\tilde{u}_{rs}) \ell_r(\xi) \ell_s(\tau).$$

Find \tilde{u}_h by cell local Galerkin method

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t \tilde{u}_h + \partial_x \tilde{f}_h) \ell_r(\xi) \ell_s(\tau) dx dt, \quad 1 \leq r, s \leq N+1.$$

Current solution

$$x \in I_e: \quad u_h^n(\xi) = \sum_{j=1}^{N+1} u_j^n \ell_j(\xi)$$

Cell local space-time solution and flux: $\tau = (t - t_n) / \Delta t$

$$\tilde{u}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} \tilde{u}_{rs} \ell_r(\xi) \ell_s(\tau), \quad \tilde{f}_h(\xi, \tau) = \sum_{r,s=1}^{N+1} f(\tilde{u}_{rs}) \ell_r(\xi) \ell_s(\tau).$$

Find \tilde{u}_h by cell local Galerkin method

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t \tilde{u}_h + \partial_x \tilde{f}_h) \ell_r(\xi) \ell_s(\tau) dx dt, \quad 1 \leq r, s \leq N+1.$$

Integrate by parts in time

$$\begin{aligned} & - \int_{t_n}^{t_{n+1}} \int_{I_e} \tilde{u}_h \ell_r(\xi) \partial_t \ell_s(\tau) dx dt + \int_{I_e} \tilde{u}_h(\xi, 1) \ell_r(\xi) \ell_s(1) dx - \int_{I_e} u_h^n(\xi) \ell_r(\xi) \ell_s(0) d\xi \\ & + \int_{t_n}^{t_{n+1}} \int_{I_e} \partial_x \tilde{f}_h \ell_r(\xi) \ell_s(\tau) dx dt = 0. \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) \, dx \, dt = 0$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) \, dx \, dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) \, dx &= \int_{I_e} u_h^n \ell_i(\xi) \, dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) \, dx \, dt \\ &\quad + \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) \, dt - \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) \, dt. \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) dx dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) dt - \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) dt. \end{aligned}$$

FR form

$$\begin{aligned} u_i^n - \int_{t^n}^{t^{n+1}} \partial_x \tilde{f}_h(\xi_i, \tau) dt \\ \Rightarrow u_i^{n+1} = -g'_L(\xi_i) \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) dt \\ - g'_R(\xi_i) \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) dt \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) \, dx \, dt = 0$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) dx dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) dt - \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) dt. \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) dx dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) dt - \int_{t^n}^{t^{n+1}} \textcolor{blue}{f}(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) dt. \end{aligned}$$

Another integration by parts

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx - \int_{t^n}^{t^{n+1}} \int_{I_e} \partial_x \tilde{f}_h \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) \right) \ell_i(0) dt \\ &\quad - \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) \right) \ell_i(1) dt, \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) dx dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) dt - \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) dt. \end{aligned}$$

Another integration by parts

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx - \int_{t^n}^{t^{n+1}} \int_{I_e} \partial_x \tilde{f}_h \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) \right) \ell_i(0) dt \\ &\quad - \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) \right) \ell_i(1) dt, \end{aligned}$$

Quadrature on solution points

$$\begin{aligned} & u_i^n - \int_{t^n}^{t^{n+1}} \partial_x \tilde{f}_h(\xi_i, \tau) dt \\ \Rightarrow u_i^{n+1} = & + \frac{\ell_i(0)}{w_i} \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) dt \\ & - \frac{\ell_i(1)}{w_i} \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) dt \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} \int_{I_e} (\partial_t u_h + \partial_x f(u_h)) \ell_i(\xi) dx dt = 0$$

Integrate by parts in space

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx + \int_{t^n}^{t^{n+1}} \int_{I_e} \tilde{f}_h \partial_x \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) \ell_i(0) dt - \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) \ell_i(1) dt. \end{aligned}$$

Another integration by parts

$$\begin{aligned} \int_{I_e} u_h^{n+1} \ell_i(\xi) dx &= \int_{I_e} u_h^n \ell_i(\xi) dx - \int_{t^n}^{t^{n+1}} \int_{I_e} \partial_x \tilde{f}_h \ell_i(\xi) dx dt \\ &\quad + \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) \right) \ell_i(0) dt \\ &\quad - \int_{t^n}^{t^{n+1}} \left(f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) \right) \ell_i(1) dt, \end{aligned}$$

Quadrature on solution points

$$\begin{aligned} & u_i^n - \int_{t^n}^{t^{n+1}} \partial_x \tilde{f}_h(\xi_i, \tau) dt \\ \Rightarrow u_i^{n+1} = & -g'_L(\xi_i) \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e-\frac{1}{2}}^-(t), \tilde{u}_{e-\frac{1}{2}}^+(t)) - \tilde{f}_h(0, \tau) dt \\ & - g'_R(\xi_i) \int_{t^n}^{t^{n+1}} f(\tilde{u}_{e+\frac{1}{2}}^-(t), \tilde{u}_{e+\frac{1}{2}}^+(t)) - \tilde{f}_h(1, \tau) dt \end{aligned}$$

For

$$u_t + u_x = 0,$$

ADER update

$$\begin{aligned} u_i^n - \partial_x \int_{t^n}^{t^{n+1}} \tilde{u}_h(\xi_i, \tau) dt \\ u_i^{n+1} = -g'_L(\xi_i) \left[f \left(\int_{t^n}^{t^{n+1}} \tilde{u}_{e-\frac{1}{2}}^-(t) dt, \int_{t^n}^{t^{n+1}} \tilde{u}_{e-\frac{1}{2}}^+(t) dt \right) - \int_{t^n}^{t^{n+1}} \tilde{u}_h(0, \tau) dt \right] \\ - g'_R(\xi_i) \left[f \left(\int_{t^n}^{t^{n+1}} \tilde{u}_{e+\frac{1}{2}}^-(t) dt, \int_{t^n}^{t^{n+1}} \tilde{u}_{e+\frac{1}{2}}^+(t) dt \right) - \int_{t^n}^{t^{n+1}} \tilde{u}_h(1, \tau) dt \right] \end{aligned} \quad (13)$$

For

$$u_t + u_x = 0,$$

ADER update

$$\begin{aligned} u_i^n - \partial_x \int_{t^n}^{t^{n+1}} \tilde{u}_h(\xi_i, \tau) dt \\ u_i^{n+1} = -g'_L(\xi_i) \left[f \left(\int_{t^n}^{t^{n+1}} \tilde{u}_{e-\frac{1}{2}}^-(t) dt, \int_{t^n}^{t^{n+1}} \tilde{u}_{e-\frac{1}{2}}^+(t) dt \right) - \int_{t^n}^{t^{n+1}} \tilde{u}_h(0, \tau) dt \right] \\ - g'_R(\xi_i) \left[f \left(\int_{t^n}^{t^{n+1}} \tilde{u}_{e+\frac{1}{2}}^-(t) dt, \int_{t^n}^{t^{n+1}} \tilde{u}_{e+\frac{1}{2}}^+(t) dt \right) - \int_{t^n}^{t^{n+1}} \tilde{u}_h(1, \tau) dt \right] \end{aligned} \quad (15)$$

LWFR-D2 update

$$u_i^{n+1} = u_i^n - \Delta t \left[\partial_x U_h(\xi_i) - g'_L(\xi_i) \left[f(U_{e-\frac{1}{2}}^-, U_{e-\frac{1}{2}}^+) - U_h(0) \right] - g'_R(\xi_i) \left[f(U_{e+\frac{1}{2}}^-, U_{e+\frac{1}{2}}^+) - U_h(1) \right] \right], \quad (16)$$

where

$$\begin{aligned} U_h^n &= u + \frac{\Delta t}{2} u_t + \frac{\Delta t^2}{3!} u_{tt} + \cdots + \frac{\Delta t^N}{(N+1)!} \frac{\partial^N u}{\partial t^N} \\ &= u - \frac{\Delta t}{2} u_x + \frac{\Delta t^2}{3!} u_{xx} + \cdots + (-1)^N \frac{\Delta t^N}{(N+1)!} \frac{\partial^N u}{\partial x^N}. \end{aligned}$$

Theorem 15. *For the linear advection equation*

$$u_t + u_x = 0,$$

the Lax-Wendroff Flux Reconstruction scheme with D2 dissipation and ADER-FR scheme are equivalent.

Theorem 16. *For the linear advection equation*

$$u_t + u_x = 0,$$

the Lax-Wendroff Flux Reconstruction scheme with D2 dissipation and ADER-FR scheme are equivalent.

Proof. Let $u_e^n = u_e^n(x)$ denote the solution polynomial at time level n in element e .

Theorem 17. *For the linear advection equation*

$$u_t + u_x = 0,$$

the Lax-Wendroff Flux Reconstruction scheme with D2 dissipation and ADER-FR scheme are equivalent.

Proof. Let $u_e^n = u_e^n(x)$ denote the solution polynomial at time level n in element e .

Then, $\tilde{u}_h(x, t) := u_e^n(x - (t - t^n))$ is a weak solution of the equation

$$\begin{aligned}\tilde{u}_t + \tilde{u}_x &= 0, & x \in [x_{e-1/2}, x_{e+1/2}], t \in (t_n, t_{n+1}], \\ \tilde{u}(x, t^n) &= u_e^n(x), & t = t_n.\end{aligned}$$

Theorem 18. *For the linear advection equation*

$$u_t + u_x = 0,$$

the Lax-Wendroff Flux Reconstruction scheme with D2 dissipation and ADER-FR scheme are equivalent.

Proof. Let $u_e^n = u_e^n(x)$ denote the solution polynomial at time level n in element e .

Then, $\tilde{u}_h(x, t) := u_e^n(x - (t - t^n))$ is a weak solution of the equation

$$\begin{aligned}\tilde{u}_t + \tilde{u}_x &= 0, & x \in [x_{e-1/2}, x_{e+1/2}], t \in (t_n, t_{n+1}], \\ \tilde{u}(x, t^n) &= u_e^n(x), & t = t_n.\end{aligned}$$

Since the predictor equation has a **unique** solution of degree N [10, 5], the specified \tilde{u}_h must be **the** predictor solution.

Theorem 19. *For the linear advection equation*

$$u_t + u_x = 0,$$

the Lax-Wendroff Flux Reconstruction scheme with D2 dissipation and ADER-FR scheme are equivalent.

Proof. Let $u_e^n = u_e^n(x)$ denote the solution polynomial at time level n in element e .

Then, $\tilde{u}_h(x, t) := u_e^n(x - (t - t^n))$ is a weak solution of the equation


$$\begin{aligned}\tilde{u}_t + \tilde{u}_x &= 0, & x \in [x_{e-1/2}, x_{e+1/2}], t \in (t_n, t_{n+1}], \\ \tilde{u}(x, t^n) &= u_e^n(x), & t = t_n.\end{aligned}$$

Since the predictor equation has a **unique** solution of degree N [10, 5], the specified \tilde{u}_h must be **the** predictor solution.


$$\begin{aligned}\tilde{u}_h(x, t) &= \tilde{u}_h(x, t^n) + (t - t^n) \frac{\partial}{\partial t} \tilde{u}_h(x, t^n) + \dots + \frac{(t - t^n)^N}{N!} \frac{\partial^N}{\partial t^N} \tilde{u}_h(x, t^n) \\ &= u^n(x) - (t - t^n) \frac{\partial}{\partial x} u^n(x) + \dots + (-1)^N \frac{(t - t^n)^N}{N!} \frac{\partial^N}{\partial x^N} u^n(x) \\ \Rightarrow \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} \tilde{u}_h(x, t) dt &= u^n(x) - \frac{\Delta t}{2} \frac{\partial}{\partial x} u^n(x) + \dots + (-1)^N \frac{\Delta t^N}{(N+1)!} \frac{\partial^N}{\partial x^N} u^n(x) \\ \Rightarrow \frac{1}{\Delta t} \int_{t^n}^{t^{n+1}} \tilde{u}_h(x, t) dt &= U_h^n(x)\end{aligned}$$

Thus, the ADER update (15) and the LWFR-D2 update (16) are the same. \square


$$u_0(x) = e^{-10x^2} \sin(10\pi x), \text{ on periodic } [-1, 1] \text{ with 120 dofs}$$




$N = 1$



$N = 2$







$N = 3$



$N = 4$

$$u_0(x) = e^{-10x^2} \sin(10\pi x), \text{ on periodic } [-1, 1] \text{ with 120 dofs}$$



$N = 3$

$N = 4$

Theorem 20. *If the predictor solution of ADER satisfies*

$$\|U^n\|_{C^{N+1}} \leq C$$

where C is a constant independent of $n, \Delta x, \Delta t$, then the ADER and LW solution will satisfy

$$\|u_{\text{ADER}}^{n+1} - u_{\text{LW}}^{n+1}\|_\infty = O(\Delta t^{N+1}).$$

Theorem 21. *If the predictor solution of ADER satisfies*

$$\|U^n\|_{C^{N+1}} \leq C$$

where C is a constant independent of $n, \Delta x, \Delta t$, then the ADER and LW solution will satisfy

$$\|u_{\text{ADER}}^{n+1} - u_{\text{LW}}^{n+1}\|_\infty = O(\Delta t^{N+1}).$$

Idea of proof. The weak formulation will give us

$$\tilde{u}_t + (\tilde{f}_h)_x = 0, \tag{18}$$

for all (x_r, t_s) where $s > 0$, i.e., $t_s > t^n$. Then, we can extrapolate to $t = t^n$ as

Theorem 22. *If the predictor solution of ADER satisfies*

$$\|U^n\|_{C^{N+1}} \leq C$$

where C is a constant independent of $n, \Delta x, \Delta t$, then the ADER and LW solution will satisfy

$$\|u_{\text{ADER}}^{n+1} - u_{\text{LW}}^{n+1}\|_\infty = O(\Delta t^{N+1}).$$

Idea of proof. The weak formulation will give us

$$\tilde{u}_t + (\tilde{f}_h)_x = 0, \quad (19)$$

for all (x_r, t_s) where $s > 0$, i.e., $t_s > t^n$. Then, we can extrapolate to $t = t^n$ as

$$\tilde{u}_t + (\tilde{f}_h)_x = O(\Delta t^N) \quad \text{at } t = t^n,$$

Theorem 23. *If the predictor solution of ADER satisfies*

$$\|U^n\|_{C^{N+1}} \leq C$$

where C is a constant independent of $n, \Delta x, \Delta t$, then the ADER and LW solution will satisfy

$$\|u_{\text{ADER}}^{n+1} - u_{\text{LW}}^{n+1}\|_\infty = O(\Delta t^{N+1}).$$

Idea of proof. The weak formulation will give us

$$\tilde{u}_t + (\tilde{f}_h)_x = 0, \quad (20)$$

for all (x_r, t_s) where $s > 0$, i.e., $t_s > t^n$. Then, we can extrapolate to $t = t^n$ as

$$\tilde{u}_t + (\tilde{f}_h)_x = O(\Delta t^N) \quad \text{at } t = t^n,$$

so that we will have

$$\tilde{u}_h(x, t^n) = \tilde{u}_h(x, t^n) + \Delta t (\tilde{f}_h)_x + \dots + \frac{\Delta t^N}{N!} \frac{\partial^{N-1}}{\partial t^{N-1}} (\tilde{f}_h)_x + O(\Delta t^{N+1}).$$

- **Modularity** - new conservation law can be added without modifying base code. User need only supply physical flux, numerical flux and wave speed estimates.
- **Portability** - Dependencies are handled by Julia's package manager
- **Parallelization** - Shared-memory via multithreading
- **Efficiency** - noticeably faster than some C++ implementations
- **Visualization** - Postprocessing to vtr format

- **Modularity** - new conservation law can be added without modifying base code. User need only supply physical flux, numerical flux and wave speed estimates.
- **Portability** - Dependencies are handled by Julia's package manager
- **Parallelization** - Shared-memory via multithreading
- **Efficiency** - noticeably faster than some C++ implementations
- **Visualization** - Postprocessing to vtr format

- **Modularity** - new conservation law can be added without modifying base code. User need only supply physical flux, numerical flux and wave speed estimates.
- **Portability** - Dependencies are handled by Julia's package manager
- **Parallelization** - Shared-memory via multithreading
- **Efficiency** - noticeably faster than some C++ implementations
- **Visualization** - Postprocessing to vtr format

- **Modularity** - new conservation law can be added without modifying base code. User need only supply physical flux, numerical flux and wave speed estimates.
- **Portability** - Dependencies are handled by Julia's package manager
- **Parallelization** - Shared-memory via multithreading
- **Efficiency** - noticeably faster than some C++ implementations
- **Visualization** - Postprocessing to vtr format

- **Modularity** - new conservation law can be added without modifying base code. User need only supply physical flux, numerical flux and wave speed estimates.
- **Portability** - Dependencies are handled by Julia's package manager
- **Parallelization** - Shared-memory via multithreading
- **Efficiency** - noticeably faster than some C++ implementations
- **Visualization** - Postprocessing to `vtr` format

```
container = Dict( "u" => u, ... )  
...  
u = container["u"]  
for cell in eachelement(grid)  
    ! heavy computation with u  
end
```

Bad version

```
container = (; u, ...)  
...  
u = container.u  
for cell in eachelement(grid)  
    ! heavy computation with u  
end
```

Good version

```
container = Dict("u" => u, ...)
```

```
...
```

```
u = container["u"]
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

```
container = (; u, ...)
```

```
...
```

```
u = container.u
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

Bad version

Good version

Tool to find type instabilities - `ProfileView.jl`

```
container = Dict("u" => u, ...)
```

```
...
```

```
u = container["u"]
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

```
container = (; u, ...)
```

```
...
```

```
u = container.u
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

Bad version

Good version

Tool to find type instabilities - `ProfileView.jl`

Tool to measure allocations - `BenchmarkTools.jl`, `TimerOutputs.jl`

```
container = Dict("u" => u, ...)
```

```
...
```

```
u = container["u"]
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

```
container = (; u, ...)
```

```
...
```

```
u = container.u
```

```
for cell in eachelement(grid)
```

```
    ! heavy computation with u
```

```
end
```

Bad version

Good version

Tool to find type instabilities - `ProfileView.jl`


Tool to measure allocations - `BenchmarkTools.jl`, `TimerOutputs.jl`

Fixing the problem - JuliaLang - forum, Zulip, Slack.

$$F' = D \times F$$

$$F = \text{Pointwise Action } U$$

$$\begin{array}{|c|c|c|} \hline & \tilde{F}' & F' \\ \hline \end{array} = \begin{array}{|c|} \hline D \\ \hline \end{array} \times \begin{array}{|c|c|c|} \hline & \tilde{F} & F \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline & \tilde{F} & F \\ \hline \end{array} = \text{Pointwise Action} \quad \begin{array}{|c|c|c|} \hline & \tilde{U} & U \\ \hline \end{array}$$


```
for cell in eachelement(grid) ! Cell loop
    for i in eachnode(basis)      ! DoF loop
        f[:,i,cell] = flux(u[:,i,cell])
    end
    BLAS.mul(D, f, res)
end
```

Bad version

```
for cell in eachelement(grid) ! Cell loop
    for i in eachnode(basis)      ! DoF loop
        u_node = get_node_vars(eq, u, i, cell)
        f_node = flux(u_node)
        for ix in eachnode(basis)
            ! res[:,ix,i,cell] += D[ix,i] * f_node
            multiply_add_to_node_vars(eq, D[ix,i],
                                         f_node, res,
                                         iix, cell)
        end
    end
end
```

Good version

Solution points and subcells



Solution points and subcells



Integrate conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ over $I_j^e \times [t^n, t^{n+1}]$

$$\Delta x_e w_j (\mathbf{u}_j^{n+1} - \mathbf{u}_j^n) + \int_{t^n}^{t^{n+1}} (\mathbf{f}_{j+\frac{1}{2}} - \mathbf{f}_{j-\frac{1}{2}}) dt = 0.$$

Solution points and subcells



Integrate conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ over $I_j^e \times [t^n, t^{n+1}]$

$$\Delta x_e w_j (\mathbf{u}_j^{n+1} - \mathbf{u}_j^n) + \int_{t^n}^{t^{n+1}} (\mathbf{f}_{j+\frac{1}{2}} - \mathbf{f}_{j-\frac{1}{2}}) dt = 0.$$

Midpoint rule: $\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_e w_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$

$$\mathbf{f}_{j+1/2}^{n+1/2} = \mathbf{f}(\mathbf{u}_{j+1/2-}^{n+1/2}, \mathbf{u}_{j+1/2+}^{n+1/2})$$

Solution points and subcells



Integrate conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ over $I_j^e \times [t^n, t^{n+1}]$

$$\Delta x_e w_j (\mathbf{u}_j^{n+1} - \mathbf{u}_j^n) + \int_{t^n}^{t^{n+1}} (\mathbf{f}_{j+\frac{1}{2}} - \mathbf{f}_{j-\frac{1}{2}}) dt = 0.$$

Midpoint rule: $\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_e w_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$
 $\mathbf{f}_{j+1/2}^{n+1/2} = \mathbf{f}(\mathbf{u}_{j+1/2-}^{n+1/2}, \mathbf{u}_{j+1/2+}^{n+1/2})$

$$\mathbf{u}_{j-\frac{1}{2}+}^{n+1/2} = \mathbf{u}_{j-\frac{1}{2}+}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}+}) - f(\mathbf{u}_{j-\frac{1}{2}-})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}, \quad \mathbf{u}_{j+\frac{1}{2}-}^{n+1/2} = \mathbf{u}_{j+\frac{1}{2}-}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}-}) - f(\mathbf{u}_{j-\frac{1}{2}+})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}.$$

Solution points and subcells



Integrate conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ over $I_j^e \times [t^n, t^{n+1}]$

$$\Delta x_e w_j (\mathbf{u}_j^{n+1} - \mathbf{u}_j^n) + \int_{t^n}^{t^{n+1}} (\mathbf{f}_{j+\frac{1}{2}} - \mathbf{f}_{j-\frac{1}{2}}) dt = 0.$$

Midpoint rule: $\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_e w_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$
 $\mathbf{f}_{j+1/2}^{n+1/2} = \mathbf{f}(\mathbf{u}_{j+1/2-}^{n+1/2}, \mathbf{u}_{j+1/2+}^{n+1/2})$

$$\mathbf{u}_{j-\frac{1}{2}+}^{n+1/2} = \mathbf{u}_{j-\frac{1}{2}+}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}+}) - f(\mathbf{u}_{j-\frac{1}{2}-})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}, \quad \mathbf{u}_{j+\frac{1}{2}-}^{n+1/2} = \mathbf{u}_{j+\frac{1}{2}-}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}-}) - f(\mathbf{u}_{j-\frac{1}{2}+})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}.$$

$$\begin{aligned} \mathbf{u}_{j-\frac{1}{2}+} &= \mathbf{u}_j(x_{j-\frac{1}{2}}), & \mathbf{u}_{j+\frac{1}{2}-} &= \mathbf{u}_j(x_{j+\frac{1}{2}}) \\ \mathbf{u}_j(x) &= \mathbf{u}_j^n + \boldsymbol{\sigma}_j (x - x_j) \end{aligned}$$

Solution points and subcells



Integrate conservation law $\mathbf{u}_t + \mathbf{f}(\mathbf{u})_x = \mathbf{0}$ over $I_j^e \times [t^n, t^{n+1}]$

$$\Delta x_e w_j (\mathbf{u}_j^{n+1} - \mathbf{u}_j^n) + \int_{t^n}^{t^{n+1}} (\mathbf{f}_{j+\frac{1}{2}} - \mathbf{f}_{j-\frac{1}{2}}) dt = 0.$$

Midpoint rule: $\mathbf{u}_j^{n+1} = \mathbf{u}_j^n - \frac{\Delta t}{\Delta x_e w_j} (\mathbf{f}_{j+1/2}^{n+1/2} - \mathbf{f}_{j-1/2}^{n+1/2})$
 $\mathbf{f}_{j+1/2}^{n+1/2} = \mathbf{f}(\mathbf{u}_{j+1/2-}^{n+1/2}, \mathbf{u}_{j+1/2+}^{n+1/2})$

$$\mathbf{u}_{j-\frac{1}{2}+}^{n+1/2} = \mathbf{u}_{j-\frac{1}{2}+}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}+}) - f(\mathbf{u}_{j-\frac{1}{2}-})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}, \quad \mathbf{u}_{j+\frac{1}{2}-}^{n+1/2} = \mathbf{u}_{j+\frac{1}{2}-}^n - \frac{\Delta t}{2} \frac{f(\mathbf{u}_{j+\frac{1}{2}-}) - f(\mathbf{u}_{j-\frac{1}{2}+})}{x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}}.$$

$$\begin{aligned} \mathbf{u}_{j-\frac{1}{2}+} &= \mathbf{u}_j(x_{j-\frac{1}{2}}), & \mathbf{u}_{j+\frac{1}{2}-} &= \mathbf{u}_j(x_{j+\frac{1}{2}}) \\ \mathbf{u}_j(x) &= \mathbf{u}_j^n + \boldsymbol{\sigma}_j (x - x_j) \end{aligned}$$

$$\begin{aligned} \boldsymbol{\sigma}_j &= \text{minmod} \left(\beta_e \frac{\mathbf{u}_{j+1} - \mathbf{u}_j}{x_{j+1} - x_j}, D_{\text{cent}}(\mathbf{u})_j, \beta_e \frac{\mathbf{u}_j^n - \mathbf{u}_{j-1}^n}{x_j - x_{j-1}} \right) \\ \beta_e &= 2 - \alpha_e \end{aligned}$$

$$\begin{aligned}
& \frac{d\mathbf{u}_{e,\mathbf{p}}^\delta}{dt} + \frac{1}{J} \nabla_{\boldsymbol{\xi}}^N \cdot \tilde{\mathbf{f}}_e(\boldsymbol{\xi}_{\mathbf{p}}) \\
& + \frac{1}{J} \sum_{i=1}^3 ((\tilde{\mathbf{f}}_e \cdot \mathbf{n}_{s,i})^* - \tilde{\mathbf{f}}_e \cdot \mathbf{n}_{s,i})(\boldsymbol{\xi}_i^R) g'_R(\xi_{p_i}) \\
& + \frac{1}{J} \sum_{i=1}^3 ((\tilde{\mathbf{f}}_e \cdot \mathbf{n}_{s,i})^* - \tilde{\mathbf{f}}_e \cdot \mathbf{n}_{s,i})(\boldsymbol{\xi}_i^L) g'_L(\xi_{p_i}) = \mathbf{0},
\end{aligned}$$

where

$$\mathbf{p} = (p_1, p_2, p_3) \text{ for } 1 \leq p_i \leq N + 1$$

$\boldsymbol{\xi}_{\mathbf{p}} = \boldsymbol{\xi}_{p_1 p_2 p_3} = (\xi_{p_1}, \xi_{p_2}, \xi_{p_3})$ is the solution point of collocation, e is the element index

$\nabla_{\boldsymbol{\xi}}^N$ is the degree N gradient

$(\tilde{\mathbf{f}}_e \cdot \mathbf{n})^*$ is the numerical flux along normal \mathbf{n}

$\mathbf{n}_{s,i}$ is the normal vector in reference cell along i^{th} direction

$\boldsymbol{\xi}_i^{L/R}$ is the point that agrees with $\boldsymbol{\xi}$ at j, k and equals 0 or 1 for L or R at i .