



[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Farmers' Innovation](#) | [Publications](#) | [Contact](#)

**Agricultural crops :: Cash crops :: Cotton**

## Tobacco Streak Virus

### Symptom

- Initially chlorotic yellow spot appear on the infected leaves
- Later spot turn into distinct necrotic purple spot with yellow halo.
- Necrotic streaks are produced on the petiole.
- Bud and flower productions are reduced.
- Severe infection resulted in drying of squares.

### Survival and Mode of Spread

- Virus survives in alternate hosts.
- Primary spread is From infected plants.
- The virus is transmitted by Thrips (Thrips tabaci).

### Favourable Conditions

- Minimum temperature, High relative humidity with leaf wetness.

### Management

- Remove infected plants and other host plants.
- Periodical weeding should be done.
- Seed treatment with imidacloprid 70WS at 7g / kg .
- Spray Dinotefuran 20% SG 60 g/acre or Flonicamid 50% WG @ 60 g/acre or Imidacloprid 17.8%SL 40- 50 ml/acre for controlling thrips

### Source of information

- Online resources
- CPG 2020



Updated on Apr 2023

[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Publications](#) | [Disclaimer](#) | [Contact Us](#)

TNAU. All Rights Reserved © 2008 - 2023 .