

PAPER • OPEN ACCESS

Long-time convergence of a nonlocal Burgers' equation towards the local N-wave

To cite this article: Giuseppe Maria Coclite *et al* 2023 *Nonlinearity* **36** 5998

View the [article online](#) for updates and enhancements.

You may also like

- [Solution ALD Processing of Halide Perovskite Thin Films Yields Superior Functional Quality and Stability Than Classical Processing](#)
Vanessa Koch, Maissa K. S. Barr, Pascal Büttner et al.
- [Coverage- and temperature-induced self-metalation of tetraphenyltransdibenzoporphyrin on Cu\(111\)](#)
Majid Shaker, Maximilian Muth, Julien Steffen et al.
- [Corrigendum: Removing the orientational degeneracy of the TS defect in 4H-SiC by electric fields and strain \(2021 New J. Phys. 23 073002\)](#)
Fabio Candolfi, Johannes A F Lehmeyer, Maximilian Rühl et al.

Long-time convergence of a nonlocal Burgers' equation towards the local N-wave

Giuseppe Maria Coclite¹ , Nicola De Nitti^{2,*} ,
Alexander Keimer³ , Lukas Pflug^{3,4} 
and Enrique Zuazua^{2,5,6} 

¹ Department of Mechanics, Mathematics, and Management, Polytechnic University of Bari, Via E. Orabona 4, 70125 Bari, Italy

² Department of Mathematics, Chair for Dynamics, Control, Machine Learning and Numerics (Alexander von Humboldt Professorship), Friedrich-Alexander-Universität Erlangen-Nürnberg, Cauerstr. 11, 91058 Erlangen, Germany

³ Department of Mathematics, Chair of Applied Mathematics (Continuous Optimization), Friedrich-Alexander-Universität Erlangen-Nürnberg, Cauerstr. 11, 91058 Erlangen, Germany

⁴ Friedrich-Alexander-Universität Erlangen-Nürnberg, Central Institute for Scientific Computing, Martensstr. 5a, 91058 Erlangen, Germany

⁵ Chair of Computational Mathematics, Fundación Deusto, Avenida de las Universidades, 24, 48007 Bilbao, Basque Country, Spain

⁶ Departamento de Matemáticas, Ciudad Universitaria de Cantoblanco, Universidad Autónoma de Madrid, 28049 Madrid, Spain

E-mail: nicola.de.nitti@fau.de

Received 1 March 2023; revised 28 June 2023

Accepted for publication 14 August 2023

Published 4 October 2023

Recommended by Dr Theodore Dimitrios Drivas



Abstract

We study the long-time behaviour of the unique weak solution of a nonlocal regularisation of the (inviscid) Burgers equation where the velocity is approximated by a one-sided convolution with an exponential kernel. The initial datum is assumed to be positive, bounded, and integrable. The asymptotic profile is given by the ‘N-wave’ entropy solution of the Burgers equation. The key ingredients of the proof are a suitable scaling argument and a nonlocal Oleinik-type estimate.

* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Keywords: nonlocal conservation laws, nonlocal flux, Burgers equation, approximation of local conservation laws, N-waves, source-type solutions, entropy solutions

Mathematics Subject Classification numbers: 35L65

(Some figures may appear in colour only in the online journal)

1. Introduction

Let us consider the following nonlocal regularisation of the Burgers equation:

$$\begin{cases} \partial_t \rho(t, x) + \partial_x (W[\rho](t, x) \rho(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ \rho(0, x) = \rho_0(x), & x \in \mathbb{R}, \end{cases} \quad (1.1)$$

supplemented by the nonlocal term

$$W[\rho](t, x) := \int_{-\infty}^x \exp(y - x) \rho(t, y) dy, \quad (t, x) \in (0, +\infty) \times \mathbb{R}, \quad (1.2)$$

which also satisfies the identity

$$\partial_x W[\rho](t, x) = \rho(t, x) - W[\rho](t, x), \quad (t, x) \in (0, +\infty) \times \mathbb{R}. \quad (1.3)$$

In what follows, we assume that the initial datum satisfies

$$\rho_0 \in L^1(\mathbb{R}; \mathbb{R}_{\geq 0}) \cap L^\infty(\mathbb{R}; \mathbb{R}_{\geq 0}) \quad (1.4)$$

and introduce the notation $M := \int_{\mathbb{R}} \rho_0(x) dx$ for its L^1 -mass.

Under these assumptions, the nonlocal conservation law (1.1) has a unique global non-negative weak solution ρ . In particular, in contrast to the case of the local Burgers equation, no entropy condition is required to select a unique weak solution; moreover, the regularity and integrability of the initial datum is essentially preserved along the evolution (see [29] and section 2 below). The aim of this paper is to study its asymptotic behaviour when $t \rightarrow +\infty$. The main result asserts that, as $t \rightarrow +\infty$, the solution $\rho(t, \cdot)$ of (1.1) converges to the (unique) *N-wave solution* (or *source-type solution*) w of the local Burgers equation (see [37, equation (2.1)]), i.e. the solution of the Burgers equation with a Dirac delta as initial data,

$$\begin{cases} \partial_t w(t, x) + \partial_x (w^2(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ w(0, \cdot) = M\delta_0, & \text{in } \mathbb{R}, \end{cases} \quad (1.5)$$

which is given explicitly by

$$w(t, x) = \begin{cases} \frac{x}{2t} & \text{if } x \in (0, \sqrt{4Mt}), \\ 0 & \text{otherwise;} \end{cases} \quad (1.6)$$

see figure 1. We refer to [37] (and to section 2 below) for the proof that (1.5) does indeed have a unique entropy solution (which is given by (1.6)) under suitable assumptions.

More precisely, our main theorem on the long-time behaviour of the solution of (1.1) can be stated as follows.




Figure 1. Plot of an N -wave solution (1.6) (with $M = 1$) for $t = 0.5$ (blue), $t = 1$ (red), and $t = 2$ (yellow).

Theorem 1.1 (long-time asymptotics). *Given ρ_0 satisfying assumption (1.4), let ρ be the unique weak solution of the nonlocal Burgers equation (1.1) and W be the corresponding nonlocal term. Then, for $p \in [1, +\infty)$, we have*

$$t^{\frac{1}{2}(1-\frac{1}{p})} \|\rho(t, \cdot) - w(t, \cdot)\|_{L^p(\mathbb{R})} \rightarrow 0 \quad \text{and} \quad t^{\frac{1}{2}(1-\frac{1}{p})} \|W(t, \cdot) - w(t, \cdot)\|_{L^p(\mathbb{R})} \rightarrow 0 \quad \text{as } t \rightarrow +\infty, \quad (1.7)$$

where w denotes the unique entropy solution (N -wave solution) of the Burgers equation (1.5) defined in (1.6).

For local conservation laws, the long-time convergence of the entropy solution to the corresponding N -wave profile, as well as existence and uniqueness of entropy solutions in case of measure initial data, was first established rigorously in [37]. In [17–20, 32, 33], the analysis was extended to classes of viscous conservation laws with flux $f(\xi) = \xi^q$ (for $\xi \in \mathbb{R}$). In case $1 < q < 2$, as $t \rightarrow +\infty$, the solutions converge to the N -wave profile of the inviscid conservation law; on the other hand, if $q = 2$, the limit profile is given by the fundamental solution of the viscous conservation law and, if $q > 2$, it is of Gaussian type (i.e. the fundamental solution of the heat equation with mass $\int_{\mathbb{R}} u_0 \, dx$). Similar results have been also obtained for scalar conservation laws with fractional diffusion (see [3, 16, 28]). For the multi-dimensional setting, we also refer to the recent work [38], where it was proved that a multidimensional Burgers-type equation with a Dirac delta distribution as initial data is not well-posed (despite the L^1 – L^∞ smoothing effect established in [39]).

For nonlocal conservation laws, this problem has not been considered in the literature. However, interestingly, it can be reduced to a type of nonlocal-to-local singular limit problem that has attracted much attention in recent years. Indeed, following [37], given $\lambda > 0$, we consider the rescaled function

$$\rho_\lambda(t, x) := \lambda \rho(\lambda^2 t, \lambda x), \quad (1.8)$$

which solves⁷

$$\begin{cases} \partial_t \rho_\lambda(t, x) + \partial_x (W_\lambda[\rho_\lambda](t, x) \rho_\lambda(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ \rho_\lambda(0, x) = \rho_{0, \lambda}(x) := \lambda \rho_0(\lambda x), & x \in \mathbb{R}, \end{cases} \quad (1.9)$$

with

$$W_\lambda[\rho_\lambda](t, x) := \lambda \int_{-\infty}^x \exp(\lambda(y - x)) \rho_\lambda(t, y) dy, \quad (t, x) \in (0, +\infty) \times \mathbb{R}, \quad (1.10)$$

which satisfies

$$\lambda^{-1} \partial_x W_\lambda[\rho_\lambda](t, x) = \rho_\lambda(t, x) - W_\lambda[\rho_\lambda](t, x), \quad (t, x) \in [0, +\infty) \times \mathbb{R}. \quad (1.11)$$

We shall prove that, for a fixed $t > 0$, $\rho_\lambda(t, \cdot) \rightarrow w(t, \cdot)$ in $L^1(\mathbb{R})$ as $\lambda \rightarrow \infty$, which, in turn, will be shown to yield

$$\|\rho(t, \cdot) - w(t, \cdot)\|_{L^1(\mathbb{R})} \rightarrow 0 \quad \text{as } t \rightarrow +\infty$$

with w defined in (1.6) (and, by interpolation, the claim in theorem 1.1).

This type of singular limit problem has been intensively studied in the case of initial data that are uniformly bounded with respect to the scaling parameter. First, in [2], it has been observed that, at least numerically, there is some hope that the solution of a nonlocal conservation law converges to the entropy solution of the corresponding local problem when the nonlocal term approaches a Dirac delta. Positive results in this direction were obtained in [30] for a large class of nonlocal conservation laws under the assumption of having monotone initial data; in [13] under the assumption that the initial datum has bounded total variation, is bounded away from zero, and satisfies a one-sided Lipschitz condition. For the case of an exponential weight, in [7, 8], Bressan and Shen proved a convergence result under the assumption that the initial datum is bounded away from zero and has bounded total variation. The core of their argument is the observation that, under suitable changes of variables, the nonlocal problem can be rewritten as a hyperbolic system with relaxation terms. The assumption on the initial data being bounded away from zero played a key role in showing a uniform total variation bound for the solution of the nonlocal problem. Indeed, in [13], a counterexample shows that the total variation of the solution may blow up if the data is not bounded away from zero.

On the other hand, in [11], by arguing on the nonlocal term W rather than on the solution of the conservation law, it was possible to remove the additional assumption on the initial data—the key observation being that the nonlocal term W enjoys further regularity and, in particular, its total variation remains uniformly bounded. From the compactness of the sequence of non-local terms, it is then possible to deduce the convergence for the sequence of solutions as well. This approach was later adapted in [14] to classes of weights more general than the exponential one.

⁷ Note that

$$W[\rho](\lambda^2 t, \lambda x) = \int_{-\infty}^{\lambda x} \exp(y - \lambda x) \rho(\lambda^2 t, y) dy = \int_{-\infty}^x \exp(\lambda(z - x)) \lambda \rho(\lambda^2 t, \lambda z) dz = \frac{1}{\lambda} W_\lambda[\rho_\lambda](t, x).$$

Then

$$\begin{aligned} \partial_t \rho(t, x) + \partial_x (W[\rho](t, x) \rho(t, x)) = 0 &\iff \partial_t \rho(\lambda^2 t, \lambda x) + \lambda \partial_x (W[\rho](\lambda^2 t, \lambda x) \rho(\lambda^2 t, \lambda x)) = 0 \\ &\iff \partial_t \lambda \rho(\lambda^2 t, \lambda x) + \lambda \partial_x (W_\lambda[\rho_\lambda](t, x) \rho(\lambda^2 t, \lambda x)) = 0 \\ &\iff \partial_t \rho_\lambda(t, x) + \partial_x (W_\lambda[\rho_\lambda](t, x) \rho_\lambda(t, x)) = 0. \end{aligned}$$

The difference and substantial added difficulty of the present contribution compared to the above-mentioned works is that, under the scaling transformation, we are considering initial data that concentrate to a Dirac delta: i.e.

$$\rho_\lambda(0, \cdot) \rightarrow M\delta_0 \quad \text{and} \quad W_\lambda[\rho_\lambda](0, \cdot) \rightarrow M\delta_0 \quad (1.12)$$

in the sense of distributions as $\lambda \rightarrow +\infty$. That is, with respect to λ , the only uniform bound for the initial data ρ_0 is given in terms the L^1 -mass.

To overcome this difficulty, we take advantage of an Oleinik-type inequality satisfied by the nonlocal term $W_\lambda[\rho_\lambda]$. Indeed, from [11], it is known that we can rewrite (1.9) as a conservation law with nonlocal source formulated purely in $W_\lambda[\rho_\lambda]$ (see (2.2) below). This motivates using the notation W_λ instead of $W_\lambda[\rho_\lambda]$ in what follows. From (2.2), arguing as in [10], we can deduce the Oleinik-type estimate (see theorem 3.2):

$$\frac{W_\lambda(t, x) - W_\lambda(t, y)}{x - y} \leq \frac{1}{t}, \quad t > 0, \quad x, y \in \mathbb{R}, \quad x \neq y.$$

Combining it with the uniform L^1 -bound

$$\int_{\mathbb{R}} W_\lambda(t, x) dx = M, \quad t > 0,$$

this inequality yields an L^∞ -bound for $t > 0$ (see lemma 3.3):

$$0 \leq W_\lambda(t, x) \lesssim \sqrt{Mt^{-1}}, \quad (t, x) \in (0, +\infty) \times \mathbb{R}.$$

With these ingredients, the approach of [17] leads to the claimed convergence of $\{W_\lambda\}_{\lambda>0}$ towards the N -wave solution of the (local) Burgers equation and, thanks to (1.3), to the convergence of $\{\rho_\lambda\}_{\lambda>0}$ as well.

The paper is organised as follows. In section 2, we recall the necessary preliminaries on the well-posedness of (1.9) (for fixed $\lambda > 0$). In section 3, we prove the key and *a priori* estimates on W_λ sketched above. Then, in section 4, we combine them and establish the convergence of $\{\rho_\lambda\}_{\lambda>0}$ and $\{W_\lambda\}_{\lambda>0}$ to the N -wave solution of the local Burgers equation as $\lambda \rightarrow +\infty$; or, equivalently, of $\{\rho(t, \cdot)\}_{t>0}$ and $\{W(t, \cdot)\}_{t>0}$ as $t \rightarrow +\infty$. This convergence result is illustrated by several numerical simulations in section 5 (together with some further conjectures). Finally, in section 6, we conclude the paper by presenting some open problems.

2. Preliminaries

For the nonlocal conservation law in (1.9), we recall the following well-posedness result and some fundamental properties of the solution. We refer to [11, theorem 2.1 and lemma 3.1] (which, in turn, relies in part on [29, theorems 2.20, 3.2 and corollary 4.3] or [12, theorem 2.1 and corollary 2.1]), [23, theorem 2.1], [14, proposition 2.1 and corollary 2.2], or [10] for the proof of a similar statement.

Theorem 2.1 (existence and uniqueness of weak solutions and maximum principle). *Let us suppose that the initial datum ρ_0 satisfies (1.4). Then, for every $\lambda > 0$, there exists a unique weak solution $\rho_\lambda \in C([0, +\infty); L^1(\mathbb{R})) \cap L^\infty((0, +\infty); L^\infty(\mathbb{R}))$ of the nonlocal Burgers equation (1.9) and the following maximum principle holds:*

$$\operatorname{ess\,inf}_{x \in \mathbb{R}} \rho_{0,\lambda}(x) \leq \rho_\lambda(t, x) \leq \|\rho_{0,\lambda}\|_{L^\infty(\mathbb{R})}, \quad \text{a.e. } (t, x) \in (0, +\infty) \times \mathbb{R}. \quad (2.1)$$

Moreover, for the nonlocal term W_λ , the following properties hold:

- (1) $W_\lambda \in W^{1,\infty}((0, +\infty) \times \mathbb{R})$ and $\operatorname{ess\,inf}_{x \in \mathbb{R}} \rho_{0,\lambda}(x) \leq W_\lambda \leq \|\rho_{0,\lambda}\|_{L^\infty(\mathbb{R})}$;
- (2) $W_\lambda \in C^0((0, +\infty); L^1(\mathbb{R}))$; in particular, if $\|\rho_\lambda(t, \cdot)\|_{L^1(\mathbb{R})} = M$, then $\|W_\lambda(t, \cdot)\|_{L^1(\mathbb{R})} = M$;
- (3) if $\rho_{0,\lambda} \in C^k(\mathbb{R})$, then $W_\lambda \in C^{k+1}((0, +\infty) \times \mathbb{R})$ for $k \geq 0$.

Furthermore, W_λ satisfies the following conservation law with nonlocal source in the strong sense:

$$\begin{cases} \partial_t W_\lambda(t, x) + \partial_x (W_\lambda^2(t, x)) \\ = \lambda \int_{-\infty}^x \exp(\lambda(y-x)) (W_\lambda(t, x) \partial_x W_\lambda(t, x) - W_\lambda(t, y) \partial_y W_\lambda(t, y)) dy, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ W_\lambda(0, x) = \lambda \int_{-\infty}^x \exp(\lambda(y-x)) \rho_{0,\lambda}(y) dy, & x \in \mathbb{R}. \end{cases} \quad (2.2)$$

For the limit problem (1.5), we rely on a more general well-posedness result from [37, theorem 1.1 and remark 1.1].

Theorem 2.2 (non-negative entropy solutions with measure initial data). *Let us consider the local conservation law*

$$\begin{cases} \partial_t u(t, x) + \partial_x f(u(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ u(0, \cdot) = \mu, & \text{in } \mathbb{R}. \end{cases} \quad (2.3)$$

Let us assume that $f: \mathbb{R} \rightarrow \mathbb{R}$ is locally Lipschitz continuous with $f(0) = 0$ and $f([0, \infty)) \subset [0, \infty)$ and that μ is a non-negative finite measure on $\mathbb{R}$. Then there exists at most one non-negative solution $u \in C((0, +\infty); L^1(\mathbb{R})) \cap L^\infty((\tau, +\infty) \times \mathbb{R})$, for all $\tau \in (0, +\infty)$, which satisfies the Kruřkov entropy condition, i.e.

$$\begin{aligned} & \forall k \in \mathbb{R}, \forall \varphi \in C_c^\infty((0, +\infty) \times \mathbb{R}), \quad \varphi \geq 0: \\ & \int_0^{+\infty} \int_{\mathbb{R}} |u - k| \partial_t \varphi + \operatorname{sign}(u - k) (f(u) - f(k)) \partial_x \varphi \, dx \, dt \geq 0, \end{aligned}$$

and achieves the initial datum in the narrow (or weak) sense of measures⁸,

$$\lim_{t \rightarrow 0} u(t, \cdot) = \mu \quad \text{narrowly in } \mathbb{R}.$$

In particular, in our setting, theorem 2.2 yields the uniqueness of the N -wave entropy solution (1.6) of (1.5).

Remark 2.3 (non-negativity condition and uniqueness). As noted in [37, remark 1.2], the uniqueness result in theorem 2.2 fails without the non-negativity assumption. This hypothesis can, however, be replaced by taking $f(\xi) = \operatorname{sign}(\xi)|\xi|^q$ (with $q > 1$) or by $f(\xi) = |\xi|^q$ (with $q > 1$) and assuming that the initial datum is achieved in a stronger sense (as shown in [37, theorem 1.2] and [37, theorem 1.3] respectively).

⁸ A sequence of signed Radon measures $\{\mu_n\}_{n \in \mathbb{N}}$ on $\mathbb{R}$ converges narrowly (or in the weak sense) to μ if, for all bounded and continuous test functions $\varphi \in C_b(\mathbb{R})$, we have

$$\lim_{n \rightarrow +\infty} \int_{\mathbb{R}} \varphi \, d\mu_n = \int_{\mathbb{R}} \varphi \, d\mu.$$

See [4, chapter 8].

3. A priori estimates

Before presenting our key *a priori* estimates, let us recall the following stability result of the nonlocal conservation law (1.1) with respect to the initial datum (see [10]).

Lemma 3.1 (stability of the nonlocal term with respect to the initial datum). *Given $\rho_{0,1}, \rho_{0,2} \in L^1(\mathbb{R})$, let us denote by $W_1, W_2 \in L^\infty((0, T); W^{1,\infty}(\mathbb{R}))$ the nonlocal terms associated to the corresponding solutions of (1.9). Then, the following stability result holds: for all $t \in [0, T]$,*

$$\begin{aligned} & \|W_1(t, \cdot) - W_2(t, \cdot)\|_{L^\infty(\mathbb{R})} \\ & \leq C(\lambda, \|\rho_{0,1}\|_{L^\infty(\mathbb{R})}, \|\rho_{0,2}\|_{L^\infty(\mathbb{R})}, \|\rho_{0,1}\|_{L^1(\mathbb{R})}, \|\rho_{0,2}\|_{L^1(\mathbb{R})}) \|\rho_{0,1} - \rho_{0,2}\|_{L^1(\mathbb{R})}, \end{aligned}$$

where C is a suitable constant that depends only on the quantities mentioned above.

Proof. From the results in [12, 29], we know that the solution of (1.9) can be written as

$$\rho_1(t, x) = \rho_{0,1}(\xi_{W_1}(t, x; 0)) \partial_2 \xi_{W_1}(t, x; 0) \quad \text{and} \quad \rho_2(t, x) = \rho_{0,2}(\xi_{W_2}(t, x; 0)) \partial_2 \xi_{W_2}(t, x; 0),$$

where ξ_{W_1} and ξ_{W_2} solve the characteristic ODEs

$$\begin{aligned} \xi_{W_1}(t, x; \tau) &= x + \int_t^\tau W_1(s, \xi_{W_1}(t, x; s)) ds, \quad \tau \in [0, T], \\ \xi_{W_2}(t, x; \tau) &= x + \int_t^\tau W_2(s, \xi_{W_2}(t, x; s)) ds, \quad \tau \in [0, T]; \end{aligned} \tag{3.1}$$

(here written as Volterra-type integral equations). In particular, we recall that the nonlocal terms corresponding to the initial data $\rho_{0,1}$ and $\rho_{0,2}$ satisfy the following fixed-point equations for $(t, x) \in (0, T) \times \mathbb{R}$:

$$\begin{aligned} W_1(t, x) &= \lambda \int_{-\infty}^x \exp(\lambda(y - x)) \rho_1(t, y) dy \\ &= \lambda \int_{-\infty}^x \exp(\lambda(y - x)) \rho_{0,1}(\xi_{W_1}(t, y; 0)) \partial_2 \xi_{W_1}(t, y; 0) dy \\ &= \lambda \int_{-\infty}^{\xi_{W_1}(t, x; 0)} \exp(\lambda(\xi_{W_1}(0, z; t) - x)) \rho_{0,1}(z) dz; \\ W_2(t, x) &= \lambda \int_{-\infty}^{\xi_{W_2}(t, x; 0)} \exp(\lambda(\xi_{W_2}(0, z; t) - x)) \rho_{0,2}(z) dz. \end{aligned}$$

Taking the absolute value of the difference, we have

$$\begin{aligned} & \lambda^{-1} |W_1(t, x) - W_2(t, x)| \\ &= \left| \int_{-\infty}^{\xi_{W_1}(t, x; 0)} \exp(\lambda(\xi_{W_1}(0, z; t) - x)) \rho_{0,1}(z) dz - \int_{-\infty}^{\xi_{W_2}(t, x; 0)} \exp(\lambda(\xi_{W_2}(0, z; t) - x)) \rho_{0,2}(z) dz \right| \\ &\leq \int_{\min\{\xi_{W_1}(t, x; 0), \xi_{W_2}(t, x; 0)\}}^{\max\{\xi_{W_1}(t, x; 0), \xi_{W_2}(t, x; 0)\}} (|\rho_{0,1}(y)| + |\rho_{0,2}(y)|) dy \\ &\quad + \int_{-\infty}^{\min\{\xi_{W_1}(t, x; 0), \xi_{W_2}(t, x; 0)\}} (\exp(\lambda(\xi_{W_1}(0, z; t) - x)) \rho_{0,1}(z) - \exp(\lambda(\xi_{W_2}(0, z; t) - x)) \rho_{0,2}(z)) dz \\ &\leq |\xi_{W_1}(t, x; 0) - \xi_{W_2}(t, x; 0)| (\|\rho_{0,1}\|_{L^\infty(\mathbb{R})} + \|\rho_{0,2}\|_{L^\infty(\mathbb{R})}) \\ &\quad + \lambda \|\xi_{W_1}(0, \cdot; t) - \xi_{W_2}(0, \cdot; t)\|_{L^\infty(\mathbb{R})} (\|\rho_{0,1}\|_{L^1(\mathbb{R})} + \|\rho_{0,2}\|_{L^1(\mathbb{R})}) + \|\rho_{0,1} - \rho_{0,2}\|_{L^1(\mathbb{R})}. \end{aligned} \tag{3.2}$$

To conclude, we need to study the stability of the characteristics with respect to W_1 and W_2 . For $(t, x, \tau) \in (0, T) \times \mathbb{R} \times (0, T)$, we compute

$$\begin{aligned} |\xi_{W_1}(t, x; \tau) - \xi_{W_2}(t, x; \tau)| &= \left| \int_t^\tau W_1(s, \xi_{W_1}(t, x; s)) - W_2(s, \xi_{W_2}(t, x; s)) \, ds \right| \\ &= \left| \int_t^\tau W_1(s, \xi_{W_1}(t, x; s)) - W_2(s, \xi_{W_1}(t, x; s)) \, ds \right| \\ &\quad + \left| \int_t^\tau W_2(s, \xi_{W_1}(t, x; s)) - W_2(s, \xi_{W_2}(t, x; s)) \, ds \right| \\ &\leq \int_{\min\{t, \tau\}}^{\max\{t, \tau\}} \|W_1(s, \cdot) - W_2(s, \cdot)\|_{L^\infty(\mathbb{R})} \, ds \\ &\quad + \|\partial_x W_2\|_{L^\infty((0, T); L^\infty(\mathbb{R}))} \int_{\min\{t, \tau\}}^{\max\{t, \tau\}} \|\xi_{W_1}(t, \cdot; s) - \xi_{W_2}(t, \cdot; s)\|_{L^\infty(\mathbb{R})} \, ds. \end{aligned}$$

Gronwall's inequality yields

$$\begin{aligned} &\|\xi_{W_1}(t, \cdot; \tau) - \xi_{W_2}(t, \cdot; \tau)\|_{L^\infty(\mathbb{R})} \\ &\leq \int_{\min\{t, \tau\}}^{\max\{t, \tau\}} \|W_1(s, \cdot) - W_2(s, \cdot)\|_{L^\infty(\mathbb{R})} \, ds \exp(|t - \tau| \|\partial_x W_2\|_{L^\infty((0, T); L^\infty(\mathbb{R}))}). \end{aligned}$$

Plugging this into (3.2), we get

$$\begin{aligned} &\|W_1(t, \cdot) - W_2(t, \cdot)\|_{L^\infty(\mathbb{R})} \\ &\leq \lambda (\|\rho_{0,1}\|_{L^\infty(\mathbb{R})} + \|\rho_{0,2}\|_{L^\infty(\mathbb{R})}) \int_{\min\{t, \tau\}}^{\max\{t, \tau\}} \|W_1(s, \cdot) - W_2(s, \cdot)\|_{L^\infty(\mathbb{R})} \, ds \\ &\quad \times \exp(|t - \tau| \|\partial_x W_2\|_{L^\infty((0, T); L^\infty(\mathbb{R}))}) \\ &\quad + \lambda^2 (\|\rho_{0,1}\|_{L^1(\mathbb{R})} + \|\rho_{0,2}\|_{L^1(\mathbb{R})}) \int_t^\tau \|W_1(s, \cdot) - W_2(s, \cdot)\|_{L^\infty(\mathbb{R})} \, ds \\ &\quad \times \exp(|t - \tau| \|\partial_x W_2\|_{L^\infty((0, T); L^\infty(\mathbb{R}))}) + \|\rho_{0,1} - \rho_{0,2}\|_{L^1(\mathbb{R})}. \end{aligned}$$

Applying again Gronwall's inequality on $W_1 - W_2$ and recalling that

$$\partial_x W_2 = \lambda(\rho_2 - W_2) \implies \|\partial_x W_2\|_{L^\infty((0, T); L^\infty(\mathbb{R}))} \leq 2\lambda \|\rho_{0,2}\|_{L^\infty(\mathbb{R})} \quad (3.3)$$

(thanks to the maximum principle in theorem 2.1), we conclude the proof. $\square$

As a first step, we prove an Oleinik-type inequality on the nonlocal term W_λ . The result is essentially contained in [10] (in a more general form). We present the proof below for the sake of completeness.

Theorem 3.2 (Oleinik-type inequality for W_λ). *Given ρ_0 such that (1.4) holds, the solution W_λ of (2.2) satisfies*

$$\frac{W_\lambda(t, x) - W_\lambda(t, y)}{x - y} \leq \frac{1}{t}, \quad t > 0, \quad x, y \in \mathbb{R}, \quad x \neq y, \quad (3.4)$$

for all $\lambda > 0$.

Proof of theorem 3.2. We consider a smoothed initial datum $\rho_{0,\lambda}^\varepsilon$ (for $\varepsilon > 0$) and call the corresponding smooth nonlocal term W_λ^ε . We then compute, differentiating the PDE in (2.2) with respect to x ,

$$\partial_{tx}^2 W_\lambda^\varepsilon = -W_\lambda^\varepsilon \partial_{xx}^2 W_\lambda^\varepsilon - (\partial_x W_\lambda^\varepsilon)^2 - \lambda W_\lambda^\varepsilon \partial_x W_\lambda^\varepsilon + \lambda^2 \int_{-\infty}^x \exp(\lambda(y-x)) W_\lambda^\varepsilon(t,y) \partial_y W_\lambda^\varepsilon(t,y) dy. \quad (3.5)$$

For $t > 0$ fixed, considering $m(t) = \sup_{y \in \mathbb{R}} \partial_y W_\lambda^\varepsilon(t,y)$ and assuming—without loss of generality—that $m(t) \geq 0$, we estimate the right-hand side of (3.5) as follows:

$$\begin{aligned} \partial_{tx}^2 W_\lambda^\varepsilon &= -W_\lambda^\varepsilon \partial_{xx}^2 W_\lambda^\varepsilon - (\partial_x W_\lambda^\varepsilon)^2 - \lambda W_\lambda^\varepsilon \partial_x W_\lambda^\varepsilon + \lambda^2 \int_{-\infty}^x \exp(\lambda(y-x)) W_\lambda^\varepsilon(t,y) \partial_y W_\lambda^\varepsilon(t,y) dy \\ &= -W_\lambda^\varepsilon \partial_{xx}^2 W_\lambda^\varepsilon - (\partial_x W_\lambda^\varepsilon)^2 - \lambda W_\lambda^\varepsilon \partial_x W_\lambda^\varepsilon \\ &\quad + \lambda^2 \int_{-\infty}^x \exp(\lambda(y-x)) \left(\rho_\lambda(t,y) - \lambda^{-1} \partial_y W_\lambda^\varepsilon(t,y) \right) \partial_y W_\lambda^\varepsilon(t,y) dy \\ &= -W_\lambda^\varepsilon \partial_{xx}^2 W_\lambda^\varepsilon - (\partial_x W_\lambda^\varepsilon)^2 - \lambda W_\lambda^\varepsilon \partial_x W_\lambda^\varepsilon \\ &\quad + \lambda^2 \int_{-\infty}^x \exp(\lambda(y-x)) \rho_\lambda^\varepsilon(t,y) \partial_y W_\lambda^\varepsilon(t,y) dy - \underbrace{\lambda \int_{-\infty}^x \exp(\lambda(y-x)) |\partial_y W_\lambda^\varepsilon(t,y)|^2 dy}_{\leq 0} \\ &\leq -W_\lambda^\varepsilon \partial_{xx}^2 W_\lambda^\varepsilon - (\partial_x W_\lambda^\varepsilon)^2 - \lambda W_\lambda^\varepsilon \partial_x W_\lambda^\varepsilon + m(t) \lambda^2 \underbrace{\int_{-\infty}^x \exp(\lambda(y-x)) \rho_\lambda^\varepsilon(t,y) dy}_{= \lambda^{-1} W_\lambda^\varepsilon(t,x)}. \end{aligned}$$

We have that, for every $t > 0$, there exists a maximum point of $\partial_y W_\lambda^\varepsilon(t,y)$ (by choosing, e.g., a compactly supported $\rho_{0,\lambda}^\varepsilon$ and relying on the regularity results of [29]). Using [15, theorem 2.1], we consider $\bar{x}(t) \in \mathbb{R}$ such that $m(t) = \partial_x W_\lambda^\varepsilon(t, \bar{x}(t))$, evaluate the previous expression at $x = \bar{x}(t)$, and compute

$$\frac{d}{dt} m(t) \leq -m^2(t).$$

Since $\tilde{m}(t) = 1/t$ is a solution of the above Riccati-type differential inequality and $\tilde{m}(0) = \infty$, we use the comparison principle for ODEs to conclude that $m(t) \leq 1/t$ and thus

$$\frac{W_\lambda^\varepsilon(t,x) - W_\lambda^\varepsilon(t,y)}{x-y} = \frac{1}{x-y} \int_y^x \partial_x W_\lambda^\varepsilon(t,\xi) d\xi \leq \frac{1}{t}, \quad t > 0, \quad x, y \in \mathbb{R}, x \neq y.$$

Taking the limit $\varepsilon \rightarrow 0^+$, thanks to lemma 3.1, we conclude the proof. $\square$

As a by-product of (3.4), we prove (arguing as in [17, lemma 1.3]) that a L^∞ -bound holds for all $t > 0$ (which blows up as $t \rightarrow 0^+$).

Lemma 3.3 (L^∞ -bounds on W_λ and ρ_λ). *The following L^∞ -bounds on W_λ and ρ_λ hold:*

$$0 \leq W_\lambda(t,x) \leq \sqrt{\frac{2M}{t}}, \quad (t,x) \in (0, +\infty) \times \mathbb{R}, \quad (3.6)$$

$$0 \leq \rho_\lambda(t,x) \leq \sqrt{\frac{2M}{t}} + \frac{1}{\lambda t}, \quad (t,x) \in (0, +\infty) \times \mathbb{R}. \quad (3.7)$$

Proof. The fact that, for all $t > 0$, $W_\lambda(t, \cdot), \rho_\lambda(t, \cdot) \geq 0$ holds is contained in point (1) of theorem 2.1. To prove the upper bound in (3.6), let us fix a time $t > 0$ and a point $\bar{x} \in \mathbb{R}$. By lemma 3.2, we have

$$W_\lambda(t, x) \geq W_\lambda(t, \bar{x}) - \frac{1}{t}(\bar{x} - x), \quad \text{for all } x \leq \bar{x},$$

i.e.

$$W_\lambda(t, x) \geq \frac{1}{t}(x - (\bar{x} + W_\lambda(t, \bar{x})t)), \quad \text{for all } 0 \leq x - (\bar{x} + W_\lambda(t, \bar{x})t) \leq W_\lambda(t, \bar{x})t.$$

Integrating over $\mathbb{R}$, we deduce

$$M = \int_{\mathbb{R}} W_\lambda(t, x) dx \geq \int_{\mathbb{R} \cap \{x \geq \bar{x} + W_\lambda(t, \bar{x})t\}} \frac{x - (\bar{x} + W_\lambda(t, \bar{x})t)}{t} dx \geq \int_0^{W_\lambda(t, \bar{x})t} \left(\frac{x}{t}\right) dx = \frac{1}{2} W_\lambda(t, \bar{x})^2 t,$$

which implies

$$W_\lambda(t, \bar{x}) \leq \sqrt{\frac{2M}{t}} \quad \text{for all } t > 0, \bar{x} \in \mathbb{R}.$$

The bound (3.7) follows from (3.6) and theorem 3.2. Indeed, by (1.11), we have

$$\begin{aligned} 0 \leq \rho_\lambda(t, x) &= W_\lambda(t, x) + \frac{1}{\lambda} \partial_x W_\lambda(t, x) \\ &\leq \sqrt{\frac{2M}{t}} + \frac{1}{\lambda t}. \end{aligned}$$

□

As a second corollary, from (3.4), we deduce the following BV_{loc} regularisation result (see [5, equation (4.3)] and [6, lemma 2.2(ii) and remark 2.3]).

Corollary 3.4 (BV-regularisation effect). *The function $W_\lambda(t, \cdot)$ belongs to $BV_{\text{loc}}(\mathbb{R})$ for every $t > 0$ and uniformly with respect to $\lambda > 0$: namely, for every compact interval $K \Subset \mathbb{R}$,*

$$|W_\lambda(t, \cdot)|_{TV(K)} \leq 2 \left(\frac{|K|}{t} + \|W_\lambda(t, \cdot)\|_{L^\infty(K)} \right), \quad t > 0. \quad (3.8)$$

Proof. Let $K := [a, b] \Subset \mathbb{R}$ be a compact interval of $\mathbb{R}$ and fix $t > 0$. Since $W_\lambda(t, \cdot) \in L^\infty(K) \subset L^1(K)$, we only need to prove, thanks to the characterisation of BV functions in [41, lemma 37.4] (see also [1, remark 2.5 and exercise 3.3] or [36, corollary 2.17]), that there exists $C > 0$ such that

$$\int_{K_h} \frac{|W_\lambda(x+h) - W_\lambda(x)|}{h} dx \leq C, \quad \forall h > 0, \text{ where } K_h := \{x \in K : x+h \in K\}.$$

Taking lemma 3.2 into account, we note that

$$\frac{W_\lambda(t, x+h) - W_\lambda(t, x)}{h} = \frac{1}{t} - \underbrace{\left(\frac{1}{t} - \frac{W_\lambda(t, x+h) - W_\lambda(t, x)}{h} \right)}_{\geq 0}, \quad (3.9)$$

which implies

$$\frac{W_\lambda(t, x+h) - W_\lambda(t, x)}{h} \leq \frac{1}{t} + \left(\frac{1}{t} - \frac{W_\lambda(t, x+h) - W_\lambda(t, x)}{h} \right).$$

Integrating over K_h and taking the absolute values on both sides yields

$$\begin{aligned}
 \int_{K_h} \frac{|W_\lambda(t, x+h) - W_\lambda(t, x)|}{h} dx &\leq \int_{K_h} \left(\frac{1}{t} + \left(\frac{1}{t} - \frac{W_\lambda(t, x+h) - W_\lambda(t, x)}{h} \right) \right) dx \\
 &= 2 \int_{K_h} \frac{1}{t} dx + \int_{\mathbb{R}} W_\lambda(t, x) \left(\frac{\mathbb{1}_K(x+h) - \mathbb{1}_K(x)}{h} \right) dx \\
 &\leq 2 \int_{K_h} \frac{1}{t} dx + \|W_\lambda(t, \cdot)\|_{L^\infty(K)} \underbrace{\int_{\mathbb{R}} \left(\frac{|\mathbb{1}_K(x+h) - \mathbb{1}_K(x)|}{h} \right) dx}_{=\text{TV}(\mathbb{1}_K)=2} \\
 &= 2 \left(\frac{|K|}{t} + \|W_\lambda(t, \cdot)\|_{L^\infty(K)} \right).
 \end{aligned}$$

□

4. Long-time behaviour

As a first step towards finishing the proof of theorem 1.1, we shall show next that $\{W_\lambda\}_{\lambda>0}$ is compact in the canonical $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$ topology. Note that the time-interval does not include $t=0$ because the L^∞ -estimate from lemma 3.3 blows up as $t \rightarrow 0^+$.

Lemma 4.1 (compactness of $\{W_\lambda\}_{\lambda>0}$ in $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$). *Let $t_0, T > 0$ be fixed. The set $\{W_\lambda\}_{\lambda>0} \subseteq C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$ of solutions to (1.1) is compactly embedded into $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$, i.e.*

$$\left\{ W_\lambda \in C([t_0, T]; L^1_{\text{loc}}(\mathbb{R})) : W_\lambda \text{ satisfies (1.10), } \lambda > 0 \right\} \xrightarrow{c} C([t_0, T]; L^1_{\text{loc}}(\mathbb{R})).$$

Proof. Arguing as in [11, theorem 4.1], we shall apply the compactness result in [40, lemma 1]: given a Banach space B , a set $F \subset C([t_0, T]; B)$ is relatively compact in $C([t_0, T]; B)$ iff

- $F(t) := \{f(t) \in B : f \in F\}$ is relatively compact in B for all $t \in [t_0, T]$;
- F is uniformly equi-continuous, i.e.

$$\forall \sigma \in \mathbb{R}_{>0} \exists \delta \in \mathbb{R}_{>0} \text{ s. t. } \forall f \in F \forall (t_1, t_2) \in [t_0, T]^2 \text{ with } |t_1 - t_2| \leq \delta : \|f(t_1) - f(t_2)\|_B \leq \sigma.$$

In our case, let us fix a compact interval $K \Subset \mathbb{R}$ and define $B = L^1(K)$ and $F(t) := \{W_\lambda(t, \cdot) \in L^1(K) : \lambda > 0\}$.

Thanks to lemma 3.2, we know that $W_\lambda(t, \cdot)$ has a uniform total variation bound; thus, by [35, theorem 13.35], the set $F(t)$ is compact in $L^1(K)$, i.e.

$$F(t) \Subset L^1(K), \quad \text{for all } t \in [t_0, T].$$

It remains to show the second point, the uniform equi-continuity. To this end, we again replace the initial datum $\rho_{0,\lambda}$ by a smooth $\rho_{0,\lambda}^\varepsilon$, with $\varepsilon > 0$, and call the corresponding smooth nonlocal term W_λ^ε . Then, we can estimate

$$\begin{aligned}
\|W_\lambda^\varepsilon(t_1, \cdot) - W_\lambda^\varepsilon(t_2, \cdot)\|_{L^1(\mathbb{R})} &= \left\| \int_{t_2}^{t_1} \partial_t W_\lambda^\varepsilon(s, \cdot) \, ds \right\|_{L^1(\mathbb{R})} \\
&\leq \left\| \int_{t_2}^{t_1} W_\lambda^\varepsilon(s, \cdot) \partial_2 W_\lambda^\varepsilon(s, \cdot) \, ds \right\|_{L^1(\mathbb{R})} \\
&\quad + \left\| \int_{t_2}^{t_1} \lambda \int_*^\infty \exp(\lambda(* - y)) \partial_y W_\lambda^\varepsilon(s, y) W_\lambda^\varepsilon(s, y) \, dy \, ds \right\|_{L^1(\mathbb{R})} \\
&\leq \|W_\lambda^\varepsilon\|_{L^\infty((0, T); L^\infty(\mathbb{R}))} \|W_\lambda^\varepsilon\|_{L^\infty((0, T); \text{TV}(\mathbb{R}))} |t_1 - t_2| \\
&\quad + \|W_\lambda^\varepsilon\|_{L^\infty((0, T); L^\infty(\mathbb{R}))} \|W_\lambda^\varepsilon\|_{L^\infty((0, T); \text{TV}(\mathbb{R}))} |t_1 - t_2|,
\end{aligned}$$

where we used Fubini-Tonelli's theorem to exchange the order of integration and estimate the last term. Thanks to lemmas 3.2 and 3.3, we have that this is a uniform bound in $\lambda > 0$ and $\varepsilon > 0$. This yields the uniform equi-continuity so that we obtain indeed the claimed compactness. $\square$

We can now complete the proof of theorem 1.1 arguing as in [17, section 2].

Proof of theorem 1.1. The core of the proof consists in showing that the family $\{\rho_\lambda\}_{\lambda>0}$ converges to the N -wave defined in (1.6). We shall divide the argument of this theorem in several steps.

Step 1. *Compactness of the family $\{W_\lambda\}_{\lambda>0}$ in $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$.* For any $0 < t_0 < T$, by lemma 4.1, we have that W_λ converges (up to extracting a subsequence) to a limit point w^* strongly in $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$; hence, we also have $W_\lambda(t, \cdot) \rightarrow w^*(t, \cdot)$ in $L^1_{\text{loc}}(\mathbb{R})$ for all $t \in [t_0, T]$ and $W_\lambda \rightarrow w^*$ pointwise (again up to subsequences) for all $t \in [t_0, T]$ and a.e. $x \in \mathbb{R}$.

Thanks to (1.11), we can deduce that ρ_λ also converges to w^* along the same subsequence. Indeed, first we observe that

$$\|W_\lambda(t, \cdot) - \rho_\lambda(t, \cdot)\|_{L^1(\mathbb{R})} = \lambda^{-1} |W_\lambda(t, \cdot)|_{\text{TV}(\mathbb{R})}$$

and thus we also obtain

$$\lim_{\lambda \rightarrow +\infty} \|\rho_\lambda - w\|_{C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))} = 0.$$

Step 2a. *Tail control and convergence of the family $\{\rho_\lambda\}_{\lambda>0}$ in $C([t_0, T], L^1(\mathbb{R}))$.* In order to pass from the convergence $\rho_\lambda \rightarrow w^*$ strongly in $C([t_0, T]; L^1_{\text{loc}}(\mathbb{R}))$ to the convergence in $C([t_0, T]; L^1(\mathbb{R}))$, we need a uniform bound on the ‘tail’ of the functions $\{\rho_\lambda\}_{\lambda>1}$. We shall prove that there exists a constant $C = C(M)$ such that

$$\int_{\{|x|>2R\}} \rho_\lambda(t, x) \, dx \leq \int_{\{|x|>R\}} \rho_0(x) \, dx + \frac{C(M)}{R} t^{1/2}, \quad t > 0. \quad (4.1)$$

Since $\rho_0 \in L^1(\mathbb{R})$, the right-hand side of (4.1) can be made arbitrarily small choosing R large enough. Then, from (4.1), the convergence

$$\rho_\lambda \rightarrow w^* \quad \text{strongly in } C([t_0, T], L^1(\mathbb{R})) \quad \text{as } \lambda \rightarrow +\infty$$

follows by considering the splitting

$$\int_{\mathbb{R}} |\rho_\lambda(t, x) - w^*(t, x)| \, dx = \int_{\{x < 2R\}} |\rho_\lambda(t, x) - w^*(t, x)| \, dx + \int_{\{x > 2R\}} |\rho_\lambda(t, x) - w^*(t, x)| \, dx.$$

In order to prove (4.1), let us consider a test function $\varphi \in C^\infty(\mathbb{R})$ such that $0 \leq \varphi \leq 1$, $\varphi \equiv 1$ for $|x| > 2$, and $\varphi \equiv 0$ for $|x| \leq 1$; we consider the rescaling $\varphi_R := \varphi(\cdot/R)$ which satisfies $\|\partial_x \varphi_R\|_{L^\infty(\mathbb{R})} \leq C/R$ for some $C > 0$. Let us multiply the PDE in (1.9) by φ_R , integrate in $(0, t) \times \mathbb{R}$ (for some $t > 0$), and perform an integration by parts (to rigorously justify this computation, we can use a smoothing argument based on lemma 3.1):

$$\int_{\mathbb{R}} \rho_\lambda(t, x) \varphi_R(x) dx = \int_{\mathbb{R}} \rho_\lambda(0, x) \varphi_R(x) dx + \int_0^t \int_{\mathbb{R}} \rho_\lambda(s, x) W_\lambda(s, x) \partial_x \varphi_R(x) dx ds.$$

We remark that

$$\begin{aligned} \int_{\mathbb{R}} \rho_\lambda(0, x) \varphi_R(x) dx &= \int_{\{|x| \geq R\}} \rho_\lambda(0, x) dx = \int_{\{|x| \geq \lambda R\}} \rho(0, x) dx \leq \int_{\{|x| > R\}} \rho(0, x) dx, \\ \int_0^t \int_{\mathbb{R}} \rho_\lambda(s, x) W_\lambda(s, x) \partial_x \varphi_R(x) dx ds &\leq \|\partial_x \varphi_R\|_{L^\infty(\mathbb{R})} \int_0^t \|\rho_\lambda(s, \cdot)\|_{L^1(\mathbb{R})} \|W_\lambda(s, \cdot)\|_{L^\infty(\mathbb{R})} ds \\ &\leq \frac{C}{R} \int_0^t M \sqrt{\frac{2M}{s}} ds \leq \frac{C}{R} \frac{1}{\sqrt{2}} M^{3/2} t^{1/2}, \end{aligned}$$

where we used lemma 3.3 in the last line.

Step 2b. Tail control and convergence of the family $\{W_\lambda\}_{\lambda > 0}$ in $C([t_0, T], L^1(\mathbb{R}))$. Since

$$\int_{\{|x| > 2R\}} W_\lambda(t, x) dx = \lambda \int_{\{|x| > 2R\}} \int_{-\infty}^x \exp(\lambda(y-x)) \rho(t, y) dy dx,$$

we use Fubini-Tonelli's theorem to deduce

$$\int_{\{|x| > 2R\}} W_\lambda(t, x) dx \leq \int_{\{|x| > 2R\}} \rho_\lambda(t, x) dx, \quad t > 0,$$

which yields, thanks to (4.1),

$$\int_{\{|x| > 2R\}} W_\lambda(t, x) dx \leq \int_{\{|x| > R\}} \rho_0(x) dx + \frac{C(M)}{R} t^{1/2}, \quad t > 0. \quad (4.2)$$

As a byproduct of Steps 1 and 2, we note that the limit point w^* satisfies

$$w^* \in C((0, +\infty); L^1(\mathbb{R}; \mathbb{R}_{\geq 0})) \cap L^\infty((\tau, +\infty); L^\infty(\mathbb{R}; \mathbb{R}_{\geq 0})) \text{ for all } \tau > 0, \quad \int_{\mathbb{R}} w^*(t, x) dx = M.$$

Step 3. Identification of the initial condition. We now identify the initial datum taken by the limit point w^* , i.e. we verify that the initial condition $M\delta_0$ is achieved in the weak sense of non-negative measures on $\mathbb{R}$. We need to prove that, for all $\varphi \in C_b(\mathbb{R})$,

$$\lim_{t \rightarrow 0^+} \int_{\mathbb{R}} w^*(t, x) \varphi(x) dx = M\varphi(0).$$

To this end, arguing as in [17, pp 52–54], we shall split the argument into two steps. First, we consider a smaller class of test functions $\varphi \in C_c^\infty(\mathbb{R}; [0, 1])$ and secondly $\varphi \in C_b(\mathbb{R})$.

We start by estimating, for a test function $\varphi \in C_c^\infty(\mathbb{R}; [0, 1])$,

$$\begin{aligned} & \left| \int_{\mathbb{R}} \rho_\lambda(t, x) \varphi(x) dx - \int_{\mathbb{R}} \rho_\lambda(0, x) \varphi(x) dx \right| \\ & \leq \left| \int_0^t \int_{\mathbb{R}} \partial_x \varphi(x) W_\lambda(s, x) \rho_\lambda(s, x) dx ds \right| \\ & \leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \|\rho_\lambda(s, \cdot)\|_{L^1(\mathbb{R})} \|W_\lambda(s, \cdot)\|_{L^\infty(\mathbb{R})} ds \\ & \leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} C(M) \int_0^t s^{-1/2} ds \leq C(M) \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \sqrt{t}. \end{aligned}$$

Then, letting $\lambda \rightarrow +\infty$, we obtain

$$\left| \int_{\mathbb{R}} w^*(t, x) - M\varphi(0) \right| \leq C(M) \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \sqrt{t},$$

which, in turn, goes to zero as $t \rightarrow 0^+$.

As a second step, let us consider the case of a bounded continuous function $\varphi \in C_b(\mathbb{R})$. We shall rely on an approximation argument and on the tail control of ρ_λ in (4.1). Let us consider a regularised test function φ_ε obtained as $\varphi_\varepsilon := \varphi * \eta_\varepsilon$ (where η_ε denotes a standard mollifier; see [21, appendix C.4]), such that $\|\varphi_\varepsilon\|_{L^\infty(\mathbb{R})} \leq \|\varphi\|_{L^\infty(\mathbb{R})}$, $\varphi_\varepsilon \rightarrow \varphi$ uniformly on compact sets of $\mathbb{R}$ as $\varepsilon \rightarrow 0^+$, and $\|\varphi_\varepsilon\|_{W^{1,\infty}(\mathbb{R})} \leq C(\varepsilon)$. We then write

$$\begin{aligned} \left| \int_{\mathbb{R}} \rho_\lambda(t, x) \varphi(x) dx - M\varphi(0) \right| & \leq \left| \int_{\mathbb{R}} \rho_\lambda(t, x) \varphi_\varepsilon(x) dx - M\varphi(0) \right| \\ & \quad + \left| \int_{\{|x|>2R\}} \rho_\lambda(t, x) (\varphi(x) - \varphi_\varepsilon(x)) dx \right| \\ & \quad + \left| \int_{\{|x|<2R\}} \rho_\lambda(t, x) (\varphi(x) - \varphi_\varepsilon(x)) dx \right|. \end{aligned}$$

The control of the first term follows by the same argument developed above. For the second and third term, we estimate

$$\begin{aligned} \left| \int_{\{|x|>2R\}} \rho_\lambda(t, x) (\varphi(x) - \varphi_\varepsilon(x)) dx \right| & \leq 2\|\varphi\|_{L^\infty(\mathbb{R})} \left(\int_{\{|x|>R\}} \rho_0(x) dx + \frac{C(M)}{R} t^{1/2} \right), \\ \left| \int_{\{|x|<2R\}} \rho_\lambda(t, x) (\varphi(x) - \varphi_\varepsilon(x)) dx \right| & \leq \|\varphi - \varphi_\varepsilon\|_{L^\infty(\{|x|<2R\})} \int_{\mathbb{R}} \rho_\lambda(t, x) dx = M\|\varphi - \varphi_\varepsilon\|_{L^\infty(\{|x|<2R\})}, \end{aligned}$$

which can both be made arbitrarily small provided that $\varepsilon > 0$ is small enough and $R > 0$ is large enough.

A similar argument can be used for $\{W_\lambda\}_{\lambda>0}$. Indeed, for $\varphi \in C_c^\infty(\mathbb{R}; [0, 1])$, we estimate

$$\begin{aligned} & \left| \int_{\mathbb{R}} W_\lambda(t, x) \varphi(x) dx - \int_{\mathbb{R}} W_\lambda(0, x) \varphi(x) dx \right| = \left| \int_0^t \int_{\mathbb{R}} \partial_t W_\lambda(s, x) \varphi(x) dx ds \right| \\ & \leq \underbrace{\int_0^t \int_{\mathbb{R}} \partial_x \varphi(x) |W_\lambda(s, x)|^2 dx ds}_{=: I_1} \\ & \quad + \lambda \underbrace{\left| \int_0^t \int_{\mathbb{R}} \varphi(x) \int_{-\infty}^x \exp(\lambda(y-x)) (W_\lambda(s, x) \partial_x W_\lambda(s, x) - W_\lambda(s, y) \partial_y W_\lambda(s, y)) dy dx ds \right|}_{=: I_2}. \end{aligned}$$

For the term I_1 , we compute

$$\begin{aligned} I_1 &\leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \|W_\lambda(s, \cdot)\|_{L^1(\mathbb{R})} \|W_\lambda(s, \cdot)\|_{L^\infty(\mathbb{R})} ds \\ &\leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} C(M) \int_0^t s^{-1/2} ds \leq C(M) \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \sqrt{t}, \end{aligned}$$

where, in the last line, we used lemma 3.3.

For I_2 , using Fubini-Tonelli's theorem, we compute

$$\begin{aligned} I_2 &= \left| \int_0^t \int_{\mathbb{R}} \varphi(x) W_\lambda(s, x) \partial_x W_\lambda(s, x) dx ds \right. \\ &\quad \left. - \lambda \int_0^t \int_{\mathbb{R}} \varphi(x) \int_{-\infty}^x \exp(\lambda(y-x)) W_\lambda(s, y) \partial_y W_\lambda(s, y) dy dx ds \right| \\ &= \left| \int_0^t \int_{\mathbb{R}} \varphi(x) W_\lambda(s, x) \partial_x W_\lambda(s, x) dx ds \right. \\ &\quad \left. - \lambda \int_0^t \int_{\mathbb{R}} W_\lambda(s, y) \partial_y W_\lambda(s, y) \int_y^\infty \varphi(x) \exp(\lambda(y-x)) dx dy ds \right|; \end{aligned}$$

integrating by parts on the term $x \mapsto \exp(\lambda(y-x))$ yields

$$I_2 = \left| \int_0^t \int_{\mathbb{R}} W_\lambda(s, y) \partial_y W_\lambda(s, y) \int_y^\infty \partial_x \varphi(x) \exp(\lambda(y-x)) dx dy ds \right|;$$

integrating by parts on the term $y \mapsto W_\lambda(s, y) \partial_y W_\lambda(s, y)$ and using the fact that $\lim_{x \rightarrow \pm\infty} W_\lambda(t, \cdot) = 0$ (which is a consequence of the fact that $W_\lambda(t, \cdot) \in L^1(\mathbb{R}) \cap \text{BV}_{\text{loc}}(\mathbb{R})$ for $t > 0$ and $\lambda > 0$), we then get

$$\begin{aligned} I_2 &= \left| -\lambda \frac{1}{2} \int_0^t \int_{\mathbb{R}} |W_\lambda(s, y)|^2 \int_y^\infty \partial_x \varphi(x) \exp(\lambda(y-x)) dx dy ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}} |W_\lambda(s, y)|^2 \partial_y \varphi(y) dy ds \right| \\ &\leq \frac{\lambda}{2} \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \int_{\mathbb{R}} |W_\lambda(s, y)|^2 \int_y^\infty \exp(\lambda(y-x)) dx dy ds + \frac{1}{2} \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \int_{\mathbb{R}} |W_\lambda(s, y)|^2 dy ds \\ &= \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \int_{\mathbb{R}} |W_\lambda(s, y)|^2 dy ds \leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \int_0^t \|W_\lambda(s, \cdot)\|_{L^1(\mathbb{R})} \|W_\lambda(s, \cdot)\|_{L^\infty(\mathbb{R})} ds \\ &\leq \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} C(M) \int_0^t s^{-1/2} ds \leq C(M) \|\partial_x \varphi\|_{L^\infty(\mathbb{R})} \sqrt{t}, \end{aligned}$$

where, in the last line, we used lemma 3.3. Thus, for any $\varepsilon > 0$, we can choose $\tau > 0$ and $\lambda_0 > 0$ such that

$$\left| \int_{\mathbb{R}} W_\lambda(t, x) \varphi(x) dx - M \varphi(0) \right| \leq \varepsilon \quad \text{for all } 0 < t < \tau, \lambda > \lambda_0.$$

The rest of the argument for $\varphi \in C_b(\mathbb{R})$ goes through as above.

Step 4. Entropy admissibility of the limit point. The limit point w^* is actually the unique entropy admissible N -wave solution w of the Burgers equation (1.5) defined in (1.6). This follows immediately from passing to the limit pointwise in the Oleinik inequality (3.4). Thanks to Urysohn's subsequence principle, from the uniqueness of the entropy solution of (1.5), we also

deduce that the whole families $\{\rho_\lambda\}_{\lambda>0}$ and $\{W_\lambda\}_{\lambda>0}$ converge to w (not just up to extracting a subsequence).

Step 5. Conclusion of the proof. From the steps above, we have that

$$\|W_\lambda(t, \cdot) - w(t, \cdot)\|_{L^1(\mathbb{R})} \rightarrow 0 \quad \text{as } \lambda \rightarrow +\infty,$$

where w denotes the N -wave solution entropy of (1.5). For $p = 1$, (1.7) is a consequence of the fact that

$$\begin{aligned} \rho_\lambda(1, x) - w(1, x) &= \lambda \rho(\lambda^2, \lambda x) - w(1, x), \\ W_\lambda(1, x) - w(1, x) &= \lambda W(\lambda^2, \lambda x) - w(1, x), \end{aligned}$$

(and that the same would hold true replacing $t = 1$ by any fixed $\bar{t} > 0$), i.e. letting $\lambda \rightarrow +\infty$ for a fixed time $\bar{t} > 0$ is equivalent to fixing $\lambda = 1$ and letting $t \rightarrow +\infty$.

To prove the result also for $p \in (1, +\infty)$, we argue by interpolation. Indeed, we have that, for $t > 0$, $\{\rho_\lambda\}_{\lambda>0}$ and $\{W_\lambda\}_{\lambda>0}$ are also uniformly bounded in $L^q(\mathbb{R})$ (being in $L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ for every $t > 0$) and $w \in L^q(\mathbb{R})$, with $q \in (1, +\infty)$. Then, Hölder's inequality yields

$$\begin{aligned} \|\rho_\lambda(t, \cdot) - w(t, \cdot)\|_{L^p(\mathbb{R})} &\leq \|\rho_\lambda(t, \cdot) - w(t, \cdot)\|_{L^1(\mathbb{R})}^{\frac{1}{2p-1}} \left(\|\rho_\lambda(t, \cdot)\|_{L^{2p}(\mathbb{R})} + \|w(t, \cdot)\|_{L^{2p}(\mathbb{R})} \right)^{\frac{2(p-1)}{2p-1}}, \\ \|W_\lambda(t, \cdot) - w(t, \cdot)\|_{L^p(\mathbb{R})} &\leq \|W_\lambda(t, \cdot) - w(t, \cdot)\|_{L^1(\mathbb{R})}^{\frac{1}{2p-1}} \left(\|W_\lambda(t, \cdot)\|_{L^{2p}(\mathbb{R})} + \|w(t, \cdot)\|_{L^{2p}(\mathbb{R})} \right)^{\frac{2(p-1)}{2p-1}}, \end{aligned}$$

from which the result follows. $\square$

5. Numerical experiments

In this section, we showcase the result in theorem 1.1 numerically. For the nonlocal problem, we rely on a non-dissipative solver based on characteristics (see [31] for further details). More precisely, the simulations illustrate the convergence

$$\bar{\rho}(t, \cdot) \rightarrow \bar{w} \quad \text{in } L^1(\mathbb{R}) \text{ as } t \rightarrow +\infty,$$

for the rescaled variables

$$\bar{\rho} := \sqrt{t}\rho, \quad \bar{w} := \sqrt{t}w, \quad y := x/\sqrt{t},$$

in which the N -wave is stationary (i.e. time-independent) and given by

$$\bar{w}(y) = \begin{cases} \frac{y}{2} & \text{if } y \in (0, \sqrt{4M}), \\ 0 & \text{otherwise.} \end{cases}$$

First, in figure 2, we present the evolution of the solution of (1.1) on long time horizons for the following initial data:

$$\begin{aligned} \text{(1)} \quad \rho_0(x) &= \mathbb{1}_{[0,1]}(x), & \text{(2)} \quad \rho_0(x) &= 2x \mathbb{1}_{[0,1]}(x), \\ \text{(3)} \quad \rho_0(x) &= 6x(1-x) \mathbb{1}_{[0,1]}(x), & \text{(4)} \quad \rho_0(x) &= 2x \mathbb{1}_{[0,0.5]}(x) + \mathbb{1}_{[0.5,1]}(x), \end{aligned}$$

for $x \in \mathbb{R}$. In all cases, we observe the convergence towards the N -wave profile of the (local) Burgers equation (1.5).

For (left) continuous initial data (as is the case in (2), (3), and (4)), the N -wave is also approximated by left-continuous functions. This is a well-known fact, as nonlocal conservation laws preserve regularity [29, corollary 5.3] (see also theorem 2.1). In particular, for the (4) case, there are two jumps downwards in the initial datum and the first one is damped out over time




Figure 2. Convergence to the N -wave profile for the nonlocal regularisation of the Burgers equation. TOP LEFT: $\rho_0(x) = \mathbb{1}_{[0,1]}(x)$. TOP RIGHT: $\rho_0(x) = 2x \mathbb{1}_{[0,1]}(x)$. BOTTOM LEFT: $\rho_0(x) = 6x(1-x) \mathbb{1}_{[0,1]}(x)$. BOTTOM RIGHT: $\rho_0(x) = 2x \mathbb{1}_{[0,0.5]}(x) + \mathbb{1}_{[0.5,1]}(x)$.

(still observable for $t = 10$ at $x \approx 1$). This can be understood when recalling that around $x \approx 1$ the velocity of the dynamics is smaller than for $x < 1$ so that the density increases between both points and the jump decreases (which is visible in particular for $t = 1$ and $t = 10$).

Secondly, in figure 3, we consider $\gamma(x) := \mathbb{1}_{(0,1)}(x)$, $x \in \mathbb{R}$, instead of an exponential weight in (1.2), i.e. we study

$$W[\rho](t, x) := \int_{x-1}^x \rho(t, y) \, dy, \quad (t, x) \in (0, T) \times \mathbb{R}.$$

The numerical simulation shows that, even in this case (which is not covered by the results of the present paper or by the ones on the singular limit problem contained in [11, 14]), a convergence result can be observed. However, the convergence seems to occur ‘less regularly’ as the constant kernel generates more and more points where the solution is not differentiable.




Figure 3. Convergence to the N -wave profile for the nonlocal regularization of Burgers equation with the constant kernel $\gamma := \mathbb{1}_{(0,1)}$. LEFT: $\rho_0(x) = \mathbb{1}_{[0,1]}(x)$. RIGHT: $\rho_0(x) = 6x(1-x)\mathbb{1}_{[0,1]}(x)$.

Indeed, in contrast to the exponential kernel case, the regularity of the solution for piece-wise constant kernels depends points-wise and locally (on the trace of backward characteristics) on initial data, kernel, and their interplay.

Finally, we present some simulations illustrating the case of a more general power-type velocity: namely,

$$\begin{cases} \partial_t \rho(t, x) + \partial_x (W^{q-1}(t, x) \rho(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ \rho(0, x) = \rho_0(x), & x \in \mathbb{R}, \end{cases} \quad (5.1)$$

for some $q \geq 2$. In this case, the explicit N -wave solution of the corresponding local conservation law

$$\begin{cases} \partial_t \rho(t, x) + \partial_x (\rho^q(t, x)) = 0, & (t, x) \in (0, +\infty) \times \mathbb{R}, \\ \rho(0, x) = \rho_0(x), & x \in \mathbb{R}, \end{cases} \quad (5.2)$$

is given by

$$w_q(t, x) = \begin{cases} \left(\frac{x}{qt}\right)^{\frac{1}{q-1}} & \text{if } x \in \left(0, q \left(\frac{M}{q-1}\right)^{\frac{q-1}{q}} t^{\frac{1}{q}}\right), \\ 0 & \text{otherwise.} \end{cases}$$

that is, in the rescaled variables

$$\begin{aligned} \bar{w}_q &:= t^{1/q} w_q, \quad y := x t^{-1/q}, \\ \bar{w}_q(y) &= \begin{cases} \left(\frac{y}{q}\right)^{\frac{1}{q-1}} & \text{if } y \in \left(0, q \left(\frac{M}{q-1}\right)^{\frac{q-1}{q}}\right), \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

(see [37, equation (2.1)]). In particular, in figure 4 (for $q = 3$), the convergence result seems to hold.




Figure 4. Convergence to the N -wave profile for the nonlocal regularization of $\partial_t \rho + \partial_x \rho^3 = 0$ with exponential kernel. LEFT: $\rho_0(x) = \mathbb{1}_{[0,1]}(x)$. RIGHT: $\rho_0(x) = 6x(1-x)\mathbb{1}_{[0,1]}(x)$.

In this case, none of the previously established results hold. However, the numerical experiments point to the fact that we may still observe the L^1 -convergence to the N -wave profile. The behaviour of the rescaled solution, which explodes at $x=0$, is particularly noteworthy. It can be explained as follows. For the conservation law

$$\partial_t \rho(t, x) + \partial_x (W[\rho]^2(t, x) \rho(t, x)) = 0, \quad (t, x) \in (0, T) \times \mathbb{R},$$

we can compute, along characteristics (see [12, 29]),

$$\begin{aligned} \frac{d}{dt} \rho(t, \xi(0, x; t)) &= \partial_t \rho(t, \xi(0, x; t)) + \partial_2 \rho(t, \xi(0, x; t)) \partial_3 \xi(0, x; t) \\ &= -\partial_2 \rho(t, \xi(0, x; t)) W[\rho]^2(t, \xi(0, x; t)) \\ &\quad - 2\rho(t, \xi(0, x; t)) W[\rho](t, \xi(0, x; t)) \partial_2 W[\rho](t, \xi(0, x; t)) \\ &\quad + \partial_2 \rho(t, \xi(0, x; t)) W[\rho]^2(t, \xi(0, x; t)) \\ &= -2\rho(t, \xi(0, x; t)) W[\rho](t, \xi(0, x; t)) \partial_2 W[\rho](t, \xi(0, x; t)). \end{aligned}$$

As W ‘looks’ to the left and the solution vanishes on the left half-space for all time $t > 0$, we have that $W[\rho](t, 0) = 0$ for all $t > 0$; thus, the value of the solution at $x=0$ never changes, i.e. $\lim_{x \searrow 0} \rho(t, x) = \rho_0(0)$ for all $t > 0$, which yields the long-time behaviour at $x=0$ observed in figure 4 upon rescaling.

6. Conclusions

In this contribution, we have proved the convergence of the solution of the nonlocal conservation law (1.1) with bounded, integrable, and non-negative initial datum to the N -wave solution of the Burgers equation as $t \rightarrow +\infty$.

Several open problems and possible generalizations of this result could be of interest for future work. We mention a few below.

- (1) *Sign-changing initial data.* The classical references that deal with the case of local conservation laws (starting from, e.g. [37]) study the case of sign-changing initial data as well; in the nonlocal setting, however, a non-negativity assumption seems to be needed in order to obtain a global existence result (see [29]).
- (2) *Non-integrable initial data.* Considering initial data merely in L^∞ instead of $L^1 \cap L^\infty$ poses a significant issue: since solutions might then have infinite mass, the compactness arguments would need to be revised and the limit profile would no longer be governed by the initial mass. We refer to [24] for the study of this problem for the heat equation. In the periodic setting, further information is available for the long-time behaviour of (local) conservation laws (see, e.g. [9, 34]).
- (3) *More general velocities.* While in the local case the available body of literature deals with general power nonlinearities (as mentioned in the introduction), in the nonlocal case a power-type velocity, as in $\partial_t \rho + \partial_x(W^{q-1}\rho) = 0$ (with $q > 2$), does not seem to generally allow for an Oleinik-type condition (see [10] for further discussion), which was pivotal in the approach used in the present paper.
- (4) *General convolution kernels.* For more general convolution kernels, nonlocal-to-local convergence results have been obtained (see [14]); however, no Oleinik inequalities are currently available (whose regularization effect is needed to treat the case of initial data that are not uniformly bounded in L^∞ with respect to the scaling parameter). The numerical simulations from section 5 still show that the convergence to the local N -wave should hold.
- (5) *Nonlocality in the velocity.* A different type of nonlocal conservation laws presents non-local effect of the form $\partial_t \rho + \partial_x(\rho(\rho^{q-1} * \gamma)) = 0$, i.e. where the averaging is not done over the solution, but over the velocity. A recent contribution on the nonlocal-to-local singular limit problem for this kind of models is contained in [22].
- (6) *Numerical schemes.* The development of numerical schemes that reproduce qualitatively the long-time behaviour of the solution has been addressed in works dealing with diffusion operators and local velocities (see, e.g. [25–27]), but it appears to be fully unexplored in the nonlocal setting.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgment

We would like to thank D Serre for his remarks on the Oleinik inequality and M Colombo, G Crippa, E Marconi, and L V Spinolo for several helpful conversations on topics related to this work. We also acknowledge the referees for their comments, which improved the quality of the manuscript.

G M Coclite and N De Nitti are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

G M Coclite has been partially supported by the Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4—Call for tender No. 3138 of 16 December 2021 of Italian Ministry of University and Research funded by the European Union—NextGenerationEUoAward Number: CN000023, Concession Decree No.

1033 of 17 December 2022 adopted by the Italian Ministry of University and Research, CUP: D93C22000410001, Centro Nazionale per la Mobilità Sostenibile and the Italian Ministry of Education, University and Research under the Programme Department of Excellence Legge 232/2016 (Grant No. CUP—D93C23000100001).

L Pflug has been supported by the Deutsche Forschungsgemeinschaft (DFG)—Project-ID 41 622 9255—SFB 1411.

E Zuazua has been funded by the Alexander von Humboldt-Professorship program, the ModConFlex Marie Curie Action, HORIZON-MSCA-2021-DN-01, the COST Action MAT-DYN-NET, the Transregio 154 Project ‘Mathematical Modelling, Simulation and Optimization Using the Example of Gas Networks’ of the DFG, Grants PID2020-11 2617GB-C22 and TED2021-13 1390B-I00 of MINECO (Spain), and by the Madrid Government—UAM Agreement for the Excellence of the University Research Staff in the context of the V PRICIT (Regional Programme of Research and Technological Innovation).

ORCID iDs

Giuseppe Maria Coclite  <https://orcid.org/0000-0001-6019-4757>

Nicola De Nitti  <https://orcid.org/0000-0003-0402-7502>

Alexander Keimer  <https://orcid.org/0000-0003-3825-5853>

Lukas Pflug  <https://orcid.org/0000-0001-8001-5832>

Enrique Zuazua  <https://orcid.org/0000-0002-1377-0958>

References

- [1] Ambrosio L, Fusco N and Pallara D 2000 *Functions of Bounded Variation and Free Discontinuity Problems (Oxford Mathematical Monographs)* (The Clarendon)
- [2] Amorim P, Colombo R M and Teixeira A 2015 On the numerical integration of scalar nonlocal conservation laws *ESAIM Math. Model. Numer. Anal.* **49** 19–37
- [3] Biler P, Karch G and Woyczyński W A 2001 Critical nonlinearity exponent and self-similar asymptotics for Lévy conservation laws *Ann. Inst. Henri Poincaré C* **18** 613–37
- [4] Bogachev V I 2007 *Measure Theory* vol I, II (Springer)
- [5] Bouchut F and James F 1998 One-dimensional transport equations with discontinuous coefficients *Nonlinear Anal.* **32** 891–933
- [6] Bouchut F, James F and Mancini S 2005 Uniqueness and weak stability for multi-dimensional transport equations with one-sided Lipschitz coefficient *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **4** 1–25
- [7] Bressan A and Shen W 2020 On traffic flow with nonlocal flux: a relaxation representation *Arch. Ration. Mech. Anal.* **237** 1213–36
- [8] Bressan A and Shen W 2021 Entropy admissibility of the limit solution for a nonlocal model of traffic flow *Commun. Math. Sci.* **19** 1447–50
- [9] Chen G-Q and Frid H 1999 Decay of entropy solutions of nonlinear conservation laws *Arch. Ration. Mech. Anal.* **146** 95–127
- [10] Coclite G M, Colombo M, Crippa G, De Nitti N, Keimer A, Marconi E, Pflug L and Spinolo L V 2023 Oleĭnik-type estimates for nonlocal conservation laws and applications to the nonlocal-to-local limit (arXiv:2304.01309)
- [11] Coclite G M, Coron J-M, De Nitti N, Keimer A and Pflug L 2022 A general result on the approximation of local conservation laws by nonlocal conservation laws: the singular limit problem for exponential kernels *Ann. Inst. Henri Poincaré C* (<https://doi.org/10.4171/aihpc/58>)
- [12] Coclite G M, De Nitti N, Keimer A and Pflug L 2022 On existence and uniqueness of weak solutions to nonlocal conservation laws with BV kernels *Z. Angew. Math. Phys.* **73** 241
- [13] Colombo M, Crippa G, Marconi E and Spinolo L V 2021 Local limit of nonlocal traffic models: convergence results and total variation blow-up *Ann. Inst. Henri Poincaré C* **38** 1653–66

- [14] Colombo M, Crippa G, Marconi E and Spinolo L V 2023 Nonlocal traffic models with general kernels: singular limit, entropy admissibility and convergence rate *Arch. Ration. Mech. Anal.* **247** 32
- [15] Constantin A and Escher J 1998 Wave breaking for nonlinear nonlocal shallow water equations *Acta Math.* **181** 229–43
- [16] Endal J, Ignat L I and Quirós F 2022 Large-time behaviour for anisotropic stable nonlocal diffusion problems with convection (arXiv:2207.01874)
- [17] Escobedo M, Vázquez J L and Zuazua E 1993 Asymptotic behaviour and source-type solutions for a diffusion-convection equation *Arch. Ration. Mech. Anal.* **124** 43–65
- [18] Escobedo M, Vázquez J L and Zuazua E 1993 A diffusion-convection equation in several space dimensions *Indiana Univ. Math. J.* **42** 1413–40
- [19] Escobedo M and Zuazua E 1997 Long-time behavior for a convection-diffusion equation in higher dimensions *SIAM J. Math. Anal.* **28** 570–94
- [20] Escobedo M and Zuazua E 1999 Long-time behaviour of diffusion waves for a viscous system of conservation laws in $\mathbf{R}^N$ *Asymptotic Anal.* **20** 133–73
- [21] Evans L C 2010 *Partial Differential Equations (Graduate Studies in Mathematics vol 19)* 2nd edn (American Mathematical Society)
- [22] Friedrich J, Göttlich S, Keimer A and Pflug L 2022 Conservation laws with nonlocal velocity—the singular limit problem (arXiv:2210.12141)
- [23] Goatin P and Scialanga S 2016 Well-posedness and finite volume approximations of the LWR traffic flow model with non-local velocity *Netw. Heterog. Media* **11** 107–21
- [24] Herranz L 1999 Asymptotic behaviour of solutions of some semilinear parabolic problems *Ann. Inst. Henri Poincaré C* **16** 49–105
- [25] Ignat L I and Pozo A 2017 A semi-discrete large-time behavior preserving scheme for the augmented Burgers equation *ESAIM Math. Model. Numer. Anal.* **51** 2367–98
- [26] Ignat L I and Pozo A 2018 A splitting method for the augmented Burgers equation *BIT Numer. Math.* **58** 73–102
- [27] Ignat L I, Pozo A and Zuazua E 2015 Large-time asymptotics, vanishing viscosity and numerics for 1-D scalar conservation laws *Math. Comput.* **84** 1633–62
- [28] Ignat L I and Stan D 2018 Asymptotic behavior of solutions to fractional diffusion-convection equations *J. London Math. Soc.* **97** 258–81
- [29] Keimer A and Pflug L 2017 Existence, uniqueness and regularity results on nonlocal balance laws *J. Differ. Equ.* **263** 4023–69
- [30] Keimer A and Pflug L 2019 On approximation of local conservation laws by nonlocal conservation laws *J. Math. Anal. Appl.* **475** 1927–55
- [31] Keimer A, Pflug L and Spinola M 2020 Nonlocal balance laws: theory of convergence for nondissipative numerical schemes submitted
- [32] Kim Y J 2004 An Oleinik-type estimate for a convection-diffusion equation and convergence to N -waves *J. Differ. Equ.* **199** 269–89
- [33] Laurencot P 1998 Long-time behaviour for diffusion equations with fast convection *Ann. Mat. Pura Appl.* **175** 233–51
- [34] Lax P D 1972 The formation and decay of shock waves *Am. Math. Mon.* **79** 227–41
- [35] Leoni G 2009 *A First Course in Sobolev Spaces (Graduate Studies in Mathematics vol 105)* (American Mathematical Society)
- [36] Leoni G 2017 *A First Course in Sobolev Spaces (Graduate Studies in Mathematics vol 181)* 2nd edn (American Mathematical Society)
- [37] Liu T-P and Pierre M 1984 Source-solutions and asymptotic behavior in conservation laws *J. Differ. Equ.* **51** 419–41
- [38] Serre D 2021 Source-solutions for the multi-dimensional Burgers equation *Arch. Ration. Mech. Anal.* **239** 95–116
- [39] Serre D and Silvestre L 2019 Multi-dimensional Burgers equation with unbounded initial data: well-posedness and dispersive estimates *Arch. Ration. Mech. Anal.* **234** 1391–411
- [40] Simon J 1987 Compact sets in the space $L^p(0, T; B)$ *Ann. Mat. Pura Appl.* **146** 65–96
- [41] Tartar L 2007 *An Introduction to Sobolev Spaces and Interpolation Spaces (Lecture Notes of the Unione Matematica Italiana vol 3)* (Springer)