

Theory of the Faraday-Effect:

Assumptions

Theory of the Faraday-Effect:

Assumptions

- ▶ There are microscopic magnetic moments in every transparent and nonconductive medium in an atomic scale

Theory of the Faraday-Effect:

Assumptions

- ▶ There are microscopic magnetic moments in every transparent and nonconductive medium in an atomic scale

Classical: Electrons orbiting the nucleus of the atom

QM: The total angular momentum \vec{J} of Electrons

Theory of the Faraday-Effect:

Assumptions

- ▶ There are microscopic magnetic moments in every transparent and nonconductive medium in an atomic scale

Classical: Electrons orbiting the nucleus of the atom

QM: The total angular momentum \vec{J} of Electrons

- ▶ Linear polarized light is a transverse electromagnetic wave (homogeneous Maxwell-Equations)

Derivation of the rotation angle β

- Gyroscopic equation:

$$\dot{\vec{\mu}} = \gamma \cdot \vec{\mu} \times \vec{\mathbf{B}} \quad (1)$$

Derivation of the rotation angle β

- Gyroscopic equation:

$$\dot{\vec{\mu}} = \gamma \cdot \vec{\mu} \times \vec{\mathbf{B}} \quad (1)$$

⇒ Precession of the magnetical moments:

$$\omega_L = \Delta E_{m_J} / \hbar = g \frac{q}{2m} B \quad (B = |\vec{\mathbf{B}}|) \quad (2)$$

Derivation of the rotation angle β

- Gyroscopic equation:

$$\dot{\vec{\mu}} = \gamma \cdot \vec{\mu} \times \vec{\mathbf{B}} \quad (1)$$

⇒ Precession of the magnetical moments:

$$\omega_L = \Delta E_{m_J} / \hbar = g \frac{q}{2m} B \quad (B = |\vec{\mathbf{B}}|) \quad (2)$$

$\omega_L \equiv$ precession frequency of $\vec{\mu}$ ('**Larmor frequency**')
(QM : $\gamma = g \frac{q}{2m}$, g the Lande-Faktor; Classically: $g = 1$)

Derivation of the rotation angle β

- Gyroscopic equation:

$$\dot{\vec{\mu}} = \gamma \cdot \vec{\mu} \times \vec{\mathbf{B}} \quad (1)$$

⇒ Precession of the magnetical moments:

$$\omega_L = \Delta E_{m_J} / \hbar = g \frac{q}{2m} B \quad (B = |\vec{\mathbf{B}}|) \quad (2)$$

$\omega_L \equiv$ precession frequency of $\vec{\mu}$ ('**Larmor frequency**')
(QM : $\gamma = g \frac{q}{2m}$, g the Lande-Faktor; Classically: $g = 1$)

- Dispersion relation:

$$\omega = k'(\omega) \cdot c = k \cdot n(\omega) \cdot c \quad (3)$$

Derivation of the rotation angle β

- Gyroscopic equation:

$$\dot{\vec{\mu}} = \gamma \cdot \vec{\mu} \times \vec{\mathbf{B}} \quad (1)$$

⇒ Precession of the magnetical moments:

$$\omega_L = \Delta E_{m_J} / \hbar = g \frac{q}{2m} B \quad (B = |\vec{\mathbf{B}}|) \quad (2)$$

$\omega_L \equiv$ precession frequency of $\vec{\mu}$ ('**Larmor frequency**')
(QM : $\gamma = g \frac{q}{2m}$, g the Lande-Faktor; Classically: $g = 1$)

- Dispersion relation:

$$\omega = k'(\omega) \cdot c = k \cdot n(\omega) \cdot c \quad (3)$$

- In the complex plane (light is propagating perpendicular to it):

$$\mathbf{E} = E_0 \cdot [e^{i(k'_l x - \omega t)} + e^{-i(k'_r x - \omega t)}] = \mathbf{E}_l + \mathbf{E}_r \quad (4)$$

► **perspective of the electron:**

► **perspective of the electron:**

electrons precessing counterclockwise around $\vec{\mathbf{B}}$ with the magnetical moments:

$k'_l = k'(\omega + \omega_L)$ for a left-handed circular wave and

$k'_r = k'(\omega - \omega_L)$ for a right-handed circular wave

► **perspective of the electron:**

electrons precessing counterclockwise around $\vec{\mathbf{B}}$ with the magnetical moments:

$k'_l = k'(\omega + \omega_L)$ for a left-handed circular wave and

$k'_r = k'(\omega - \omega_L)$ for a right-handed circular wave

► Assuming $\omega_L \ll \omega$

$$\Rightarrow k'_\pm := k'(\omega \pm \omega_L) \approx k'(\omega) \pm \frac{dk'}{d\omega}(\omega) \cdot \omega_L := k' \pm \frac{dk'}{d\omega} \omega_L$$

► **perspective of the electron:**

electrons precessing counterclockwise around \vec{B} with the magnetical moments:

$k'_l = k'(\omega + \omega_L)$ for a left-handed circular wave and

$k'_r = k'(\omega - \omega_L)$ for a right-handed circular wave

► Assuming $\omega_L \ll \omega$

$$\Rightarrow k'_\pm := k'(\omega \pm \omega_L) \approx k'(\omega) \pm \frac{dk'}{d\omega}(\omega) \cdot \omega_L := k' \pm \frac{dk'}{d\omega} \omega_L$$

► combined with (4) one obtains:

$$\mathbf{E} = E_0 \cdot [e^{i \frac{dk'}{d\omega} \omega_L \cdot x} (e^{i(k'x - \omega t)} + e^{-i(k'x - \omega t)})]$$

► **perspective of the electron:**

electrons precessing counterclockwise around \vec{B} with the magnetical moments:

$k'_l = k'(\omega + \omega_L)$ for a left-handed circular wave and

$k'_r = k'(\omega - \omega_L)$ for a right-handed circular wave

► Assuming $\omega_L \ll \omega$

$$\Rightarrow k'_\pm := k'(\omega \pm \omega_L) \approx k'(\omega) \pm \frac{dk'}{d\omega}(\omega) \cdot \omega_L := k' \pm \frac{dk'}{d\omega} \omega_L$$

► combined with (4) one obtains:

$$\begin{aligned} \mathbf{E} &= E_0 \cdot \left[e^{i \frac{dk'}{d\omega} \omega_L \cdot x} (e^{i(k'x - \omega t)} + e^{-i(k'x - \omega t)}) \right] \\ \Rightarrow \beta &= \frac{dk'}{d\omega} \omega_L \cdot \Delta x = \frac{dk'}{d\omega} \frac{e}{2m_e} B \cdot \Delta x \end{aligned}$$

($\Delta x \equiv$ length of the interval the light has travelled in the medium
 $g = 1$ und $q = e$ in (2) for simplicity)

- Using the chainrule and the dispersion relation:

$$\frac{dk'}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \frac{d\lambda}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \cdot \frac{-\lambda^2}{2\pi c} = -\frac{\lambda}{c} \cdot \frac{dn}{d\lambda} \quad (5)$$

- Using the chainrule and the dispersion relation:

$$\frac{dk'}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \frac{d\lambda}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \cdot \frac{-\lambda^2}{2\pi c} = -\frac{\lambda}{c} \cdot \frac{dn}{d\lambda} \quad (5)$$

$$\Rightarrow \beta = -\frac{\lambda}{c} \frac{dn}{d\lambda} \frac{e}{2m_e} \cdot \Delta x \cdot B = \Delta x \cdot V \cdot B \quad (6)$$

where V is the Verdet constant of the medium

- Using the chainrule and the dispersion relation:

$$\frac{dk'}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \frac{d\lambda}{d\omega} = \frac{2\pi}{\lambda} \cdot \frac{dn}{d\lambda} \cdot \frac{-\lambda^2}{2\pi c} = -\frac{\lambda}{c} \cdot \frac{dn}{d\lambda} \quad (5)$$

$$\Rightarrow \beta = -\frac{\lambda}{c} \frac{dn}{d\lambda} \frac{e}{2m_e} \cdot \Delta x \cdot B = \Delta x \cdot V \cdot B \quad (6)$$

where V is the Verdet constant of the medium

- For non homogeneous **B**-field we let $\Delta x \rightarrow 0$ to get the more general formula for β

$$\beta_{l_1 \rightarrow l_2} = V \cdot \int_{l_1}^{l_2} B dx \quad (7)$$

Comments to characteristic features of the Faraday-Effect

- ▶ With a strong **B**-Field we know $g > 1$, so the precession frequency will be bigger than the classical Larmor frequency

Comments to characteristic features of the Faraday-Effect

- ▶ With a strong **B**-Field we know $g > 1$, so the precession frequency will be bigger than the classical Larmor frequency
- ▶ The Verdet constant depends on the temperature of the medium ($\frac{dn}{d\lambda}(T)$ in the area of small variations negligible)

Comments to characteristic features of the Faraday-Effect

- ▶ With a strong **B**-Field we know $g > 1$, so the precession frequency will be bigger than the classical Larmor frequency
- ▶ The Verdet constant depends on the temperature of the medium ($\frac{dn}{d\lambda}(T)$ in the area of small variations negligible)
- ▶ β is independent on the direction of propagation of the light.

Comments to characteristic features of the Faraday-Effect

- ▶ With a strong **B**-Field we know $g > 1$, so the precession frequency will be bigger than the classical Larmor frequency
- ▶ The Verdet constant depends on the temperature of the medium ($\frac{dn}{d\lambda}(T)$ in the area of small variations negligible)
- ▶ β is independent on the direction of propagation of the light.

Left/right-handed circular polarized light changes to right/left-handed circular polarized light by changing the direction of propagation and you see, that eq. (4) is invariant under the transformation

$$\begin{pmatrix} k_r \\ k_l \\ \omega \end{pmatrix} \rightarrow \begin{pmatrix} -k_l \\ -k_r \\ -\omega \end{pmatrix} \quad (8)$$

Comments to characteristic features of the Faraday-Effect

- ▶ With a strong **B**-Field we know $g > 1$, so the precession frequency will be bigger than the classical Larmor frequency
- ▶ The Verdet constant depends on the temperature of the medium ($\frac{dn}{d\lambda}(T)$ in the area of small variations negligible)
- ▶ β is independent on the direction of propagation of the light.

Left/right-handed circular polarized light changes to right/left-handed circular polarized light by changing the direction of propagation and you see, that eq. (4) is invariant under the transformation

$$\begin{pmatrix} k_r \\ k_l \\ \omega \end{pmatrix} \rightarrow \begin{pmatrix} -k_l \\ -k_r \\ -\omega \end{pmatrix} \quad (8)$$

As a consequence, we obtain $\beta = \beta'$
(the specific difference from the Faraday-Effect to other double refraction phenomena)