

This is a part of SFQED-Loops script collection
developed for calculating loop processes in
Strong-Field Quantum Electrodynamics.
The scripts are available on <https://github.com/ArsenyMironov/SFQED-Loops>

If you use this script in your research, please, consider
citing our papers:

- A. A. Mironov, S. Meuren, and A. M. Fedotov, PRD 102, 053005 (2020),
<https://doi.org/10.1103/PhysRevD.102.053005>

- A. A. Mironov, A. M. Fedotov, arXiv:2109.00634 (2021)

If you have any questions, please, don't hesitate to contact:
mironov.hep@gmail.com

Copyright (C) 2021 Arseny Mironov, <https://scholar.google.ru/citations?user=GQwNEuwAAAAJ>

This program is free software: you can redistribute it and/or modify
it under the terms of the GNU General Public License as published by
the Free Software Foundation, either version 3 of the License, or
(at your option) any later version.

This program is distributed in the hope that it will be useful,
but WITHOUT ANY WARRANTY; without even the implied warranty of
MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the
GNU General Public License for more details.

You should have received a copy of the GNU General Public License
along with this program. If not, see <https://www.gnu.org/licenses/>.

$$x_1 \equiv x_2$$

```
In[1]:= NotebookEvaluate[NotebookOpen[NotebookDirectory[] <> "definitions.nb"]]
```

FeynCalc 9.3.1 (stable version). For help, use the documentation center, check out the wiki or visit the forum.

To save your and our time, please check our FAQ for answers to some common FeynCalc questions. See also the supplied examples. If you use FeynCalc in your research, please cite

- V. Shtabovenko, R. Mertig and F. Orellana, Comput.Phys.Commun. 256 (2020) 107478, arXiv:2001.04407.
- V. Shtabovenko, R. Mertig and F. Orellana, Comput.Phys.Commun. 207 (2016) 432–444, arXiv:1601.01167.
- R. Mertig, M. Böhm, and A. Denner, Comput. Phys. Commun. 64 (1991) 345–359.

Electron propagator (proper time is dimensionless)

$$S^c(x_2, x_1) = \Lambda^{4-D} \int \frac{d^D p}{(2\pi)^D} E_p(x_2) \frac{i((\gamma p) + m)}{p^2 - m^2 + i0} \bar{E}_p(x_1)$$

$$x = x_2 - x_1,$$

$$X = \frac{1}{2}(x_1 + x_2),$$

$$\xi^2 = -\frac{e^2 a^2}{m^2},$$

$$[\Lambda] = m - \text{mass scale},$$

$$E_p(x_2) = \left[1 - \frac{e(\gamma k)(\gamma a)}{2(kp)}(kx_2) \right] \text{Exp} \left[-i(p \cdot x_2) + i \frac{e(a \cdot p)}{2(k \cdot p)}(k \cdot x_2)^2 + i \frac{a^2 e^2}{6(k \cdot p)}(k \cdot x_2)^3 \right];$$

$$(\gamma p - m) S^c(p) = i - \text{in E - p representation}$$

$$S^c(p, q) = (2\pi)^4 \delta(p - q) S^c(p)$$

```
In[2]:= NewMomentum["p"]
NewCoordinate["x1"]
NewCoordinate["x2"]
NewCoordinate["x"]
NewCoordinate["X"]
```

$$\begin{aligned}
& \left\{ p^\alpha, p^2, k \cdot p, Fp^\alpha, FFp^\alpha, FDP^\alpha, a \cdot p, 0, 0, 0, -a^2 (k \cdot p), 0, 0, -\frac{m^6 \chi p^2}{e^2}, -\frac{m^6 \chi p^2}{e^2}, \frac{m^6 \chi p^2}{e^2}, 0, 0, 0, 0, 0, 0 \right\} \\
& \left\{ x1^\alpha, x1^2, k \cdot x1, a \cdot x1, Fx1^\alpha, FFx1^\alpha, FDX1^\alpha, k \cdot x1, 0, 0, 0, -a^2 (k \cdot x1), \right. \\
& \quad \left. 0, 0, -\frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}, -\frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}, \frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}, 0, 0, 0, 0, 0, 0 \right\} \\
& \left\{ x2^\alpha, x2^2, k \cdot x2, a \cdot x2, Fx2^\alpha, FFx2^\alpha, FDX2^\alpha, k \cdot x2, 0, 0, 0, -a^2 (k \cdot x2), \right. \\
& \quad \left. 0, 0, -\frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}, -\frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}, \frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}, 0, 0, 0, 0, 0, 0 \right\} \\
& \left\{ x^\alpha, x^2, k \cdot x, a \cdot x, Fx^\alpha, FFx^\alpha, FDX^\alpha, k \cdot x, 0, 0, 0, -a^2 (k \cdot x), \right. \\
& \quad \left. 0, 0, -\frac{m^2 \xi^2 (k \cdot x)^2}{e^2}, -\frac{m^2 \xi^2 (k \cdot x)^2}{e^2}, \frac{m^2 \xi^2 (k \cdot x)^2}{e^2}, 0, 0, 0, 0, 0, 0 \right\} \\
& \left\{ X^\alpha, X^2, k \cdot X, a \cdot X, FX^\alpha, FFX^\alpha, FDX^\alpha, k \cdot X, 0, 0, 0, -a^2 (k \cdot X), \right. \\
& \quad \left. 0, 0, -\frac{m^2 \xi^2 (k \cdot X)^2}{e^2}, -\frac{m^2 \xi^2 (k \cdot X)^2}{e^2}, \frac{m^2 \xi^2 (k \cdot X)^2}{e^2}, 0, 0, 0, 0, 0, 0 \right\}
\end{aligned}$$

In[7]:= **EpX2 = Ep[x2, p]**

EpBarx1 = EpC[x1, p]

$$\text{Out[7]} = \left\{ 1 - \frac{e(k \cdot x2)(\gamma \cdot k)(\gamma \cdot a)}{2(k \cdot p)}, \frac{a^2 e^2 (k \cdot x2)^3}{6(k \cdot p)} + \frac{e(a \cdot p)(k \cdot x2)^2}{2(k \cdot p)} - p \cdot x2 \right\}$$

$$\text{Out[8]} = \left\{ 1 - \frac{e(k \cdot x1)(\gamma \cdot a)(\gamma \cdot k)}{2(k \cdot p)}, -\frac{a^2 e^2 (k \cdot x1)^3}{6(k \cdot p)} - \frac{e(a \cdot p)(k \cdot x1)^2}{2(k \cdot p)} + p \cdot x1 \right\}$$

In[9]:= **Matrix = EpX2[[1]].(GAD[α] × Pair[Momentum[p, D], LorentzIndex[α, D]] + m).EpBarx1[[1]]**

Coeff = i Λ^(4-D) / (2 π)^D / (pv2 - m^2)

Phase = EpX2[[2]] + EpBarx1[[2]]

$$\text{Out[9]} = \left(1 - \frac{e(k \cdot x2)(\gamma \cdot k)(\gamma \cdot a)}{2(k \cdot p)} \right) (m + \gamma^\alpha p^\alpha) \cdot \left(1 - \frac{e(k \cdot x1)(\gamma \cdot a)(\gamma \cdot k)}{2(k \cdot p)} \right)$$

$$\text{Out[10]} = \frac{i (2\pi)^{-D} \Lambda^{4-D}}{p^2 - m^2}$$

$$\text{Out[11]} = -\frac{a^2 e^2 (k \cdot x1)^3}{6(k \cdot p)} + \frac{a^2 e^2 (k \cdot x2)^3}{6(k \cdot p)} - \frac{e(a \cdot p)(k \cdot x1)^2}{2(k \cdot p)} + \frac{e(a \cdot p)(k \cdot x2)^2}{2(k \cdot p)} + p \cdot x1 - p \cdot x2$$

In[12]:= **Matrix1 = Contract[DiracSimplify[Matrix]]**

Coeff1 = Coeff;

Phase1 = Phase;

$$\text{Out[12]} = -\frac{a^2 e^2 \gamma \cdot k (k \cdot x1) (k \cdot x2)}{2 (k \cdot p)} - \frac{e m (k \cdot x1) (\gamma \cdot a) (\gamma \cdot k)}{2 (k \cdot p)} -$$

$$\frac{e m (k \cdot x2) (\gamma \cdot k) (\gamma \cdot a)}{2 (k \cdot p)} - \frac{e (k \cdot x1) (\gamma \cdot p) (\gamma \cdot a) (\gamma \cdot k)}{2 (k \cdot p)} - \frac{e (k \cdot x2) (\gamma \cdot k) (\gamma \cdot a) (\gamma \cdot p)}{2 (k \cdot p)} + m + \gamma \cdot p$$

In[15]:= **Matrix2 =**

Expand[ExpandScalarProduct[Matrix1 /. {Momentum[x1, D] → Momentum[X, D] - Momentum[x, D] / 2, Momentum[x2, D] → Momentum[X, D] + Momentum[x, D] / 2}]]

Coeff2 = Coeff1;

Phase2 =

Expand[ExpandScalarProduct[Phase1 /. {Momentum[x1, D] → Momentum[X, D] - Momentum[x, D] / 2, Momentum[x2, D] → Momentum[X, D] + Momentum[x, D] / 2}]]

$$\text{Out[15]} = \frac{a^2 e^2 \gamma \cdot k (k \cdot x)^2}{8 (k \cdot p)} - \frac{a^2 e^2 \gamma \cdot k (k \cdot X)^2}{2 (k \cdot p)} + \frac{e m (k \cdot x) (\gamma \cdot a) (\gamma \cdot k)}{4 (k \cdot p)} - \frac{e m (k \cdot x) (\gamma \cdot k) (\gamma \cdot a)}{4 (k \cdot p)} -$$

$$\frac{e m (k \cdot X) (\gamma \cdot a) (\gamma \cdot k)}{2 (k \cdot p)} - \frac{e m (k \cdot X) (\gamma \cdot k) (\gamma \cdot a)}{2 (k \cdot p)} - \frac{e (k \cdot x) (\gamma \cdot k) (\gamma \cdot a) (\gamma \cdot p)}{4 (k \cdot p)} +$$

$$\frac{e (k \cdot x) (\gamma \cdot p) (\gamma \cdot a) (\gamma \cdot k)}{4 (k \cdot p)} - \frac{e (k \cdot X) (\gamma \cdot k) (\gamma \cdot a) (\gamma \cdot p)}{2 (k \cdot p)} - \frac{e (k \cdot X) (\gamma \cdot p) (\gamma \cdot a) (\gamma \cdot k)}{2 (k \cdot p)} + m + \gamma \cdot p$$

$$\text{Out[17]} = \frac{a^2 e^2 (k \cdot x) (k \cdot X)^2}{2 (k \cdot p)} + \frac{a^2 e^2 (k \cdot x)^3}{24 (k \cdot p)} + \frac{e (a \cdot p) (k \cdot x) (k \cdot X)}{k \cdot p} - p \cdot x$$

In[18]:= **Matrix3 = (((Expand[Matrix2 /. TripleGamma]) /. EpsToF) /. FieldSubstitutions) /. akToF /.**

{DiracGamma[LorentzIndex[β, D], D].DiracGamma[5] ×

Pair[LorentzIndex[β, D], Momentum[FDp, D]] → FVD[γγ5FD, α] × pv[α]}

Coeff3 = Coeff2;

Phase3 = Phase2;

$$\text{Out[18]} = \frac{a^2 e^2 \gamma \cdot k (k \cdot x)^2}{8 (k \cdot p)} - \frac{a^2 e^2 \gamma \cdot k (k \cdot X)^2}{2 (k \cdot p)} - \frac{e (a \cdot p) \gamma \cdot k (k \cdot X)}{k \cdot p} +$$

$$e \gamma \cdot a (k \cdot X) + \frac{i e m \sigma F (k \cdot x)}{4 (k \cdot p)} + \frac{i e \gamma \gamma 5 F D^\alpha p^\alpha (k \cdot x)}{2 (k \cdot p)} + m + \gamma \cdot p$$

Expanding scalar products into components and changing variables

$$p \rightarrow \{p_- = 1/2 (p^0 - p^3), p_+ = p^0 + p^3, p_\perp\}$$

$$p_- = x_- / 2 s$$

$$p_+ = (p^2 - p_\perp^2) / 2 p_- = s (p^2 + p_\perp^2) / x_-$$

Integration measure

$$\int d^D p \dots = \int \frac{ds}{2s} dp^2 d^{D-2} p_\perp$$

$$x_- = kx / m$$

$$p_- = kx / 2 m s$$

$$kp = m p_- = m xm / 2 s = kx / 2 s$$

$$ap = -at pt$$

$$\gamma p = \gamma_- p_+ + \gamma_+ p_- - \gamma_\perp p_\perp = Gm \frac{s}{x_-} (p^2 + p_\perp^2) + Gp \frac{x_-}{2s} - Gt pt$$

$$\gamma k = \gamma_- k_+ = m Gm$$

$$kx = k_+ x_- = m xm$$

$$(\gamma F^*)^\mu \cdot \gamma^5 \rightarrow \{(\gamma F^*)_- \cdot \gamma^5 = 0, (\gamma F^*)_+ \cdot \gamma^5, (\gamma F^*)_\perp \cdot \gamma^5\} = \{0, \gamma\gamma 5FDp, \gamma\gamma 5FDt\}$$

$$(\gamma F^*)_\mu k^\mu = 0 \rightarrow (\gamma F^*)_- = 0$$

$$(\gamma F^*)_\mu a^\mu = 0 \rightarrow \gamma\gamma 5FDt at = 0$$

$$(\gamma F^*)^\mu \cdot \gamma^5 p_\mu = \gamma\gamma 5FDm * pp - \gamma\gamma 5FDt * pt$$

In[21]:= D[{xm / 2 / s, s (p2 + pt2) / xm}, {{s, p2}}]

Abs[Det[%]]

$$\text{Out[21]=} \begin{pmatrix} -\frac{xm}{2s^2} & 0 \\ \frac{p2+pt2}{xm} & \frac{s}{xm} \end{pmatrix}$$

$$\text{Out[22]=} \frac{1}{2 |s|}$$

```
In[23]:= Matrix4 =
  Collect[Expand[Matrix3 /. {DiracGamma[Momentum[p, D], D] → Gp * pm + Gm * pp - Gt * pt, Pair[
    Momentum[a, D], Momentum[p, D]] → -at pt, FVD[γγ5FD, α_] * pv[α_] →
    γγ5FDp * pm - γγ5FDt * pt, DiracGamma[Momentum[k, D], D] → m Gm} /.
    {kp → kx / 2 / s, pm → xm / 2 / s} /. {kx → m xm} /. {pp → s (p2 + pt^2) / xm}], {pt, p2}]
```

```
Coeff4 = Coeff3 / 2 / s
```

```
Phase4 =
```

```
Collect[Expand[Phase3 /. {Pair[Momentum[p, D], Momentum[x, D]] → pp xm + pm xp - pt * xt,
  kp → m pm, Pair[Momentum[a, D], Momentum[p, D]] → -at pt} /.
  {kp → kx / 2 / s, pm → xm / 2 / s} /. {pp → s (p2 + pt^2) / xm} /. {xm → kx / m}], {pt, pp}]
```

$$\text{Out[23]} = -\frac{a^2 e^2 G_m s (k \cdot X)^2}{x m} + \frac{1}{4} a^2 e^2 G_m m^2 s x m + e \gamma \cdot a (k \cdot X) + p t \left(\frac{2 a t e G_m s (k \cdot X)}{x m} - i \gamma \gamma 5 F D t e s - G t \right) +$$

$$\frac{1}{2} i e m s \sigma F + \frac{1}{2} i \gamma \gamma 5 F D p e x m + \frac{G_m p^2 s}{x m} + \frac{G_m p t^2 s}{x m} + \frac{G_p x m}{2 s} + m$$

$$\text{Out[24]} = \frac{i 2^{-D-1} \pi^{-D} \Lambda^{4-D}}{s(p^2 - m^2)}$$

$$\text{Out[25]} = \frac{1}{12} a^2 e^2 s (k \cdot x)^2 + a^2 e^2 s (k \cdot X)^2 + p t (x t - 2 a t e s (k \cdot X)) - \frac{x p (k \cdot x)}{2 m s} - p^2 s + p t^2 (-s)$$

Integration over

$$\int d^{D-2} p_{\perp} \dots$$

$$\begin{aligned} I_0 &= \int d^{D-2} p_{\perp} \text{Exp}[-I A p_{\perp}^2 + I (J \cdot p_{\perp})] = \\ &= \text{Exp}\left[-I \frac{\pi}{2} \frac{D-2}{2}\right] \pi^{\frac{D-2}{2}} (\det A)^{-\frac{1}{2}} \text{Exp}\left[I \frac{1}{4} J \cdot A^{-1} \cdot J\right] \end{aligned}$$

$$\begin{aligned} I_1 &= \int d^{D-2} p_{\perp} p_{\perp} \text{Exp}[-I A p_{\perp}^2 + I (J \cdot p_{\perp})] = \\ &= \frac{1}{2} A^{-1} \cdot J I_0 \end{aligned}$$

$$\begin{aligned} I_2 &= \int d^{D-2} p_{\perp} p_{\perp}^2 \text{Exp}[-I A p_{\perp}^2 + I (J \cdot p_{\perp})] = \\ &= \left[-i \frac{1}{2} \text{Tr} A^{-1} + \left(\frac{1}{2} A^{-1} \cdot J\right)^2\right] I_0 \end{aligned}$$

where

$$A = s,$$

$$J = x_{\perp} - 2 e a_{\perp} s k X,$$

$$\det A = s^{D-2},$$

$$A^{-1} = 1/s$$

We perform integrations

Integrations changes the coefficient (Coeff) and phase

Then recollect some scalar products

$$a_{\perp}^2 = -a^2$$

$$a_{\perp} x_{\perp} = -(ax)$$

$$x_{\perp}^2 = 2 x_{-} x_{+} - x^2$$

```
In[26]:= Amatr = -Coefficient[Phase4, pt^2]
J = Coefficient[Phase4, pt]
CI0 = Exp[-I Pi / 2 (D / 2 - 1)] Pi^(D / 2 - 1) / Amatr^(D / 2 - 1)

Out[26]= s

Out[27]= xt - 2 at es(k · X)

Out[28]= e^(-1/2 i pi (D/2 - 1)) pi^(D/2 - 1) s^(1 - D/2)
```

```
In[29]:= Phase5 = Expand[Expand[((Phase4 /. {pt -> 0}) + 1 / 4 J^2 / Amatr)] /.
  {at^2 -> -av2, at xt -> -ax, xt^2 -> 2 xm xp - xv2} /. {xm -> kx / m}]
Coeff5 = Coeff4 * CI0
Matrix5 =
  Collect[Expand[Expand[(Matrix4 /. {pt -> 0}) + Coefficient[Matrix4, pt] * 1 / 2 / Amatr * J +
    Coefficient[Matrix4, pt^2] * (-I / 2 * (D - 2) / Amatr + (1 / 2 / Amatr * J)^2)] /.
    {γγ5FDtat -> 0, at^2 -> -av2, at xt -> -ax, xt^2 -> 2 xm xp - xv2}],
    {p2, Gm, Gp, Gt, γγ5FDm, γγ5FDp, γγ5FDt}]
```

$$\text{Out[29]} = \frac{1}{12} a^2 e^2 s (k \cdot x)^2 + e(a \cdot x)(k \cdot X) - p_2 s - \frac{x^2}{4 s}$$

$$\text{Out[30]} = \frac{i 2^{-D-1} e^{-\frac{1}{2} i \pi \left(\frac{D}{2}-1\right)} \pi^{-\frac{D}{2}-1} \Lambda^{4-D} s^{-D/2}}{p^2 - m^2}$$

$$\begin{aligned} \text{Out[31]} = & Gm \left(\frac{1}{4} a^2 e^2 m^2 s xm - \frac{i D}{2 xm} - \frac{x^2}{4 s xm} + \frac{xp}{2 s} + \frac{i}{xm} \right) + e \gamma \cdot a (k \cdot X) + Gt \left(at e (k \cdot X) - \frac{xt}{2 s} \right) + \\ & \frac{1}{2} i e m s \sigma F + \frac{1}{2} i \gamma \gamma 5 F D p e xm - \frac{1}{2} i \gamma \gamma 5 F D t e xt + \frac{Gm p_2 s}{xm} + \frac{Gp xm}{2 s} + m \end{aligned}$$

Next we substitute

$$\gamma_{\perp} a_{\perp} = -\gamma a$$

$$\gamma_{\perp} x_{\perp} = \gamma_{-} x_{+} - \gamma_{+} x_{-} - \gamma x$$

$$\gamma_{-} x_{-} = \frac{\cancel{\gamma k}}{m} x_{-}$$

$$x_{-} = kx / m$$


```
In[32]:= Matrix6 = Collect[Expand[Matrix5] /.
  {Gt at → -Contract[GAD[α] × av[α]], Gt xt → Gm xp + Gp xm - Contract[GAD[α] × xv[α]] ,
  γγ5FDtxt → γγ5FDp xm - DiracSlash[FDxv, Dimension → D].GA[5]} /.
  {Gm xm → DiracGamma[Momentum[k, D], D] / m * xm} /. {xm → kx / m},
  {p2, Gm, Gp, Gt, γγ5FDm, γγ5FDp, γγ5FDt}]
Coeff6 = Coeff5
Phase6 = Phase5
```

$$\text{Out[32]} = \frac{1}{2} i e (\gamma \cdot \text{FDxv}) \cdot \gamma^5 + \frac{1}{4} a^2 e^2 s \gamma \cdot k (k \cdot x) +$$

$$Gm \left(-\frac{i D m}{2 (k \cdot x)} - \frac{m x^2}{4 s (k \cdot x)} + \frac{i m}{k \cdot x} \right) + \frac{1}{2} i e m s \sigma F + \frac{Gm m p^2 s}{k \cdot x} + m + \frac{\gamma \cdot x}{2 s}$$

$$\text{Out[33]} = \frac{i 2^{-D-1} e^{-\frac{1}{2} i \pi \left(\frac{D}{2} - 1 \right)} \pi^{-\frac{D}{2} - 1} \Lambda^{4-D} s^{-D/2}}{p^2 - m^2}$$

$$\text{Out[34]} = \frac{1}{12} a^2 e^2 s (k \cdot x)^2 + e(a \cdot x) (k \cdot X) - p^2 s - \frac{x^2}{4 s}$$

$$v = p^2 - m^2$$

$$\int \frac{dv}{v + i0} \exp[-i s v] = -2 \pi i \theta(s)$$

$$\int dv \exp[-i s v] = 2 \pi \delta(s)$$

$$\int_{-\infty}^{\infty} \frac{ds}{s^{D/2}} s e^{i g(s)} 2 \pi \delta(s) = -2 \pi i \int_{-\infty}^{\infty} \frac{ds}{s^{D/2}} e^{i g(s)} \theta(s) [i (D/2 - 1) + s g'(s)]$$

where

$$g(s) = \text{Phase6 with } p^2 = m^2$$

Finally,

$$Gm = \gamma_- = \gamma k / m$$

```

In[35]:= Phase7 = Phase6 /. {p2 → m^2}
Coeff7 = Coeff6 * (pv2 - m^2) * (-2 π i)
g = Phase7;
Matrix7 =
  Expand[(Matrix6 /. {p2 → m^2}) + Coefficient[Matrix6, p2 s] * (i (D/2 - 1) + s D[g, s])] /.
    {Gm → DiracGamma[Momentum[k, D], D] / m}

Out[35]=  $\frac{1}{12} a^2 e^2 s (k \cdot x)^2 + e(a \cdot x) (k \cdot X) + m^2 (-s) - \frac{x^2}{4 s}$ 

Out[36]=  $2^{-D} e^{-\frac{1}{2} i \pi \left(\frac{D}{2}-1\right)} \pi^{-D/2} \Lambda^{4-D} s^{-D/2}$ 

Out[38]=  $\frac{1}{2} i e (\gamma \cdot \text{FDxv}) \cdot \bar{\gamma}^5 + \frac{1}{3} a^2 e^2 s \gamma \cdot k (k \cdot x) + \frac{1}{2} i e m s \sigma F + m + \frac{\gamma \cdot x}{2 s}$ 

In[39]:= Matrix8 =
  Expand[Matrix7 / m /. {DiracGamma[Momentum[k, D], D] → Contract[GAD[α] × FFxv[α]] / (-av2 kx)}]
Coeff8 = Coeff7 * m
Phase8 = Phase7 /. {av2 kx^2 → Fx^2}

Out[39]=  $\frac{i e (\gamma \cdot \text{FDxv}) \cdot \bar{\gamma}^5}{2 m} - \frac{e^2 s \gamma \cdot \text{FFx}}{3 m} + \frac{1}{2} i e s \sigma F + \frac{\gamma \cdot x}{2 m s} + 1$ 

Out[40]=  $2^{-D} e^{-\frac{1}{2} i \pi \left(\frac{D}{2}-1\right)} \pi^{-D/2} m \Lambda^{4-D} s^{-D/2}$ 

Out[41]=  $e(a \cdot x) (k \cdot X) + \frac{1}{12} e^2 Fx^2 s - m^2 s - \frac{x^2}{4 s}$ 

In[42]:= Matrix9 = Matrix8 /. {s → s1 / m^2}
Coeff9 = Simplify[Coeff8 / m^2 /. {s → s1 / m^2}, Assumptions → m > 0]
Coeff9 /. {D → 4}
Phase9 = Phase8 /. {s → s1 / m^2}

Out[42]=  $\frac{i e (\gamma \cdot \text{FDxv}) \cdot \bar{\gamma}^5}{2 m} - \frac{e^2 s1 \gamma \cdot \text{FFx}}{3 m^3} + \frac{i e s1 \sigma F}{2 m^2} + \frac{m \gamma \cdot x}{2 s1} + 1$ 

Out[43]=  $i 2^{-D} e^{-\frac{1}{4} i \pi D} \pi^{-D/2} \Lambda^{4-D} m^{D-1} s1^{-D/2}$ 

Out[44]=  $-\frac{i m^3}{16 \pi^2 s1^2}$ 

Out[45]=  $e(a \cdot x) (k \cdot X) + \frac{e^2 Fx^2 s1}{12 m^2} - \frac{m^2 x^2}{4 s1} - s1$ 

```

```
In[46]:= Matrix91 = Expand[Matrix9 /. {DiracGamma[Momentum[FDxv, D], D].GA[5] →
Contract[DiracGamma[LorentzIndex[μ, D], D].GA[5] × FDxv[μ] /.
{FDxv[μ_] → -LCD[μ, α1, α2, α3] × xv[α1] × av[α2] × kv[α3]}, EpsContract → False]} /.
antiTripleGamma /. {DiracGamma[Momentum[FFx, D], D] →
-av2 kx DiracGamma[Momentum[k, D], D]} /.
{σF → -2 i DiracGamma[Momentum[a, D], D].DiracGamma[Momentum[k, D], D]}]
Out[46]= 
$$\frac{a^2 e^2 s1 \gamma \cdot k (k \cdot x)}{3 m^3} + \frac{e s1 (\gamma \cdot a) (\gamma \cdot k)}{m^2} + \frac{e (a \cdot x) \gamma \cdot k}{2 m} - \frac{e \gamma \cdot a (k \cdot x)}{2 m} + \frac{e (\gamma \cdot a) (\gamma \cdot k) (\gamma \cdot x)}{2 m} + \frac{m \gamma \cdot x}{2 s1} + 1$$

```

```
In[47]:= Expand[(Matrix91 / (m / 2 / s1) /. {s1 → s * m ^ 2}) / 2 / s]
Out[47]= 
$$\frac{1}{3} a^2 e^2 s \gamma \cdot k (k \cdot x) + e m s (\gamma \cdot a) (\gamma \cdot k) + \frac{1}{2} e (a \cdot x) \gamma \cdot k - \frac{1}{2} e \gamma \cdot a (k \cdot x) + \frac{1}{2} e (\gamma \cdot a) (\gamma \cdot k) (\gamma \cdot x) + m + \frac{\gamma \cdot x}{2 s}$$

```

Another representation of the γ – matrix preexponent term

```
In[48]:= MartixAnother = (2 m s1 * 1 / 2 / m / s1 GAD[α] (xv[α] - s1 e Fxv[α] + s1 ^ 2 / 3 e ^ 2 FFxv[α]) + 2 m s1 * 1) .
(1 + e s1 DiracSlash[a, k, Dimension → D]) /. {s1 → s1 / m ^ 2}
MartixAnother1 = DiracSimplify[Contract[DotSimplify[
MartixAnother / (2 s1 / m) /. {Fxv[α_] → kv[α] ax - av[α] kx, FFxv[α_] → -av2 kx kv[α]}]]]
CoeffAnother = Coeff9 * m / 2 / s1
Out[48]= 
$$\left( \gamma^\alpha \left( \frac{e^2 s1^2 FFx^\alpha}{3 m^4} - \frac{e s1 Fx^\alpha}{m^2} + x^\alpha \right) + \frac{2 s1}{m} \right) \left( \frac{e s1 (\gamma \cdot a) (\gamma \cdot k)}{m^2} + 1 \right)$$

Out[49]= 
$$\frac{a^2 e^2 s1 \gamma \cdot k (k \cdot x)}{3 m^3} + \frac{e s1 (\gamma \cdot a) (\gamma \cdot k)}{m^2} - \frac{e (a \cdot x) \gamma \cdot k}{2 m} + \frac{e \gamma \cdot a (k \cdot x)}{2 m} + \frac{e (\gamma \cdot x) (\gamma \cdot a) (\gamma \cdot k)}{2 m} + \frac{m \gamma \cdot x}{2 s1} + 1$$

Out[50]= 
$$i 2^{-D-1} e^{-\frac{1}{4} i \pi D} \pi^{-D/2} \Lambda^{4-D} m^D s1^{-\frac{D}{2}-1}$$

In[51]:= Matrix91 - Expand[MartixAnother1]
Out[51]= 
$$\frac{e (a \cdot x) \gamma \cdot k}{m} - \frac{e \gamma \cdot a (k \cdot x)}{m} + \frac{e (\gamma \cdot a) (\gamma \cdot k) (\gamma \cdot x)}{2 m} - \frac{e (\gamma \cdot x) (\gamma \cdot a) (\gamma \cdot k)}{2 m}$$

```

Final result for the electron propagator in a CCF

$$S^c(x_2, x_1) = \Lambda^{4-D} \int \frac{d^D p}{(2\pi)^D} E_p(x_2) \frac{i((\gamma p) + m)}{p^2 - m^2 + i0} E_p^{\text{bar}}(x_1) =$$

$$e^{-i \frac{\pi}{2} \left(\frac{D}{2} - 1 \right)} \frac{\Lambda^{4-D}}{2^D \pi^{D/2}} m^{D-1} e^{i \eta} \int_0^\infty \frac{ds}{s^{D/2}} \left[1 + \frac{m(\gamma x)}{2s} - \frac{e^2 s (\gamma FFx)}{3 m^3} + \right.$$

$$\left. \frac{i e s (\sigma^{\alpha\beta} F_{\alpha\beta})}{2 m^2} + \frac{i e F_{\alpha\beta}^* x^\beta \gamma^\alpha \gamma^5}{2 m} \right] e^{-i s - i \frac{m^2 x^2}{4 s} + i \frac{\pi}{12} \frac{e^2}{m^2} (Fx)^2} =$$

$$\begin{aligned}
&= e^{-i \frac{\pi}{2} \binom{D-1}{2}} \frac{\Lambda^{4-D}}{2^D \pi^{D/2}} m^{D-1} e^{i \eta} \int_0^\infty \frac{ds}{s^{D/2}} \left[1 + \frac{m(\gamma x)}{2s} + \frac{e(\gamma a)(kx)}{2m} - \right. \\
&\quad \frac{e(\gamma k)(ax)}{2m} + \frac{e(\gamma x)(\gamma a)(\gamma k)}{2m} + \frac{es(\gamma a)(\gamma k)}{m^2} + \\
&\quad \left. \frac{e^2 a^2 s(\gamma k)(kx)}{3m^3} \right] e^{-i s - i \frac{m^2 \gamma^2}{4s} + i \frac{\pi}{12} \frac{a^2}{m^2} (Fx)^2} = \\
&= e^{-i \frac{\pi}{2} \binom{D-1}{2}} \frac{\Lambda^{4-D}}{2^{D+1} \pi^{D/2}} m^D e^{i \eta} \int_0^\infty \frac{ds}{s^{D/2+1}} \left[\frac{2s}{m} + \gamma^\alpha \left(g_{\alpha\beta} - \frac{es}{m^2} F_{\alpha\beta} + \frac{e^2 s^2}{3m^4} F_{\alpha\lambda} F^\lambda{}_\beta \right) \right. \\
&\quad \left. x^\beta \right] \left(1 + \frac{ies}{2} \sigma^{\alpha\beta} F_{\alpha\beta} \right) e^{-i s - i \frac{m^2 \gamma^2}{4s} + i \frac{\pi}{12} \frac{a^2}{m^2} (Fx)^2}
\end{aligned}$$

$$\eta = e(ax)(k, (x_1 + x_2)/2),$$

$$x = x_2 - x_1,$$

$$e > 0,$$

$$\sigma^{\alpha\beta} = \frac{i}{2} (\gamma^\alpha \gamma^\beta - \gamma^\beta \gamma^\alpha),$$

$$\gamma^5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3,$$

$$e^{-i \frac{\pi}{2} \binom{D-1}{2}} \frac{\Lambda^{4-D}}{2^D \pi^{D/2}} m^{D-1} \rightarrow \frac{(-i)m^3}{16 \pi^2}, \quad D \rightarrow 4$$