

This is a part of SFQED-Loops script collection
developed for calculating loop and tree-level processes in
Strong-Field Quantum Electrodynamics.
The scripts are available on <https://github.com/ArsenyMironov/SFQED-Loops>

If you use this script in your research, please, consider
citing our papers:

- A. A. Mironov, S. Meuren, and A. M. Fedotov, PRD 102, 053005 (2020); arXiv:2003.06909
<https://doi.org/10.1103/PhysRevD.102.053005>
- A. A. Mironov, A. M. Fedotov, 105 (3), 033005 (2022); arXiv:2109.00634
<https://doi.org/10.1103/PhysRevD.105.033005>
- Y. V. Selivanov, A. A. Mironov, A. I. Alexeenko, A. M. Fedotov arXiv:2605.12479 (2026)

If you have any questions, please, don't hesitate to contact:
mironov.hep@gmail.com

Copyright (C) 2026 Arseny Mironov, <https://scholar.google.ru/citations?user=GQwNEuwAAAAJ>

This program is free software: you can redistribute it and/or modify
it under the terms of the GNU General Public License as published by
the Free Software Foundation, either version 3 of the License, or
(at your option) any later version.

This program is distributed in the hope that it will be useful,
but WITHOUT ANY WARRANTY; without even the implied warranty of
MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the
GNU General Public License for more details.


You should have received a copy of the GNU General Public License
along with this program. If not, see <https://www.gnu.org/licenses/>.

This notebook is a supplement for our work
[Y. V. Selivanov, A. A. Mironov, A. I. Alexeenko, A. M. Fedotov arXiv:2605.12479 (2026)]

The notebook was prepared by A. A. Mironov and Y. V. Selivanov

The notebook presents a tree-level calculation of the trident process rate (the direct term contribution) initiated by a relativistic electron propagaing in a strong constant crossed electromagnetic field, W_{dir}

The corresponding diagram considered in this notebook reads

$$\sum_{\text{spin}} \int d\Pi(q, g, f) \left| \begin{array}{c} \text{diagram} \\ \text{"direct" diagram} \end{array} \right|^2$$


The notebook depends on two additional files
 definitions.nb
 TridentDiracSimplify.nb
 which should be contained in the same directory

```
In[1]:= NotebookEvaluate[NotebookOpen[NotebookDirectory[] <> "definitions.nb"]]
```

FeynCalc 10.1.0 (stable version). For help, use the
 online documentation, visit the forum and have a look at the supplied
 examples. The PDF-version of the manual can be downloaded [here](#).

If you use FeynCalc in your research, please
 evaluate FeynCalcHowToCite[] to learn how to cite this software.

*Please keep in mind that the proper academic attribution
 of our work is crucial to ensure the future development of this package!*

Trident process ($e > 0$)

$$S_{fi}^{\text{trident, full}} = S_{fi}^{\text{dir}} + S_{fi}^{\text{ex}},$$

where S_{fi}^{dir} is the direct term and S_{fi}^{ex}
 is the exchange term. We will consider only S_{fi}^{dir}

$$\text{Note that } \int d\Pi \left| S_{fi}^{\text{dir}} \right|^2 = \int d\Pi \left| S_{fi}^{\text{ex}} \right|^2,$$

where $\int d\Pi$ is the integral over all the outgoing particle momenta

The direct term matrix element is defined by*

$$S_{fi}^{\text{dir}} = (ie)^2 \int d^4x_1 d^4x_2 \bar{\psi}_q(x_1) \gamma^\mu \psi_p(x_1) D_{\mu\nu}^c(x_1 - x_2) \bar{\psi}_f(x_2) \gamma^\nu \psi_g^+(x_2);$$

x_1^μ, x_2^μ – position of the left and right vertices of the diagram;

p^μ, q^μ – initial and final electron momenta;

$S^c, D_{\mu\nu}^c$ – electron and photon casual propagators;

Superscript + stands for Hermitian conjugation

The Ritus E_p – function

$$E_p(x_1) = \left[1 - \frac{e(\gamma k)(\gamma a)}{2(kp)} (kx_2) \right]$$

$$\text{Exp}\left[-i(\mathbf{p} \cdot \mathbf{x}_1) + i \frac{e(a \cdot \mathbf{p})}{2(k \cdot \mathbf{p})} (k \cdot \mathbf{x}_1)^2 + i \frac{e^2 a^2}{6(k \cdot \mathbf{p})} (k \cdot \mathbf{x}_1)^3\right];$$

$$\bar{E}_q(x_2) = \gamma^0 E_q(x_2) \gamma^0;$$

Volkov states

$$\psi_{p\lambda}(x_1) = E_p(x_1) u_{p\lambda}$$

$$\bar{\psi}_{p\lambda}(x_1) = \bar{u}_{p\lambda} \bar{E}_p(x_1)$$

$$\psi_{p\lambda}^+(x_1) = E_{-p}(x_1) v_{p\lambda}$$

$$\bar{u}_{p\lambda} = u_{p\lambda} \gamma^0$$

Photon propagator

$$D_{\mu\nu}^c(x) = \int \frac{d^4 l}{(2\pi)^4} e^{ilx} \frac{-i}{l^2 + i0} g_{\mu\nu}$$

* Note some differences in notations used in this

Notebook and in section 6.1 of [arXiv 2605.12479]. Here, $e > 0$ denotes the absolute value of the electron charge, and momenta f and g are interchanged. While this slightly changes some intermediate steps, this does not affect the final result.

Main

Preliminary definitions

We assume that

$\mathbf{F}$ is the crossed field tensor,

$\mathbf{k}$ is the 'wave' 4 - vector, normalised such that $k^0 = m$,

$\mathbf{a}$ is the transverse (space - like) constant 4 - vector,

χp denotes the χ parameter of a particle with 4 - momentum p ,

All objects such as vectors and tensors introduced via commands from `definitions.nb` are assumed to be D - dimensional. However, since there is no regularization involved in this calculation, we will set $D \rightarrow 4$ along the Notebook

Command **NewMomentum** is defined in `definitions.nb`. It introduces a set of 4 - vectors and scalars induced by a 4 - momenta and tensor F , see `definitions.nb` for more details

```

In[2]:= NewMomentum["p"] (*Initial electron*)
NewMomentum["q"] (*Scattered electron*)
NewMomentum["g"] (*New positron, created by a photon*)
NewMomentum["f"] (*New electron, created by a photon*)
NewMomentum["l"] (*Intermediate photon*)
NewMomentum["l1"] (*Also intermediate photon,
l1 is used when mod-sqaring the amplitude*)
NewMomentum["l2"] (*Same as above*)
NewCoordinate["x1"] (*Coordinate at which emission happens*)
NewCoordinate["x2"] (*Coordinate at which pair creation happens*)
ScalarProduct[p, p] = m^2;
ScalarProduct[q, q] = m^2;
ScalarProduct[f, f] = m^2;
ScalarProduct[g, g] = m^2;

{p^α, p^2, k · p, Fp^α, FFp^α, FDp^α, a · p, 0, 0, 0, -a^2 (k · p), 0, 0, - $\frac{m^6 \chi p^2}{e^2}$ , - $\frac{m^6 \chi p^2}{e^2}$ ,  $\frac{m^6 \chi p^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{q^α, q^2, k · q, Fq^α, FFq^α, FDq^α, a · q, 0, 0, 0, -a^2 (k · q), 0, 0, - $\frac{m^6 \chi q^2}{e^2}$ , - $\frac{m^6 \chi q^2}{e^2}$ ,  $\frac{m^6 \chi q^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{g^α, g^2, g · k, Fg^α, FFg^α, FDg^α, a · g, 0, 0, 0, -a^2 (g · k), 0, 0, - $\frac{m^6 \chi g^2}{e^2}$ , - $\frac{m^6 \chi g^2}{e^2}$ ,  $\frac{m^6 \chi g^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{f^α, f^2, f · k, Ff^α, FFf^α, FDf^α, a · f, 0, 0, 0, -a^2 (f · k), 0, 0, - $\frac{m^6 \chi f^2}{e^2}$ , - $\frac{m^6 \chi f^2}{e^2}$ ,  $\frac{m^6 \chi f^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{l^α, l^2, k · l, Fl^α, FFl^α, FDll^α, a · l, 0, 0, 0, -a^2 (k · l), 0, 0, - $\frac{m^6 \chi l^2}{e^2}$ , - $\frac{m^6 \chi l^2}{e^2}$ ,  $\frac{m^6 \chi l^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{l1^α, l1^2, k · l1, Fl1^α, FFl1^α, FDll1^α, a · l1, 0, 0, 0, -a^2 (k · l1), 0, 0, - $\frac{m^6 \chi l1^2}{e^2}$ , - $\frac{m^6 \chi l1^2}{e^2}$ ,  $\frac{m^6 \chi l1^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{l2^α, l2^2, k · l2, Fl2^α, FFl2^α, FDll2^α, a · l2, 0, 0, 0, -a^2 (k · l2), 0, 0, - $\frac{m^6 \chi l2^2}{e^2}$ , - $\frac{m^6 \chi l2^2}{e^2}$ ,  $\frac{m^6 \chi l2^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{x1^α, x1^2, k · x1, a · x1, Fx1^α, FFx1^α, FDx1^α, k · x1, 0, 0, 0,
-a^2 (k · x1), 0, 0, - $\frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}$ , - $\frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}$ ,  $\frac{m^2 \xi^2 (k \cdot x1)^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

{x2^α, x2^2, k · x2, a · x2, Fx2^α, FFx2^α, FDx2^α, k · x2, 0, 0, 0,
-a^2 (k · x2), 0, 0, - $\frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}$ , - $\frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}$ ,  $\frac{m^2 \xi^2 (k \cdot x2)^2}{e^2}$ , 0, 0, 0, 0, 0, 0}

```

Commands **Ep** and **EpC** introduce E_p –

functions in the form of an array {prefactor, phase},
see definitions.nb for details

In[15]:= **Ep**x1 = **Ep**[x1, p]

EqBarx1 = **EpC**[x1, q]

EfBarx2 = **EpC**[x2, f]

Emingx2 = **Ep**[x2, -g]

$$\text{Out[15]} = \left\{ 1 - \frac{e(k \cdot x1)(\gamma \cdot k)(\gamma \cdot a)}{2(k \cdot p)}, \frac{a^2 e^2 (k \cdot x1)^3}{6(k \cdot p)} + \frac{e(a \cdot p)(k \cdot x1)^2}{2(k \cdot p)} - p \cdot x1 \right\}$$

$$\text{Out[16]} = \left\{ 1 - \frac{e(k \cdot x1)(\gamma \cdot a)(\gamma \cdot k)}{2(k \cdot q)}, -\frac{a^2 e^2 (k \cdot x1)^3}{6(k \cdot q)} - \frac{e(a \cdot q)(k \cdot x1)^2}{2(k \cdot q)} + q \cdot x1 \right\}$$

$$\text{Out[17]} = \left\{ 1 - \frac{e(k \cdot x2)(\gamma \cdot k)(\gamma \cdot k)}{2(f \cdot k)}, -\frac{a^2 e^2 (k \cdot x2)^3}{6(f \cdot k)} - \frac{e(a \cdot f)(k \cdot x2)^2}{2(f \cdot k)} + f \cdot x2 \right\}$$

$$\text{Out[18]} = \left\{ \frac{e(k \cdot x2)(\gamma \cdot k)(\gamma \cdot a)}{2(g \cdot k)} + 1, -\frac{a^2 e^2 (k \cdot x2)^3}{6(g \cdot k)} + \frac{e(a \cdot g)(k \cdot x2)^2}{2(g \cdot k)} + g \cdot x2 \right\}$$

The dressed vertex (with an outgoing photon line)

$$\Gamma^\mu(l, q, p) = \int d^4x \bar{E}_q(x) \gamma^\mu E_p(x) e^{i l x};$$

x^μ – position of the vertex;

p^μ, q^μ – initial and final electron momenta;

l^μ, q^μ – momentum of the outgoing photon

$-\infty < s < \infty$ – measure for the momentum exchanged

with the field (may be viewed as the 'photon number')

$$\Gamma^\mu(l, q, p) = \int ds (2\pi)^4 \delta(p + sk - q - l) \tilde{\Gamma}^\mu(s, q, p)$$

$$\tilde{\Gamma}^\mu(s, q, p) = \gamma^\mu A_0(s, q, p) -$$

$$\frac{e}{2} \left(\frac{(\gamma a)(\gamma k) \gamma^\mu}{kq} + \frac{\gamma^\mu (\gamma k)(\gamma a)}{kp} \right) A_1(s, q, p) - \frac{e^2 a^2 k^\mu (\gamma k)}{2 kq kp} A_2(s, q, p)$$

$$A_0(s, q, p) = \int \frac{d\phi}{2\pi} e^{is\phi} e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} = \frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \text{Ai}(y)$$

$$A_1(s, q, p) = -i \frac{dA_0}{ds} = \int \frac{d\phi}{2\pi} e^{is\phi} \phi e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} =$$

$$\frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \left\{ \frac{\alpha}{8\beta} \text{Ai}(y) - i \frac{1}{(4\beta)^{1/3}} \text{Ai}'(y) \right\}$$

$$A_2(s, q, p) = (-i)^2 \frac{d^2 A_0}{ds^2} = \int \frac{d\phi}{2\pi} e^{is\phi} \phi^2 e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} =$$

$$\frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \left\{ \left[2\left(\frac{\alpha}{8\beta}\right)^2 - \frac{s}{4\beta}\right] \text{Ai}(y) - i \frac{\alpha}{(4\beta)^{4/3}} \text{Ai}'(y) \right\}$$

where

$$\alpha = \alpha(q, p) = -e \left(\frac{ap}{kp} - \frac{aq}{kq} \right)$$

$$\beta = \beta(q, p) = \frac{e^2 a^2}{8} \left(\frac{1}{kp} - \frac{1}{kq} \right)$$

$$y = y(s, q, p) = (4\beta(q, p))^{2/3} \left[\frac{s}{4\beta(q, p)} - \left(\frac{\alpha(q, p)}{8\beta(q, p)} \right)^2 \right]$$

The amplitude expressed via $\tilde{\Gamma}^\mu$

$$S_{fi}^{\text{dir}} = (ie)^2 \int d^4x_1 d^4x_2 \bar{\psi}_q(x_1) \gamma^\mu \psi_p(x_1) D_{\mu\nu}^c(x_1 - x_2) \bar{\psi}_f(x_2) \gamma^\nu \psi_g(x_2)$$

$$= (ie)^2 \int d^4x_1 d^4x_2 \frac{d^4l}{(2\pi)^4} \frac{-i}{l^2 + i0} g_{\mu\nu}$$

$$\left(\bar{u}_{q\lambda_2} \bar{E}_q(x_1) \gamma^\mu E_p(x_1) u_{p\lambda_1} e^{ilx_1} \right) \left(\bar{u}_{f\sigma_2} \bar{E}_f(x_2) \gamma^\mu E_g(x_2) v_{g\sigma_1} e^{-ilx_2} \right)$$

$$= ie^2 \int \frac{d^4l}{(2\pi)^4} \frac{g_{\mu\nu}}{l^2 + i0} \left(\bar{u}_{q\lambda_2} \int d^4s_1 (2\pi)^4 \delta(p + s_1 k - q - l) \tilde{\Gamma}^\mu(s_1, q, p) u_{p\lambda_1} \right)$$

$$\left(\bar{u}_{f\sigma_2} \int d^4s_2 (2\pi)^4 \delta(l + s_2 k - f - g) \tilde{\Gamma}^\mu(s_2, g, -f) v_{g\sigma_1} \right)$$

$$= ie^2 \int d^4s_1 d^4s_2 (2\pi)^4 \delta(p + (s_1 + s_2)k - q - f - g)$$

$$\frac{g_{\mu\nu}}{(p + s_1 k - q)^2 + i0} \left(\bar{u}_{q\lambda_2} \tilde{\Gamma}^\mu(s_1, q, p) u_{p\lambda_1} \right) \left(\bar{u}_{f\sigma_2} \tilde{\Gamma}^\mu(s_2, g, -f) v_{g\sigma_1} \right)$$

Let us introduce partial amplitude $m(s_1, s_2)$

(curly M in [arXiv 2605.12479], see Eq. (6.1), note that here $f \leftrightarrow g$)

$$S_{fi}^{\text{dir}} = \int ds_1 ds_2 i m(s_1, s_2) (2\pi)^4 \delta(g + f + q - p - (s_1 + s_2)k)$$

$$i m(s_1, s_2) = ie^2 \left[\bar{u}_{q\lambda_2} \tilde{\Gamma}^\mu(s_1, q, p) u_{p\lambda_1} \right] \frac{g_{\mu\nu}}{l^2 + i0} \left[\bar{u}_{f\sigma_2} \tilde{\Gamma}^\nu(s_2, f, -g) v_{g\sigma_1} \right]$$

$$m(s_1, s_2) = e^2 \frac{1}{l^2 + i0} \left[\bar{u}_{q\lambda_2} \tilde{\Gamma}^\mu(s_1, q, p) u_{p\lambda_1} \right] \left[\bar{u}_{g\sigma_2} \tilde{\Gamma}_\mu(s_2, f, -g) v_{g\sigma_1} \right]$$

$$m^+(t_1, t_2) =$$

$$e^2 \frac{1}{l'^2 - i0} \left[\bar{u}_{p\lambda_1} \left(\tilde{\Gamma}^\mu(t_1, q, p) \right)^+ u_{q\lambda_2} \right] \left[\bar{v}_{g\sigma_1} \left(\tilde{\Gamma}_\mu(t_2, f, -g) \right)^+ u_{f\sigma_2} \right]$$

$$\mathbf{l} = \mathbf{p} + \mathbf{s}_1 \mathbf{k} - \mathbf{q}$$

$$\mathbf{l}' = \mathbf{p} + \mathbf{t}_1 \mathbf{k} - \mathbf{q}$$

$$\sum u_{\mathbf{p}\lambda}^\alpha \bar{u}_{\mathbf{p}\lambda}^\beta = (\mathbf{p}\gamma + m)^{\alpha\beta}$$

$$\sum v_{\mathbf{p}\lambda}^\alpha \bar{v}_{\mathbf{p}\lambda}^\beta = (\mathbf{p}\gamma - m)^{\alpha\beta}$$

$$dW = \frac{dw}{VT 2 p^0};$$

$$\frac{1}{2} \sum_{\lambda\sigma} - \text{averaging over the initial electron spin states}$$

and summation over the final particle spin projections

$$\frac{1}{2} - \text{a symmetry factor to account for the two identical final electrons}$$

$$dw = \frac{1}{2} \times \frac{1}{2} \sum_{\lambda\sigma} \frac{d^3 f}{(2\pi)^3 2 f^0} \frac{d^3 g}{(2\pi)^3 2 g^0} \frac{d^3 q}{(2\pi)^3 2 q^0} (|S_{fi}^{dir}|)^2;$$

$$(|S_{fi}^{dir}|)^2 = \frac{VT}{L_\phi} \int ds_1 ds_2 dt_1 dt_2 M(s_1, s_2, t_1, t_2) \\ (2\pi)^4 \delta(\mathbf{g} + \mathbf{f} + \mathbf{q} - \mathbf{p} - (\mathbf{s}_1 + \mathbf{s}_2) \mathbf{k}) (2\pi) \delta(s_1 + s_2 - t_1 - t_2)$$

$$M(s_1, s_2, t_1, t_2) = m(s_1, s_2) m^+(t_1, t_2)$$

Steps 1-2: Integrations using δ -functions

Step 1 : integrate over $d^3 f$ using δ - function

$$\int \frac{d^3 f}{(2\pi)^3 2 f^0} (2\pi)^4 \delta(\mathbf{g} + \mathbf{f} + \mathbf{q} - \mathbf{p} - (\mathbf{s}_1 + \mathbf{s}_2) \mathbf{k}) = \\ \frac{2\pi}{2 f^0} \delta(g^0 + f^0 + q^0 - p^0 - (s_1 + s_2) k^0)$$

$\delta(\mathbf{g} + \mathbf{f} + \mathbf{q} - \mathbf{p} - (\mathbf{s}_1 + \mathbf{s}_2) \mathbf{k})$ also implies the conservation of :

$$\mathbf{k}\mathbf{g} + \mathbf{k}\mathbf{f} + \mathbf{k}\mathbf{q} - \mathbf{k}\mathbf{p} = 0, \quad \mathbf{g}_\perp + \mathbf{f}_\perp + \mathbf{q}_\perp - \mathbf{p}_\perp = 0$$

$$\text{Consider } g^0 + f^0 + q^0 - p^0 - (s_1 + s_2) k^0 = 0$$

By substituting $f^0 = \sqrt{m^2 + \left(\vec{f}\right)^2} = \sqrt{m^2 + \left(\vec{p} + \vec{s} \mathbf{k} - \vec{g} - \vec{q}\right)^2}$, we get

$$\sqrt{m^2 + \left(\vec{p} + (s_1 + s_2) \vec{k} - \vec{g} - \vec{q}\right)^2} = p^0 + (s_1 + s_2) k^0 - g^0 - q^0$$

If we square this expression and rearrange terms, we will arrive at

$$s_1 + s_2 = \tilde{s} = \frac{(pq + pg - qg - m^2)}{kp - kq - kg} = \frac{(pq + pg - qg - m^2)}{kf}$$

Hence,

$$\delta (g^0 + f^0 + q^0 - p^0 - (s_1 + s_2) k^0) = \delta (F (s_1 + s_2)) = \frac{\delta (s_1 + s_2 - \tilde{s})}{|F'(\tilde{s})|}$$

$$\text{where } F(s) = p^0 + s k^0 - g^0 - q^0 - \sqrt{m^2 + (\vec{p} + s \vec{k} - \vec{g} - \vec{q})^2}$$

By taking the derivative and substituting

$$\text{back } \sqrt{m^2 + (\vec{p} + \tilde{s} \vec{k} - \vec{g} - \vec{q})^2} = (p^0 + \tilde{s} k^0 - g^0 - q^0) = f^0,$$

it is straightforward to show that

$$F'(\tilde{s}) = \frac{(kf)}{f^0} = \frac{m^2 \chi_f}{\xi f^0}$$

hence

$$\delta (g^0 + f^0 + q^0 - p^0 - \tilde{s} k^0) = \frac{\xi f^0}{m^2 \chi_f} \delta (s_1 + s_2 - \tilde{s})$$

$$\int \frac{d^3 f}{(2\pi)^3 2 f^0} (2\pi)^4 \delta (g + f + q - p - (s_1 + s_2) k) = \frac{\pi \xi}{m^2 \chi_f} \delta (s_1 + s_2 - \tilde{s})$$

Step 2 : intergate over s_2, t_2

$$\int ds_1 ds_2 dt_1 dt_2 (2\pi) \delta$$

$$(s_1 + s_2 - t_1 - t_2) \frac{\pi \xi}{m^2 \chi_f} \delta (s_1 + s_2 - \tilde{s}) M(s_1, s_2, t_1, t_2)$$

$$= \frac{2\pi^2 \xi}{m^2 \chi_f} \int ds_1 dt_1 dt_2 \delta (\tilde{s} - t_1 - t_2) M(s_1, \tilde{s} - s_1, t_1, t_2)$$

$$= \frac{2\pi^2 \xi}{m^2 \chi_f} \int ds dt M(s_1, \tilde{s} - s, t, \tilde{s} - t), \text{ where } s = s_1, t = t_1$$

In the end, we are calculating :

$$dW =$$

$$\frac{1}{2} \times \frac{1}{2} \frac{1}{2 p^0 L_\phi} \frac{2\pi^2 \xi}{(2\pi)^6 2 g^0 2 q^0 m^2 \chi_f} \sum_{\lambda\sigma} d^3 g d^3 q ds dt M(s, \tilde{s} - s, t, \tilde{s} - t)$$

$$M(s, \tilde{s} - s, t, \tilde{s} - t) = m(s, \tilde{s} - s) m^+(t, \tilde{s} - t)$$

$$m(s, \tilde{s} - s) = e^2 \frac{1}{l^2 + i0} [\bar{u}_{q\lambda_2} \tilde{\Gamma}^\mu(s, q, p) u_{p\lambda_1}] [\bar{u}_{g\sigma_2} \tilde{\Gamma}_\mu(\tilde{s} - s, f, -g) v_{g\sigma_1}]$$

$$m^+(t, \tilde{s} - t) =$$

$$e^2 \frac{1}{l'^2 - i0} [\bar{u}_{p\lambda_1} (\tilde{\Gamma}^{\mu 1} (t, q, p))^+ u_{q\lambda_2}] [\bar{v}_{g\sigma_1} (\tilde{\Gamma}^{\nu 1} (\tilde{s} - t, f, -g))^+ u_{f\sigma_2}]$$

$$l = p + s - k - q$$

$$l' = p + t - k - q$$

Step 3: Spin sums

For technical convenience,
we first define the operation for a dummy matrix element mod squared
 $[\bar{u}_{q\lambda_2} \gamma^\mu u_{p\lambda_1}] [\bar{u}_{g\sigma_2} \gamma^\nu v_{g\sigma_1}] g_{\mu\nu} [\bar{u}_{p\lambda_1} \gamma^{\mu 1} u_{q\lambda_2}] [\bar{v}_{g\sigma_1} \gamma^{\nu 1} u_{f\sigma_2}] g_{\mu 1 \nu 1},$
then perform the formal spin summation, and after this we substitute
 $\gamma^\mu \rightarrow \tilde{\Gamma}^\mu (s, q, p)$
 $\gamma^\nu \rightarrow \tilde{\Gamma}^\nu (\tilde{s} - s, f, -g)$
 $\gamma^{\mu 1} \rightarrow (\tilde{\Gamma}^{\mu 1} (t, q, p))^+$
 $\gamma^{\nu 1} \rightarrow (\tilde{\Gamma}^{\nu 1} (\tilde{s} - t, f, -g))^+$

```
In[19]:= DummyExpr = (SpinorUBarD[q, m].GAD[μ].SpinorUD[p, m])
              (SpinorUBarD[f, m].GAD[ν].SpinorVD[g, m])
DummyExprC =
  ComplexConjugate[DummyExpr, FCRenameDummyIndices → False] /. {μ → μ1, ν → ν1}

Out[19]= u(q, m).γμ.u(p, m) u(f, m).γν.v(g, m)

Out[20]= (φ(-g, m)).γν1.(φ(f, m)) (φ(p, m)).γμ1.(φ(q, m))

In[21]:= DummyExpr1 = FermionSpinSum[DummyExpr DummyExprC]

Out[21]= tr((γ.f + m).γν.(γ.g - m).γν1) tr((m + γ.p).γμ1.(m + γ.q).γμ)
```

We define

$$\tilde{\Gamma}^\mu (s, q, p) = \gamma^\mu A_0 (s, q, p) - \frac{e}{2} \left(\frac{(\gamma a) (\gamma k) \gamma^\mu}{kq} + \frac{\gamma^\mu (\gamma k) (\gamma a)}{kp} \right) A_1 (s, q, p) - \frac{e^2 a^2 k^\mu (\gamma k)}{2 kq kp} A_2 (s, q, p)$$

$$(\tilde{\Gamma}^{\mu 1} (t, q, p))^+ = \gamma^\mu B_0 (t, q, p) - \frac{e}{2} \left(\frac{\gamma^\mu (\gamma k) (\gamma a)}{kq} + \frac{(\gamma a) (\gamma k) \gamma^\mu}{kp} \right) B_1 (t, q, p) - \frac{e^2 a^2 k^\mu (\gamma k)}{2 kq kp} B_2 (t, q, p)$$

$$B_n = (A_n)^*$$

It what follows,
our goal is to integrate the expression for the trident rate;

This involves some variable changes
in the integrand and algebraic simplifications;

To track the calculation, we split the expression into variables

- **Expr** - the integrand
- **Coeff** -
the overall coefficient (also dependent on the integration variables)
- **Diffs** - the integration measure to track which integrations are left,
namely

$$W = \int [\text{Diffs}] * \text{Coeff} * \text{Expr}$$

We will be modifying there three variables,
in each step adding a number to the name (DiffsN, CoeffN, ExprN)

Commands **rtilde** and **rtildeC** are introduced
in definitions.nb in accordance with formulas for

$\tilde{\Gamma}^{\mu}(s, q, p)$ and $(\tilde{\Gamma}^{\mu}(t, q, p))^+$, respectively;

chif[p] is also a handy command introduced in definitions.nb,
it returns symbol ' χ_p ' for a corresponding argument 'p'

```
In[22]:= Diffs = ds dt d3q d3g
Coeff = (1 / 2) (1 / 2) (1 / Lφ / 2 / p0) 2 π^2 ξ / ((2 π) ^6 2 g0 2 q0 m^2 chif[f])
e^4 / (l1v2 + iε) / (l2v2 - iε) /. {e^4 → (4 π α) ^2}
Expr = DummyExpr1 /.
{DiracGamma[LorentzIndex[μ, D], D] → rtilde[μ, s, q, p],
DiracGamma[LorentzIndex[ν, D], D] → rtilde[μ, stilde - s, f, -g],
DiracGamma[LorentzIndex[μ1, D], D] → rtildeC[μ1, t, q, p],
DiracGamma[LorentzIndex[ν1, D], D] → rtildeC[μ1, stilde - t, f, -g]}
```

```
Out[22]= d3g d3q ds dt
```

$$\text{Out[23]} = \frac{\alpha^2 \xi}{64 \pi^2 g_0 L \phi m^2 p_0 q_0 \chi f (i\epsilon + l1^2) (l2^2 - i\epsilon)}$$

$$\begin{aligned} \text{Out[24]} = & \text{tr} \left((m + \gamma \cdot p) \left(-\frac{a^2 e^2 k^{\mu 1} \gamma \cdot k B_2(t, q, p)}{2 (k \cdot p) (k \cdot q)} - \frac{1}{2} e B_1(t, q, p) \left(\frac{(\gamma \cdot a) \cdot (\gamma \cdot k) \cdot \gamma^{\mu 1}}{k \cdot p} + \frac{\gamma^{\mu 1} \cdot (\gamma \cdot k) \cdot (\gamma \cdot a)}{k \cdot q} \right) + \gamma^{\mu 1} B_0(t, q, p) \right) (m + \gamma \cdot q) \cdot \right. \\ & \left. \left(-\frac{a^2 e^2 k^{\mu} \gamma \cdot k A_2(s, q, p)}{2 (k \cdot p) (k \cdot q)} - \frac{1}{2} e A_1(s, q, p) \left(\frac{\gamma^{\mu} \cdot (\gamma \cdot k) \cdot (\gamma \cdot a)}{k \cdot p} + \frac{(\gamma \cdot a) \cdot (\gamma \cdot k) \cdot \gamma^{\mu}}{k \cdot q} \right) + \gamma^{\mu} A_0(s, q, p) \right) \right) \\ & \text{tr} \left((\gamma \cdot f + m) \left(\frac{a^2 e^2 k^{\mu} \gamma \cdot k A_2(\text{stilde} - s, f, -g)}{2 (f \cdot k) (g \cdot k)} - \frac{1}{2} e A_1(\text{stilde} - s, f, -g) \left(\frac{(\gamma \cdot a) \cdot (\gamma \cdot k) \cdot \gamma^{\mu}}{f \cdot k} - \frac{\gamma^{\mu} \cdot (\gamma \cdot k) \cdot (\gamma \cdot a)}{g \cdot k} \right) + \right. \right. \\ & \left. \left. \gamma^{\mu} A_0(\text{stilde} - s, f, -g) \right) (\gamma \cdot g - m) \left(\frac{a^2 e^2 k^{\mu 1} \gamma \cdot k B_2(\text{stilde} - t, f, -g)}{2 (f \cdot k) (g \cdot k)} - \right. \right. \\ & \left. \left. \frac{1}{2} e B_1(\text{stilde} - t, f, -g) \left(\frac{\gamma^{\mu 1} \cdot (\gamma \cdot k) \cdot (\gamma \cdot a)}{f \cdot k} - \frac{(\gamma \cdot a) \cdot (\gamma \cdot k) \cdot \gamma^{\mu 1}}{g \cdot k} \right) + \gamma^{\mu 1} B_0(\text{stilde} - t, f, -g) \right) \right) \end{aligned}$$

```
In[25]:= Diffs1 = Diffs;
Coeff1 = Coeff;
```

The following step is saved as a pre - evaluated expression stored in TridentDiracSimplify.nb. To perform the actual computation, uncomment commented line. The calculation takes several minutes

```
In[27]:= (*Expr1=DiracSimplify[Expr];*)
NotebookEvaluate[
NotebookOpen[NotebookDirectory[] <> "TridentDiracSimplify.nb"]];
```

Step 4: Rearranging terms in Expr and further simplifications

SortTerms and **SortTermsSimplify** are two lists, which we use as an argument in command Collect to collect terms in Expr

```
In[28]:= SortTerms[X_] := Collect[X, {A0 B0, A0 B1, A0 B2, A1 B0, A1 B1, A1 B2, A2 B0,
A2 B1, A2 B2, C0 D0, C0 D1, C0 C2, C1 D0, C1 D1, C1 C2, C2 D0, C2 D1, C2 C2}]
SortTermsSimplify[X_] :=
Collect[X, {A0 B0, A0 B1, A0 B2, A1 B0, A1 B1, A1 B2, A2 B0, A2 B1, A2 B2,
C0 D0, C0 D1, C0 C2, C1 D0, C1 D1, C1 C2, C2 D0, C2 D1, C2 C2}, Simplify]
```

For brevity, we substitute A_n and B_n functions with symbols :
 $A_n[s, q, p] \rightarrow A_n$

$$B_n[t, q, p] \rightarrow B_n$$

$$A_n[\tilde{s} - s, f, -g] \rightarrow C_n$$

$$B_n[\tilde{s} - t, f, -g] \rightarrow D_n$$

Then terms in the expressions are collected according to the rules introduced in the previous input cell

```
In[30]:= DiffS2 = DiffS1;
Coeff2 = Coeff1;
Expr1 /. {A0[s, q, p] → A0, A1[s, q, p] → A1, A2[s, q, p] → A2,
  B0[t, q, p] → B0, B1[t, q, p] → B1, B2[t, q, p] → B2,
  A0[stilde - s, f, -g] → C0,
  A1[stilde - s, f, -g] → C1, A2[stilde - s, f, -g] → C2,
  B0[stilde - t, f, -g] → D0, B1[stilde - t, f, -g] → D1, B2[stilde - t, f, -g] → D2};
Expr2 = SortTerms[%];
Expr2 /. {D → 4}
```

$$\begin{aligned} \text{Out[34]} = & \frac{16 A_2 B_2 C_0 D_0 \bar{a}^4 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^4}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} + A_1 B_2 \left(C_1 D_0 \left(\frac{8 \bar{a}^4 (\bar{f} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^4 (\bar{g} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{p}} + \frac{8 \bar{a}^4 (\bar{f} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{q}} - \frac{8 \bar{a}^4 (\bar{g} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{q}} \right) + \right. \\ & C_0 D_0 \left(-\frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{q}} - \right. \\ & \left. \left. \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{q}} + \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^3}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} + \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^3}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} \right) \right) + \\ & A_2 B_1 \left(C_0 D_1 \left(\frac{8 \bar{a}^4 (\bar{f} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^4 (\bar{g} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{p}} + \frac{8 \bar{a}^4 (\bar{f} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{q}} - \frac{8 \bar{a}^4 (\bar{g} \cdot \bar{k}) e^4}{\bar{k} \cdot \bar{q}} \right) + \right. \\ & C_0 D_0 \left(-\frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{q}} - \right. \\ & \left. \left. \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) e^3}{\bar{k} \cdot \bar{q}} + \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^3}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} + \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^3}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} \right) \right) + \\ & A_0 B_0 \left(\frac{16 C_2 D_2 \bar{a}^4 (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^4}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} + \frac{8 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^3}{\bar{f} \cdot \bar{k}} - \frac{8 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^3}{\bar{g} \cdot \bar{k}} + \right. \\ & \frac{16 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - \frac{16 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} + \frac{8 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{\bar{f} \cdot \bar{k}} - \\ & \frac{8 C_1 D_2 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{\bar{g} \cdot \bar{k}} + 16 C_0 D_2 m^2 \bar{a}^2 e^2 + \frac{8 C_0 D_2 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} + \\ & \frac{8 C_0 D_2 \bar{a}^2 (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{g} \cdot \bar{k}} + \frac{8 C_0 D_2 \bar{a}^2 (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{f} \cdot \bar{k}} + \frac{8 C_0 D_2 \bar{a}^2 (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{g} \cdot \bar{k}} - \\ & \left. \frac{16 C_0 D_2 m^2 \bar{a}^2 (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - \frac{16 C_0 D_2 \bar{a}^2 (\bar{f} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - 16 C_0 D_2 \bar{a}^2 (\bar{p} \cdot \bar{q}) e^2 + \right. \end{aligned}$$

$$\begin{aligned}
& \text{C1 D0} \left(16 e (\bar{a} \cdot \bar{f}) m^2 - 16 e (\bar{a} \cdot \bar{g}) m^2 + \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) m^2}{\bar{f} \cdot \bar{k}} - \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) m^2}{\bar{g} \cdot \bar{k}} - 16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p}) - \right. \\
& \quad 16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q}) + 16 e (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) + 16 e (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q}) - \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p})}{\bar{f} \cdot \bar{k}} + \\
& \quad \left. \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p})}{\bar{g} \cdot \bar{k}} - \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q})}{\bar{f} \cdot \bar{k}} + \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q})}{\bar{g} \cdot \bar{k}} \right) + \\
& \text{C0 D1} \left(16 e (\bar{a} \cdot \bar{f}) m^2 - 16 e (\bar{a} \cdot \bar{g}) m^2 + \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{k}) m^2}{\bar{f} \cdot \bar{k}} - \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{k}) m^2}{\bar{g} \cdot \bar{k}} - 16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p}) - \right. \\
& \quad 16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q}) + 16 e (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) + 16 e (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q}) - \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p})}{\bar{f} \cdot \bar{k}} + \\
& \quad \left. \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p})}{\bar{g} \cdot \bar{k}} - \frac{16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q})}{\bar{f} \cdot \bar{k}} + \frac{16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q})}{\bar{g} \cdot \bar{k}} \right) + \\
& \text{C2 D1} \left(\frac{8 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^3}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^3}{\bar{g} \cdot \bar{k}} + \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - \right. \\
& \quad \left. \frac{16 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^3}{\bar{g} \cdot \bar{k}} \right) + \\
& \text{C0 D0} (-64 m^4 - 32 (\bar{f} \cdot \bar{g}) m^2 + 32 (\bar{p} \cdot \bar{q}) m^2 + 32 (\bar{f} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) + 32 (\bar{f} \cdot \bar{p}) (\bar{g} \cdot \bar{q})) + \\
& \text{C2 D0} \left(16 m^2 \bar{a}^2 e^2 + \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{f} \cdot \bar{k}} + \right. \\
& \quad \left. \frac{8 \bar{a}^2 (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{g} \cdot \bar{k}} - \frac{16 m^2 \bar{a}^2 (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - \frac{16 \bar{a}^2 (\bar{f} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - 16 \bar{a}^2 (\bar{p} \cdot \bar{q}) e^2 \right) + \\
& \text{C1 D1} \left(-64 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) e^2 + \frac{16 m^2 \bar{a}^2 (\bar{g} \cdot \bar{k}) e^2}{\bar{f} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} - \right. \\
& \quad \frac{16 \bar{a}^2 (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{g} \cdot \bar{k}} + \frac{32 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} + \frac{32 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{g} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{\bar{g} \cdot \bar{k}} - \\
& \quad \frac{16 \bar{a}^2 (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{f} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{f} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{g} \cdot \bar{k}} + \frac{32 m^2 \bar{a}^2 (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} - \\
& \quad \frac{64 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} + \frac{32 \bar{a}^2 (\bar{f} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k})} + \frac{32 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{f} \cdot \bar{k}} + \\
& \quad \left. \frac{32 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{g} \cdot \bar{k}} - \frac{16 \bar{a}^2 (\bar{f} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{\bar{g} \cdot \bar{k}} + 32 \bar{a}^2 (\bar{p} \cdot \bar{q}) e^2 + \frac{16 m^2 \bar{a}^2 (\bar{f} \cdot \bar{k}) e^2}{\bar{g} \cdot \bar{k}} \right) + \\
& \text{A0 B1} \left(\text{C2 D1} \left(\frac{8 \bar{a}^4 (\bar{k} \cdot \bar{p}) e^4}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^4 (\bar{k} \cdot \bar{p}) e^4}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^4 (\bar{k} \cdot \bar{q}) e^4}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^4 (\bar{k} \cdot \bar{q}) e^4}{\bar{g} \cdot \bar{k}} \right) + \right. \\
& \quad \left. \text{C2 D0} \left(-16 \bar{a}^2 (\bar{a} \cdot \bar{p}) e^3 - 16 \bar{a}^2 (\bar{a} \cdot \bar{q}) e^3 + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) e^3}{\bar{f} \cdot \bar{k}} + \right. \right.
\end{aligned}$$

$$\begin{aligned}
& \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{p}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} + \frac{16 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} + \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} - \\
& \frac{16 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{q}) e^2}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} - \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{p} \cdot \bar{q}) e^2}{\bar{k} \cdot \bar{p}} + \frac{8 \bar{a}^2 (\bar{g} \cdot \bar{k}) (\bar{p} \cdot \bar{q}) e^2}{\bar{k} \cdot \bar{p}} - \\
& \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{p} \cdot \bar{q}) e^2}{\bar{k} \cdot \bar{q}} + \frac{8 \bar{a}^2 (\bar{g} \cdot \bar{k}) (\bar{p} \cdot \bar{q}) e^2}{\bar{k} \cdot \bar{q}} + \frac{8 m^2 \bar{a}^2 (\bar{f} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{p}} + \frac{16 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{p}} - \\
& \frac{8 m^2 \bar{a}^2 (\bar{g} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{p}} - \frac{16 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{p}} + \frac{8 m^2 \bar{a}^2 (\bar{f} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{q}} + \frac{16 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{q}} - \\
& \frac{8 m^2 \bar{a}^2 (\bar{g} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{q}} - \frac{16 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{q}} - \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} + \\
& \frac{16 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p}) e^2}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} + \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) e^2}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} - \frac{16 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{p}) e^2}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} \Bigg) + \\
& \text{A1 B0} \left(\frac{8 \text{C1 D2 } \bar{a}^4 (\bar{k} \cdot \bar{p}) e^4}{\bar{f} \cdot \bar{k}} - \frac{8 \text{C1 D2 } \bar{a}^4 (\bar{k} \cdot \bar{p}) e^4}{\bar{g} \cdot \bar{k}} + \frac{8 \text{C1 D2 } \bar{a}^4 (\bar{k} \cdot \bar{q}) e^4}{\bar{f} \cdot \bar{k}} - \frac{8 \text{C1 D2 } \bar{a}^4 (\bar{k} \cdot \bar{q}) e^4}{\bar{g} \cdot \bar{k}} - \right. \\
& \frac{16 \text{C0 D2 } \bar{a}^2 (\bar{a} \cdot \bar{p}) e^3 - 16 \text{C0 D2 } \bar{a}^2 (\bar{a} \cdot \bar{q}) e^3 +}{\bar{f} \cdot \bar{k}} + \frac{8 \text{C0 D2 } \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) e^3}{\bar{g} \cdot \bar{k}} + \\
& \frac{8 \text{C0 D2 } \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{q}) e^3}{\bar{f} \cdot \bar{k}} + \frac{8 \text{C0 D2 } \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{q}) e^3}{\bar{g} \cdot \bar{k}} + \\
& \text{C0 D0} \left(16 e (\bar{a} \cdot \bar{p}) m^2 + 16 e (\bar{a} \cdot \bar{q}) m^2 - \frac{16 e (\bar{a} \cdot \bar{p}) (\bar{k} \cdot \bar{q}) m^2}{\bar{k} \cdot \bar{p}} - \frac{16 e (\bar{a} \cdot \bar{q}) (\bar{k} \cdot \bar{p}) m^2}{\bar{k} \cdot \bar{q}} + 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p}) + \right. \\
& 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q}) + 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p}) + 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q}) - \frac{16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q}) (\bar{g} \cdot \bar{k})}{\bar{k} \cdot \bar{p}} - \\
& \left. \frac{16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{q})}{\bar{k} \cdot \bar{p}} - \frac{16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p}) (\bar{g} \cdot \bar{k})}{\bar{k} \cdot \bar{q}} - \frac{16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{p})}{\bar{k} \cdot \bar{q}} \right) + \\
& \text{C1 D1} \left(-16 \bar{a}^2 (\bar{a} \cdot \bar{p}) e^3 - 16 \bar{a}^2 (\bar{a} \cdot \bar{q}) e^3 + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) e^3}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) e^3}{\bar{f} \cdot \bar{k}} - \right. \\
& \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{p}) e^3}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) e^3}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{q}) e^3}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{q}) e^3}{\bar{f} \cdot \bar{k}} - \\
& \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{f}) (\bar{k} \cdot \bar{q}) e^3}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{g}) (\bar{k} \cdot \bar{q}) e^3}{\bar{g} \cdot \bar{k}} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} + \\
& \left. \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{q}) e^3}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p})} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{p}) e^3}{(\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} + \frac{8 \bar{a}^2 (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{k}) (\bar{k} \cdot \bar{p}) e^3}{(\bar{g} \cdot \bar{k}) (\bar{k} \cdot \bar{q})} \right) + \text{C1 D0} \\
& \left(-8 \bar{a}^2 (\bar{f} \cdot \bar{p}) e^2 - 8 \bar{a}^2 (\bar{f} \cdot \bar{q}) e^2 + 8 \bar{a}^2 (\bar{g} \cdot \bar{p}) e^2 + 8 \bar{a}^2 (\bar{g} \cdot \bar{q}) e^2 - \frac{8 m^2 \bar{a}^2 (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} - \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{g}) (\bar{k} \cdot \bar{p}) e^2}{\bar{f} \cdot \bar{k}} + \right.
\end{aligned}$$

$$\left. \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{p}) e^2}{\bar{k} \cdot \bar{p}} - \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{q}) (\bar{g} \cdot \bar{k}) e^2}{\bar{k} \cdot \bar{q}} - \frac{8 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{q}) e^2}{\bar{k} \cdot \bar{q}} - \frac{16 m^2 \bar{a}^2 (\bar{f} \cdot \bar{k}) (\bar{g} \cdot \bar{k}) e^2}{(\bar{k} \cdot \bar{p}) (\bar{k} \cdot \bar{q})} \right)$$

To simplify further, contractions of momenta with k , such as (kp) , are replaced by χ – parameters

$$(kp) = \frac{m^2 \chi_p}{\xi}$$

ξ is defined from

$$\xi^2 = -\frac{e^2 a^2}{m^2}$$

Function **ktochiSubs** is defined in `definitions.nb`, it generates substitutions of scalar products (kp) , (kq) , etc, to pass to the corresponding χ parameters as defined above

In[35]:= **(*example*)**

kp /. ktochiSubs[p, q, f, g]

Out[35]=
$$\frac{m^2 \chi_p}{\xi}$$

In[36]:= **Diffs3 = Diffs2;**

Coeff3 = Coeff2;

Expr3 = Expr2 /. ktochiSubs[p, q, f, g] /. {av2 → -m^2 ξ^2 / e^2};

% /. {D → 4}

Out[39]=
$$\frac{16 A2 B2 C0 D0 m^4 \chi f \chi g \xi^4}{\chi p \chi q} + A1 B2$$

$$\left(C1 D0 \left(\frac{8 m^4 \chi f \xi^4}{\chi p} - \frac{8 m^4 \chi g \xi^4}{\chi p} + \frac{8 m^4 \chi f \xi^4}{\chi q} - \frac{8 m^4 \chi g \xi^4}{\chi q} \right) + C0 D0 \left(\frac{8 e m^2 \chi g (\bar{a} \cdot \bar{f}) \xi^2}{\chi p} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{f}) \xi^2}{\chi q} + \right. \right.$$

$$\left. \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{g}) \xi^2}{\chi p} + \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{g}) \xi^2}{\chi q} - \frac{16 e m^2 \chi f \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p \chi q} - \frac{16 e m^2 \chi f \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi p \chi q} \right) + A2 B1$$

$$\left(C0 D1 \left(\frac{8 m^4 \chi f \xi^4}{\chi p} - \frac{8 m^4 \chi g \xi^4}{\chi p} + \frac{8 m^4 \chi f \xi^4}{\chi q} - \frac{8 m^4 \chi g \xi^4}{\chi q} \right) + C0 D0 \left(\frac{8 e m^2 \chi g (\bar{a} \cdot \bar{f}) \xi^2}{\chi p} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{f}) \xi^2}{\chi q} + \right. \right.$$

$$\left. \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{g}) \xi^2}{\chi p} + \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{g}) \xi^2}{\chi q} - \frac{16 e m^2 \chi f \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p \chi q} - \frac{16 e m^2 \chi f \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi p \chi q} \right) +$$

$$A0 B0 \left(-16 C0 D2 \xi^2 m^4 + \frac{16 C2 D2 \xi^4 \chi p \chi q m^4}{\chi f \chi g} + \frac{16 C0 D2 \xi^2 \chi p \chi q m^4}{\chi f \chi g} - \frac{16 C1 D2 e \xi^2 \chi p \chi q (\bar{a} \cdot \bar{f}) m^2}{\chi f \chi g} + \right.$$

$$\frac{16 C1 D2 e \xi^2 \chi p \chi q (\bar{a} \cdot \bar{g}) m^2}{\chi f \chi g} - \frac{8 C1 D2 e \xi^2 \chi q (\bar{a} \cdot \bar{p}) m^2}{\chi f} + \frac{8 C1 D2 e \xi^2 \chi q (\bar{a} \cdot \bar{p}) m^2}{\chi g} -$$

$$\left. \frac{8 C1 D2 e \xi^2 \chi p (\bar{a} \cdot \bar{q}) m^2}{\chi f} + \frac{8 C1 D2 e \xi^2 \chi p (\bar{a} \cdot \bar{q}) m^2}{\chi g} + \frac{16 C0 D2 \xi^2 \chi p \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi f \chi g} - \right.$$

$$\begin{aligned}
& \frac{8 \text{ C0 D2 } \xi^2 \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi^f} - \frac{8 \text{ C0 D2 } \xi^2 \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi^f} - \frac{8 \text{ C0 D2 } \xi^2 \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi^g} - \frac{8 \text{ C0 D2 } \xi^2 \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi^g} + \\
& 16 \text{ C0 D2 } \xi^2 (\bar{p} \cdot \bar{q}) m^2 + \text{C2 D1} \left(-\frac{16 e m^2 \chi p \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi^f \chi^g} + \frac{16 e m^2 \chi p \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi^f \chi^g} - \right. \\
& \quad \left. \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi^f} + \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi^g} - \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi^f} + \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi^g} \right) + \\
& \text{C1 D0} \left(16 e (\bar{a} \cdot \bar{f}) m^2 + \frac{16 e \chi^g (\bar{a} \cdot \bar{f}) m^2}{\chi^f} - 16 e (\bar{a} \cdot \bar{g}) m^2 - \frac{16 e \chi^f (\bar{a} \cdot \bar{g}) m^2}{\chi^g} + \right. \\
& \quad \frac{16 e \chi q (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p})}{\chi^g} - 16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p}) + \frac{16 e \chi p (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q})}{\chi^g} - 16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q}) - \\
& \quad \left. \frac{16 e \chi q (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p})}{\chi^f} + 16 e (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) - \frac{16 e \chi p (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q})}{\chi^f} + 16 e (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q}) \right) + \\
& \text{C0 D1} \left(16 e (\bar{a} \cdot \bar{f}) m^2 + \frac{16 e \chi^g (\bar{a} \cdot \bar{f}) m^2}{\chi^f} - 16 e (\bar{a} \cdot \bar{g}) m^2 - \frac{16 e \chi^f (\bar{a} \cdot \bar{g}) m^2}{\chi^g} + \right. \\
& \quad \frac{16 e \chi q (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p})}{\chi^g} - 16 e (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p}) + \frac{16 e \chi p (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q})}{\chi^g} - 16 e (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q}) - \\
& \quad \left. \frac{16 e \chi q (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p})}{\chi^f} + 16 e (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) - \frac{16 e \chi p (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q})}{\chi^f} + 16 e (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q}) \right) + \\
& \text{C0 D0} (-64 m^4 - 32 (\bar{f} \cdot \bar{g}) m^2 + 32 (\bar{p} \cdot \bar{q}) m^2 + 32 (\bar{f} \cdot \bar{q}) (\bar{g} \cdot \bar{p}) + 32 (\bar{f} \cdot \bar{p}) (\bar{g} \cdot \bar{q})) + \\
& \text{C1 D1} \left(-\frac{16 \xi^2 \chi^g m^4}{\chi^f} - \frac{32 \xi^2 \chi p \chi q m^4}{\chi^f \chi^g} - \frac{16 \xi^2 \chi^f m^4}{\chi^g} - \frac{32 \xi^2 \chi p \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi^f \chi^g} + \right. \\
& \quad \frac{16 \xi^2 \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi^f} + \frac{16 \xi^2 \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi^g} + \frac{16 \xi^2 \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi^f} + \frac{16 \xi^2 \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi^g} + \\
& \quad \frac{16 \xi^2 \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi^f} + \frac{16 \xi^2 \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi^g} + \frac{16 \xi^2 \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi^f} + \frac{16 \xi^2 \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi^g} - \\
& \quad 32 \xi^2 (\bar{p} \cdot \bar{q}) m^2 - \frac{64 e^2 \chi p \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi^f \chi^g} + \frac{32 e^2 \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p})}{\chi^f} + \frac{32 e^2 \chi q (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p})}{\chi^g} + \\
& \quad \left. \frac{32 e^2 \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q})}{\chi^f} + \frac{32 e^2 \chi p (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q})}{\chi^g} - 64 e^2 (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q}) \right) + \\
& \text{C2 D0} \left(-16 \xi^2 m^4 + \frac{16 \xi^2 \chi p \chi q m^4}{\chi^f \chi^g} + \frac{16 \xi^2 \chi p \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi^f \chi^g} - \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi^f} - \right. \\
& \quad \left. \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi^f} - \frac{8 \xi^2 \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi^g} - \frac{8 \xi^2 \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi^g} + 16 \xi^2 (\bar{p} \cdot \bar{q}) m^2 \right) + \\
& \text{A1 B0} \left(\frac{8 \text{ C1 D2 } m^4 \chi p \xi^4}{\chi^f} - \frac{8 \text{ C1 D2 } m^4 \chi p \xi^4}{\chi^g} + \frac{8 \text{ C1 D2 } m^4 \chi q \xi^4}{\chi^f} - \frac{8 \text{ C1 D2 } m^4 \chi q \xi^4}{\chi^g} - \right. \\
& \quad \left. \frac{8 \text{ C0 D2 } e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi^f} - \frac{8 \text{ C0 D2 } e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi^f} - \frac{8 \text{ C0 D2 } e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi^g} - \right.
\end{aligned}$$

$$\begin{aligned}
& \frac{8 \text{ C0 D2 } e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + 16 \text{ C0 D2 } e m^2 (\bar{a} \cdot \bar{p}) \xi^2 + 16 \text{ C0 D2 } e m^2 (\bar{a} \cdot \bar{q}) \xi^2 + \\
& \text{C1 D1} \left(-\frac{8 e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} + \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi g} - \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} + \right. \\
& \quad \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi g} + \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi f} - \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi f} - \\
& \quad \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + 16 e m^2 (\bar{a} \cdot \bar{p}) \xi^2 - \frac{8 e m^2 \chi g \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi f \chi p} - \frac{8 e m^2 \chi f \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi g \chi p} + \\
& \quad \left. 16 e m^2 (\bar{a} \cdot \bar{q}) \xi^2 - \frac{8 e m^2 \chi g \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi f \chi q} - \frac{8 e m^2 \chi f \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi g \chi q} \right) + \\
& \text{C0 D0} \left(16 e (\bar{a} \cdot \bar{p}) m^2 - \frac{16 e \chi q (\bar{a} \cdot \bar{p}) m^2}{\chi p} + 16 e (\bar{a} \cdot \bar{q}) m^2 - \frac{16 e \chi p (\bar{a} \cdot \bar{q}) m^2}{\chi q} + \right. \\
& \quad 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p}) - \frac{16 e \chi g (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p})}{\chi q} + 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q}) - \frac{16 e \chi g (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q})}{\chi p} + \\
& \quad \left. 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p}) - \frac{16 e \chi f (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p})}{\chi q} + 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q}) - \frac{16 e \chi f (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q})}{\chi p} \right) + \\
& \text{C0 D1} \left(-\frac{8 \xi^2 \chi p m^4}{\chi f} + \frac{8 \xi^2 \chi p m^4}{\chi g} - \frac{8 \xi^2 \chi q m^4}{\chi f} + \frac{8 \xi^2 \chi q m^4}{\chi g} - \frac{8 \xi^2 \chi f m^4}{\chi p} + \frac{8 \xi^2 \chi g m^4}{\chi p} - \right. \\
& \quad \frac{8 \xi^2 \chi f m^4}{\chi q} + \frac{8 \xi^2 \chi g m^4}{\chi q} - \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi f} + \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi g} - \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi f} + \\
& \quad \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi g} + 8 \xi^2 (\bar{f} \cdot \bar{p}) m^2 + \frac{8 \xi^2 \chi g \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi f \chi p} + 8 \xi^2 (\bar{f} \cdot \bar{q}) m^2 + \frac{8 \xi^2 \chi g \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi f \chi q} - \\
& \quad 8 \xi^2 (\bar{g} \cdot \bar{p}) m^2 - \frac{8 \xi^2 \chi f \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi g \chi p} - 8 \xi^2 (\bar{g} \cdot \bar{q}) m^2 - \frac{8 \xi^2 \chi f \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi g \chi q} + \\
& \quad \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi p} - \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p} + \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi q} - \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi q} - \\
& \quad \frac{16 e^2 \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi f} + \frac{16 e^2 \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi g} - \frac{16 e^2 \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi f} + \frac{16 e^2 \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi g} - \\
& \quad 16 e^2 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p}) + \frac{16 e^2 \chi g \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p})}{\chi f \chi p} + 16 e^2 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p}) - \frac{16 e^2 \chi f \chi q (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p})}{\chi g \chi p} - \\
& \quad 16 e^2 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q}) + \frac{16 e^2 \chi g \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q})}{\chi f \chi q} + 16 e^2 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q}) - \frac{16 e^2 \chi f \chi p (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q})}{\chi g \chi q} + \\
& \quad \left. \frac{16 e^2 \chi f (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi p} - \frac{16 e^2 \chi g (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi p} + \frac{16 e^2 \chi f (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi q} - \frac{16 e^2 \chi g (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi q} \right) + \\
& \text{C1 D0} \left(\frac{8 \xi^2 \chi p m^4}{\chi f} - \frac{8 \xi^2 \chi p m^4}{\chi g} + \frac{8 \xi^2 \chi q m^4}{\chi f} - \frac{8 \xi^2 \chi q m^4}{\chi g} + \frac{8 \xi^2 \chi f m^4}{\chi p} - \frac{8 \xi^2 \chi g m^4}{\chi p} + \right. \\
& \quad \frac{8 \xi^2 \chi f m^4}{\chi q} - \frac{8 \xi^2 \chi g m^4}{\chi q} + \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi f} - \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi g} + \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi f} - \\
& \quad \left. \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi g} \right)
\end{aligned}$$

$$\begin{aligned}
& \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi g} + 8 \xi^2 (\bar{f} \cdot \bar{p}) m^2 - \frac{8 \xi^2 \chi g \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi f \chi p} + 8 \xi^2 (\bar{f} \cdot \bar{q}) m^2 - \frac{8 \xi^2 \chi g \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi f \chi q} - \\
& 8 \xi^2 (\bar{g} \cdot \bar{p}) m^2 + \frac{8 \xi^2 \chi f \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi g \chi p} - 8 \xi^2 (\bar{g} \cdot \bar{q}) m^2 + \frac{8 \xi^2 \chi f \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi g \chi q} - \\
& \left. \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi p} + \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p} - \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi q} + \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi q} \right) \Bigg) + \\
A0 B1 & \left(C2 D1 \left(\frac{8 m^4 \chi p \xi^4}{\chi f} - \frac{8 m^4 \chi p \xi^4}{\chi g} + \frac{8 m^4 \chi q \xi^4}{\chi f} - \frac{8 m^4 \chi q \xi^4}{\chi g} \right) + C2 D0 \left(-\frac{8 e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} - \right. \right. \\
& \left. \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} - \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} - \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + 16 e m^2 (\bar{a} \cdot \bar{p}) \xi^2 + 16 e m^2 (\bar{a} \cdot \bar{q}) \xi^2 \right) + \\
C1 D1 & \left(-\frac{8 e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} + \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi g} - \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi f} + \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi g} + \right. \\
& \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi f} - \frac{8 e m^2 \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi f} - \frac{8 e m^2 \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g} + \\
& 16 e m^2 (\bar{a} \cdot \bar{p}) \xi^2 - \frac{8 e m^2 \chi g \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi f \chi p} - \frac{8 e m^2 \chi f \chi q (\bar{a} \cdot \bar{p}) \xi^2}{\chi g \chi p} + \\
& \left. 16 e m^2 (\bar{a} \cdot \bar{q}) \xi^2 - \frac{8 e m^2 \chi g \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi f \chi q} - \frac{8 e m^2 \chi f \chi p (\bar{a} \cdot \bar{q}) \xi^2}{\chi g \chi q} \right) + \\
C0 D0 & \left(16 e (\bar{a} \cdot \bar{p}) m^2 - \frac{16 e \chi q (\bar{a} \cdot \bar{p}) m^2}{\chi p} + 16 e (\bar{a} \cdot \bar{q}) m^2 - \frac{16 e \chi p (\bar{a} \cdot \bar{q}) m^2}{\chi q} + \right. \\
& 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{p}) - \frac{16 e \chi g (\bar{a} \cdot \bar{q}) (\bar{f} \cdot \bar{p})}{\chi q} + 16 e (\bar{a} \cdot \bar{g}) (\bar{f} \cdot \bar{q}) - \frac{16 e \chi g (\bar{a} \cdot \bar{p}) (\bar{f} \cdot \bar{q})}{\chi p} + \\
& \left. 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{p}) - \frac{16 e \chi f (\bar{a} \cdot \bar{q}) (\bar{g} \cdot \bar{p})}{\chi q} + 16 e (\bar{a} \cdot \bar{f}) (\bar{g} \cdot \bar{q}) - \frac{16 e \chi f (\bar{a} \cdot \bar{p}) (\bar{g} \cdot \bar{q})}{\chi p} \right) + \\
C1 D0 & \left(-\frac{8 \xi^2 \chi p m^4}{\chi f} + \frac{8 \xi^2 \chi p m^4}{\chi g} - \frac{8 \xi^2 \chi q m^4}{\chi f} + \frac{8 \xi^2 \chi q m^4}{\chi g} - \frac{8 \xi^2 \chi f m^4}{\chi p} + \frac{8 \xi^2 \chi g m^4}{\chi p} - \right. \\
& \frac{8 \xi^2 \chi f m^4}{\chi q} + \frac{8 \xi^2 \chi g m^4}{\chi q} - \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi f} + \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi g} - \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi f} + \\
& \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi g} + 8 \xi^2 (\bar{f} \cdot \bar{p}) m^2 + \frac{8 \xi^2 \chi g \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi f \chi p} + 8 \xi^2 (\bar{f} \cdot \bar{q}) m^2 + \frac{8 \xi^2 \chi g \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi f \chi q} - \\
& 8 \xi^2 (\bar{g} \cdot \bar{p}) m^2 - \frac{8 \xi^2 \chi f \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi g \chi p} - 8 \xi^2 (\bar{g} \cdot \bar{q}) m^2 - \frac{8 \xi^2 \chi f \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi g \chi q} + \\
& \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi p} - \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p} + \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi q} - \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi q} - \\
& \frac{16 e^2 \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi f} + \frac{16 e^2 \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi g} - \frac{16 e^2 \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi f} + \frac{16 e^2 \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g})}{\chi g} - \\
& 16 e^2 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p}) + \frac{16 e^2 \chi g \chi q (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p})}{\chi f \chi p} + 16 e^2 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p}) - \frac{16 e^2 \chi f \chi q (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p})}{\chi g \chi p} -
\end{aligned}$$

$$\begin{aligned}
& 16 e^2 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q}) + \frac{16 e^2 \chi g \chi p (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q})}{\chi f \chi q} + 16 e^2 (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q}) - \frac{16 e^2 \chi f \chi p (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q})}{\chi g \chi q} + \\
& \left. \frac{16 e^2 \chi f (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi p} - \frac{16 e^2 \chi g (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi p} + \frac{16 e^2 \chi f (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi q} - \frac{16 e^2 \chi g (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi q} \right) + \\
& \text{C0 D1} \left(\frac{8 \xi^2 \chi p m^4}{\chi f} - \frac{8 \xi^2 \chi p m^4}{\chi g} + \frac{8 \xi^2 \chi q m^4}{\chi f} - \frac{8 \xi^2 \chi q m^4}{\chi g} + \frac{8 \xi^2 \chi f m^4}{\chi p} - \frac{8 \xi^2 \chi g m^4}{\chi p} + \right. \\
& \frac{8 \xi^2 \chi f m^4}{\chi q} - \frac{8 \xi^2 \chi g m^4}{\chi q} + \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi f} - \frac{8 \xi^2 \chi p (\bar{f} \cdot \bar{g}) m^2}{\chi g} + \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi f} - \\
& \frac{8 \xi^2 \chi q (\bar{f} \cdot \bar{g}) m^2}{\chi g} + 8 \xi^2 (\bar{f} \cdot \bar{p}) m^2 - \frac{8 \xi^2 \chi g \chi q (\bar{f} \cdot \bar{p}) m^2}{\chi f \chi p} + 8 \xi^2 (\bar{f} \cdot \bar{q}) m^2 - \frac{8 \xi^2 \chi g \chi p (\bar{f} \cdot \bar{q}) m^2}{\chi f \chi q} - \\
& 8 \xi^2 (\bar{g} \cdot \bar{p}) m^2 + \frac{8 \xi^2 \chi f \chi q (\bar{g} \cdot \bar{p}) m^2}{\chi g \chi p} - 8 \xi^2 (\bar{g} \cdot \bar{q}) m^2 + \frac{8 \xi^2 \chi f \chi p (\bar{g} \cdot \bar{q}) m^2}{\chi g \chi q} - \\
& \left. \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi p} + \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p} - \frac{8 \xi^2 \chi f (\bar{p} \cdot \bar{q}) m^2}{\chi q} + \frac{8 \xi^2 \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi q} \right) + \\
& \text{A0 B2} \left(-16 \text{C2 D0} m^4 \xi^4 + \text{C1 D0} \left(16 e m^2 (\bar{a} \cdot \bar{f}) \xi^2 - 16 e m^2 (\bar{a} \cdot \bar{g}) \xi^2 - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} + \right. \right. \\
& \left. \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} \right) + \\
& \text{C0 D0} \left(-16 \xi^2 m^4 + \frac{16 \xi^2 \chi f \chi g m^4}{\chi p \chi q} - 16 \xi^2 (\bar{f} \cdot \bar{g}) m^2 + \frac{8 \xi^2 \chi g (\bar{f} \cdot \bar{p}) m^2}{\chi p} + \frac{8 \xi^2 \chi g (\bar{f} \cdot \bar{q}) m^2}{\chi q} + \right. \\
& \left. \frac{8 \xi^2 \chi f (\bar{g} \cdot \bar{p}) m^2}{\chi p} + \frac{8 \xi^2 \chi f (\bar{g} \cdot \bar{q}) m^2}{\chi q} - \frac{16 \xi^2 \chi f \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p \chi q} \right) + \\
& \text{A2 B0} \left(-16 \text{C0 D2} m^4 \xi^4 + \text{C0 D1} \left(16 e m^2 (\bar{a} \cdot \bar{f}) \xi^2 - 16 e m^2 (\bar{a} \cdot \bar{g}) \xi^2 - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} + \right. \right. \\
& \left. \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} \right) + \\
& \text{C0 D0} \left(-16 \xi^2 m^4 + \frac{16 \xi^2 \chi f \chi g m^4}{\chi p \chi q} - 16 \xi^2 (\bar{f} \cdot \bar{g}) m^2 + \frac{8 \xi^2 \chi g (\bar{f} \cdot \bar{p}) m^2}{\chi p} + \frac{8 \xi^2 \chi g (\bar{f} \cdot \bar{q}) m^2}{\chi q} + \right. \\
& \left. \frac{8 \xi^2 \chi f (\bar{g} \cdot \bar{p}) m^2}{\chi p} + \frac{8 \xi^2 \chi f (\bar{g} \cdot \bar{q}) m^2}{\chi q} - \frac{16 \xi^2 \chi f \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p \chi q} \right) + \\
& \text{A1 B1} \left(\text{C1 D1} \left(-32 m^4 \xi^4 + \frac{8 m^4 \chi g \chi q \xi^4}{\chi f \chi p} + \frac{8 m^4 \chi f \chi q \xi^4}{\chi g \chi p} + \frac{8 m^4 \chi g \chi p \xi^4}{\chi f \chi q} + \frac{8 m^4 \chi f \chi p \xi^4}{\chi g \chi q} \right) + \right. \\
& \text{C1 D0} \left(16 e m^2 (\bar{a} \cdot \bar{f}) \xi^2 - \frac{8 e m^2 \chi g \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi f \chi p} - \frac{8 e m^2 \chi g \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi f \chi q} - \right. \\
& 16 e m^2 (\bar{a} \cdot \bar{g}) \xi^2 + \frac{8 e m^2 \chi f \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g \chi p} + \frac{8 e m^2 \chi f \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi g \chi q} - \\
& \left. \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi q} \right) -
\end{aligned}$$

$$\begin{aligned}
& \left. \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi p} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} \right) + \\
\text{C0 D1} & \left(16 e m^2 (\bar{a} \cdot \bar{f}) \xi^2 - \frac{8 e m^2 \chi g \chi q (\bar{a} \cdot \bar{f}) \xi^2}{\chi f \chi p} - \frac{8 e m^2 \chi g \chi p (\bar{a} \cdot \bar{f}) \xi^2}{\chi f \chi q} - 16 e m^2 (\bar{a} \cdot \bar{g}) \xi^2 + \right. \\
& \frac{8 e m^2 \chi f \chi q (\bar{a} \cdot \bar{g}) \xi^2}{\chi g \chi p} + \frac{8 e m^2 \chi f \chi p (\bar{a} \cdot \bar{g}) \xi^2}{\chi g \chi q} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} + \\
& \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{p}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{p}) \xi^2}{\chi q} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi p} + \\
& \left. \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi p} - \frac{8 e m^2 \chi f (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} + \frac{8 e m^2 \chi g (\bar{a} \cdot \bar{q}) \xi^2}{\chi q} \right) + \text{C0 D0} \\
& \left(\frac{16 \xi^2 \chi q m^4}{\chi p} + \frac{16 \xi^2 \chi p m^4}{\chi q} - \frac{32 \xi^2 \chi f \chi g m^4}{\chi p \chi q} + 32 \xi^2 (\bar{f} \cdot \bar{g}) m^2 - \frac{16 \xi^2 \chi g (\bar{f} \cdot \bar{p}) m^2}{\chi p} + \frac{16 \xi^2 \chi g (\bar{f} \cdot \bar{p}) m^2}{\chi q} + \right. \\
& \frac{16 \xi^2 \chi g (\bar{f} \cdot \bar{q}) m^2}{\chi p} - \frac{16 \xi^2 \chi g (\bar{f} \cdot \bar{q}) m^2}{\chi q} - \frac{16 \xi^2 \chi f (\bar{g} \cdot \bar{p}) m^2}{\chi p} + \frac{16 \xi^2 \chi f (\bar{g} \cdot \bar{p}) m^2}{\chi q} + \frac{16 \xi^2 \chi f (\bar{g} \cdot \bar{q}) m^2}{\chi p} - \\
& \frac{16 \xi^2 \chi f (\bar{g} \cdot \bar{q}) m^2}{\chi q} + \frac{32 \xi^2 \chi f \chi g (\bar{p} \cdot \bar{q}) m^2}{\chi p \chi q} + 64 e^2 (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{g}) - \frac{32 e^2 \chi g (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{p})}{\chi p} - \\
& \left. \frac{32 e^2 \chi f (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{p})}{\chi p} - \frac{32 e^2 \chi g (\bar{a} \cdot \bar{f}) (\bar{a} \cdot \bar{q})}{\chi q} - \frac{32 e^2 \chi f (\bar{a} \cdot \bar{g}) (\bar{a} \cdot \bar{q})}{\chi q} + \frac{64 e^2 \chi f \chi g (\bar{a} \cdot \bar{p}) (\bar{a} \cdot \bar{q})}{\chi p \chi q} \right)
\end{aligned}$$

Step 5: Passing to invariant variables ρ_1 , τ_1 , ρ_2 , τ_2 in the integral

In the following step, we consider only the integrand, namely, $\text{Coeff} * \text{Expr}$. We will introduce the Jacobian of the tranformation later

Recall the conservation laws

$$p + sk = q + l$$

$$\chi p = \chi q + \chi l$$

Variables ρ_1 , τ_1 read

$$\begin{aligned}
\rho_1 &= \frac{e(qFp)}{\xi m^4 \chi l} = - \frac{e(lFp)}{\xi m^4 \chi l} = \frac{e(qFl)}{\xi m^4 \chi l} \\
\tau_1 &= - \frac{e(qF^* p)}{m^4 \chi l} = \frac{e(lF^* p)}{m^4 \chi l} = \frac{e(qF^* l)}{m^4 \chi l}
\end{aligned}$$

Similarly, we have ρ_2 and τ_2

$$t k + l = f + g$$

$$\chi l = \chi f + \chi g$$

$$\rho_2 = -\frac{e (gFf)}{\xi m^4 \chi l} = -\frac{e (gFl)}{\xi m^4 \chi l} = \frac{e (fFl)}{\xi m^4 \chi l}$$

$$\tau_2 = -\frac{e (gF^* f)}{m^4 \chi l} = -\frac{e (gF^* l)}{m^4 \chi l} = \frac{e (fF^* l)}{m^4 \chi l}$$

$$-\infty < \rho_{1,2}, \tau_{1,2} < +\infty$$

Also, notice that

$$(q F^2 p) = -a^2 k p k q$$

Expr contains contractions of momenta in pairs

$(pq), (pf), (pg), (qf), (qg), (fg)$

Let us express all such contractions through ρ_1 ,

τ_1, ρ_2, τ_2 and the χ parameters

The key idea is projecting a given 4 – momentum on a basis spanned by the field tensor and another 4 – momentum :

$$q^\mu = C_1 p^\mu + C_2 (Fp)^\mu + C_3 (F^* p)^\mu + C_4 (FFp)^\mu$$

$$C_1 = \frac{\chi_q}{\chi_p}$$

$$C_2 = -\frac{e^2 (qFp)}{m^6 \chi_p^2}$$

$$C_3 = -\frac{e^2 (qF^* p)}{m^6 \chi_p^2}$$

$$C_4 = \frac{e^2}{2 m^4 \chi_p \chi_q} \left(1 - \frac{\chi_q^2}{\chi_p^2} + \frac{e^2 (qFp)^2}{m^8 \chi_p^2} + \frac{e^2 (qF^* p)^2}{m^8 \chi_p^2} \right)$$

This projection is specific to a crossed field,

as in such field $[F^{n>2}]^{\mu\nu} = [F^* F]^{\mu\nu} = 0$

(a non – crossed field may have higher – order terms in powers of F)

Then we substitute the projected vector to

the initial scalar product. Terms such as (qFp) are re – expressed via invariant parameters

Note that expressing combinations such as (qg) via

the invariant parameters ρ_n, τ_n requires an intermediate step. The issue is that q and g enter different vertices,

and their projection results in appearance of combinations such as (pFg) , which do not enter the definitions of ρ_n , τ_n . We use that the vertices are connected by l .

Command **ExpandVector**[**p, q, μ**] is defined in definitions.nb. It allows projecting 4 – vector p on the basis spanned by q and F , and the resulting vector carries index μ ;
See the illustrating example below

Each projection is stored in a function of the Lorentz index μ , given by command

qpexpansion[μ]

where q is the vector being projected and p – the momentum spanning the basis

When defining a projection, we also introduce related substitutions of scalar combinations such as (qFp) , that reexpress them via invariant ρ_n , τ_n ;

These substitutions are stored in arrays, named, for example, as $\text{subs}_{\rho_1 \tau_1 qp}$, so that the last two letters are attributed to the momenta in the same order as in 'qpexpansion'

In[40]:= (**example**)

ExpandVector[**p, q, μ**]

$$\text{Out[40]} = \frac{e^2 FFp^\mu (e^2 (FDp \cdot q)^2 + e^2 (Fp \cdot q)^2 + m^8 \chi p^2 - m^8 \chi q^2)}{2 m^{12} \chi p^3 \chi q} - \frac{e^2 FDp^\mu (FDp \cdot q)}{m^6 \chi p^2} - \frac{e^2 Fp^\mu (Fp \cdot q)}{m^6 \chi p^2} + \frac{\chi q p^\mu}{\chi p}$$

(pq)

- project q on p
- contract with p , reexpress via ρ_n , τ_n

```
In[41]:= subs $\rho$ 1 $\tau$ 1qp = {Pair[Momentum[Fp, D], Momentum[q, D]]  $\rightarrow$   $\rho$ 1  $\xi$  m^4  $\chi$ l / e,
  Pair[Momentum[FDp, D], Momentum[q, D]]  $\rightarrow$  - $\tau$ 1 m^4  $\chi$ l / e}
qpexpansion[ $\mu$ _] = Collect[ExpandVector[p, q,  $\mu$ ] /. subs $\rho$ 1 $\tau$ 1qp,
  {pv[ $\mu$ ], Fpv[ $\mu$ ], FDPv[ $\mu$ ], FFpv[ $\mu$ ]}, Simplify]
pq = Contract[pv[ $\mu$ ]  $\times$  qv[ $\mu$ ] /. {qv[ $\mu$ ]  $\rightarrow$  qpexpansion[ $\mu$ ]}
```

$$\text{Out[41]} = \left\{ \text{Fp} \cdot q \rightarrow \frac{m^4 \xi \rho^1 \chi^1}{e}, \text{FDp} \cdot q \rightarrow -\frac{m^4 \tau^1 \chi^1}{e} \right\}$$

$$\text{Out[42]} = \frac{e^2 \text{FFp}^\mu (\xi^2 \rho^1 \chi^1 + \tau^1 \chi^1 + \chi p^2 - \chi q^2)}{2 m^4 \chi p^3 \chi q} + \frac{e \tau^1 \chi^1 \text{FDp}^\mu}{m^2 \chi p^2} - \frac{e \xi \rho^1 \chi^1 \text{Fp}^\mu}{m^2 \chi p^2} + \frac{\chi q p^\mu}{\chi p}$$

$$\text{Out[43]} = \frac{m^2 (\xi^2 \rho^1 \chi^1 + \tau^1 \chi^1 + \chi p^2 - \chi q^2)}{2 \chi p \chi q} + \frac{m^2 \chi q}{\chi p}$$

(lp), required later as an intermediate step
 – project l on p
 – contract with p, reexpress via ρ_n, τ_n

```
In[44]:= subs $\rho$ 1 $\tau$ 1lp = {Pair[Momentum[Fp, D], Momentum[l, D]]  $\rightarrow$  - $\rho$ 1  $\xi$  m^4  $\chi$ l / e,
  Pair[Momentum[FDp, D], Momentum[l, D]]  $\rightarrow$   $\tau$ 1 m^4  $\chi$ l / e,
  Pair[Momentum[FFp, D], Momentum[l, D]]  $\rightarrow$  -av2 kl kp} /.
  ktochiSubs[p, q, f, g, l] /. {av2  $\rightarrow$  -m^2  $\xi$ ^2 / e^2}
lpexpansion[ $\mu$ _] = Collect[ExpandVector[p, l,  $\mu$ ] /. subs $\rho$ 1 $\tau$ 1lp,
  {lv[ $\mu$ ], Flv[ $\mu$ ], FDllv[ $\mu$ ], FFlv[ $\mu$ ]}, Simplify]
lp = Contract[pv[ $\mu$ ]  $\times$  lv[ $\mu$ ] /. {lv[ $\mu$ ]  $\rightarrow$  lpexpansion[ $\mu$ ]}) /. subs $\rho$ 1 $\tau$ 1lp
```

$$\text{Out[44]} = \left\{ \text{Fp} \cdot l \rightarrow -\frac{m^4 \xi \rho^1 \chi^1}{e}, \text{FDp} \cdot l \rightarrow \frac{m^4 \tau^1 \chi^1}{e}, \text{FFp} \cdot l \rightarrow \frac{m^6 \chi^1 \chi p}{e^2} \right\}$$

$$\text{Out[45]} = \frac{e^2 \text{FFp}^\mu (\chi^1 (\xi^2 \rho^1 + \tau^1 - 1) + \chi p^2) - 2 e m^2 \tau^1 \chi^1 \chi p \text{FDp}^\mu + 2 m^2 \chi^1 \chi p (e \xi \rho^1 \text{Fp}^\mu + m^2 \chi p p^\mu)}{2 m^4 \chi^1 \chi p^3}$$

$$\text{Out[46]} = \frac{m^6 \chi p^2 (\chi^1 (\xi^2 \rho^1 + \tau^1 - 1) + \chi p^2) + 2 m^6 \chi^1 \chi p^2}{2 m^4 \chi^1 \chi p^3}$$

Projection of p on l,
 required later as an intermediate step (particularly, subs ρ 1 τ 1pl)
 Note that subs ρ 1 τ 1pl is not exactly
 the same as subs ρ 1 τ 1lp! We use both further

```
In[47]:= subsp1τ1pl = {Pair[Momentum[Fl, D], Momentum[p, D]] → ρ1 ξ m^4 χl / e,
  Pair[Momentum[FDl, D], Momentum[p, D]] → -τ1 m^4 χl / e,
  Pair[Momentum[FFl, D], Momentum[p, D]] → -av2 kl kp} /.
  ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}
plexpansion[μ_] = Collect[ExpandVector[l, p, μ] /. subsp1τ1pl,
  {lv[μ], Flv[μ], FDlv[μ], FFlv[μ]}, Simplify]
```

$$\text{Out[47]} = \left\{ \text{Fl} \cdot p \rightarrow \frac{m^4 \xi \rho_1 \chi^1}{e}, \text{FDl} \cdot p \rightarrow -\frac{m^4 \tau_1 \chi^1}{e}, \text{FFl} \cdot p \rightarrow \frac{m^6 \chi^1 \chi^p}{e^2} \right\}$$

$$\text{Out[48]} = \frac{e^2 \text{FFl}^\mu (\chi^1{}^2 (\xi^2 \rho_1{}^2 + \tau_1{}^2 + 1) - \chi^p{}^2)}{2 m^4 \chi^1{}^3 \chi^p} + \frac{e \tau_1 \text{FDl}^\mu}{m^2 \chi^1} - \frac{e \xi \rho_1 \text{Fl}^\mu}{m^2 \chi^1} + \frac{\chi^p l^\mu}{\chi^1}$$

(fg)
 -project g on f
 -contract with f, reexpress via ρ_n , τ_n

```
In[49]:= subsp2τ2gf = {Pair[Momentum[Ff, D], Momentum[g, D]] → -ρ2 ξ m^4 χl / e,
  Pair[Momentum[FDf, D], Momentum[g, D]] → -τ2 m^4 χl / e}
gfexpansion[μ_] = Collect[ExpandVector[f, g, μ] /. subsp2τ2gf,
  {fv[μ], Ffv[μ], FDFv[μ], FFFv[μ]}, Simplify]
fg = Contract[gfexpansion[μ] × fv[μ]]
```

$$\text{Out[49]} = \left\{ \text{Ff} \cdot g \rightarrow -\frac{m^4 \xi \rho_2 \chi^1}{e}, \text{FDf} \cdot g \rightarrow -\frac{m^4 \tau_2 \chi^1}{e} \right\}$$

$$\text{Out[50]} = \frac{e^2 \text{FFF}^\mu (\chi^1{}^2 (\xi^2 \rho_2{}^2 + \tau_2{}^2) + \chi^f{}^2 - \chi^g{}^2)}{2 m^4 \chi^1{}^3 \chi^g} + \frac{e \tau_2 \chi^1 \text{FDf}^\mu}{m^2 \chi^1{}^2} + \frac{e \xi \rho_2 \chi^1 \text{Ff}^\mu}{m^2 \chi^1{}^2} + \frac{\chi^g f^\mu}{\chi^f}$$

$$\text{Out[51]} = \frac{m^2 (\chi^1{}^2 (\xi^2 \rho_2{}^2 + \tau_2{}^2) + \chi^f{}^2 - \chi^g{}^2)}{2 \chi^f \chi^g} + \frac{m^2 \chi^g}{\chi^f}$$

Projection of g on l, required later as an intermediate step

```
In[52]:= subsp2τ2gl = {Pair[Momentum[Fl, D], Momentum[g, D]] → -ρ2 ξ m^4 χl / e,
  Pair[Momentum[FDl, D], Momentum[g, D]] → -τ2 m^4 χl / e}
glexpansion[μ_] = Collect[ExpandVector[l, g, μ] /. subsp2τ2gl,
  {lv[μ], Flv[μ], FDlv[μ], FFlv[μ]}, Simplify]
```

$$\text{Out[52]} = \left\{ \text{Fl} \cdot g \rightarrow -\frac{m^4 \xi \rho_2 \chi^1}{e}, \text{FDl} \cdot g \rightarrow -\frac{m^4 \tau_2 \chi^1}{e} \right\}$$

$$\text{Out[53]} = -\frac{e^2 \text{FFl}^\mu (\chi^g{}^2 - \chi^1{}^2 (\xi^2 \rho_2{}^2 + \tau_2{}^2 + 1))}{2 m^4 \chi^g \chi^1{}^3} + \frac{e \tau_2 \text{FDl}^\mu}{m^2 \chi^1} + \frac{e \xi \rho_2 \text{Fl}^\mu}{m^2 \chi^1} + \frac{\chi^g l^\mu}{\chi^1}$$

(pg)
 -project g on l
 -contract with p, reexpress via ρ_n ,
 τ_n using also substitutions subsp1τ1pl

and expression for (lp) calculated above

```
In[54]:= gpexpansion[μ_] =
  Collect[ExpandVector[p, g, μ], {pv[μ], Fpv[μ], FDpv[μ], FFpv[μ]}, Simplify]
pg = Simplify[Contract[pv[μ] × gv[μ] /. {gv[μ] → glexpansion[μ]}] /. subsρ1τ1pl /.
  {Contract[pv[μ] × lv[μ]] → lp}]
gp = pg;
```

$$\text{Out[54]} = \frac{\text{FFp}^\mu (e^4 (\text{FDp} \cdot g)^2 + e^4 (\text{Fp} \cdot g)^2 + e^2 m^8 (\chi p^2 - \chi g^2))}{2 m^{12} \chi g \chi p^3} - \frac{e^2 \text{FDp}^\mu (\text{FDp} \cdot g)}{m^6 \chi p^2} - \frac{e^2 \text{Fp}^\mu (\text{Fp} \cdot g)}{m^6 \chi p^2} + \frac{\chi g p^\mu}{\chi p}$$

$$\text{Out[55]} = \frac{m^2 (\chi g^2 (\xi^2 \rho_1^2 + \tau_1^2 + 1) + 2 \chi g \chi p (\xi^2 \rho_1 \rho_2 - \tau_1 \tau_2) + \chi p^2 (\xi^2 \rho_2^2 + \tau_2^2 + 1))}{2 \chi g \chi p}$$

(qg)
 – project q on p (calculated above)
 – project g on l (calculated above)
 – contract, reexpress via ρ_n, τ_n
 – the result will contain (lp),
 which we calculated above and substitute here;
 – In the last step,
 we remove all terms containing $(F)^{n>2}$ and $(F^* F)$ as they are zero

```
In[57]:= qg = Simplify[
  Contract[qv[μ] × gv[μ] /. {qv[μ] → qpexpansion[μ], gv[μ] → glexpansion[μ]}] /.
    subsρ1τ1qp /. subsρ1τ1pl /. subsρ1τ1lp /. {Contract[pv[μ] × lv[μ]] →
      lp} /. {Pair[Momentum[FFp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[FFp, D], Momentum[FDl, D]] → 0,
      Pair[Momentum[FDp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[FFp, D], Momentum[Ffl, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[FDl, D]] → 0,
      Pair[Momentum[FDp, D], Momentum[Ffl, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[Ffl, D]] → av2 kp kl,
      Pair[Momentum[FDp, D], Momentum[FDl, D]] → av2 kp kl} /.
    ktouchSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}]
gq = qg;
```

$$\text{Out[57]} = \frac{m^2 (\xi^2 (\rho_1 \chi g (\chi l + \chi q) + \rho_2 \chi p \chi q)^2 + \tau_1^2 \chi g^2 (\chi l + \chi q)^2 - 2 \tau_1 \tau_2 \chi g \chi p \chi q (\chi l + \chi q) + \chi p^2 ((\tau_2^2 + 1) \chi q^2 + \chi g^2))}{2 \chi g \chi p^2 \chi q}$$

Projection of f on l, required later as an intermediate step

```
In[59]:= subsp2τ2fl = {Pair[Momentum[Fl, D], Momentum[f, D]] → ρ2 ξ m^4 χl / e,
  Pair[Momentum[FDl, D], Momentum[f, D]] → τ2 m^4 χl / e,
  Pair[Momentum[FFl, D], Momentum[f, D]] → -av2 kl kf} /.
  ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}
flexpansion[μ_] = Collect[ExpandVector[l, f, μ] /. subsp2τ2fl,
  {lv[μ], Flv[μ], Fdlv[μ], FFlv[μ]}, Simplify]
```

$$\text{Out[59]} = \left\{ f \cdot Fl \rightarrow \frac{m^4 \xi \rho^2 \chi^l}{e}, f \cdot FDI \rightarrow \frac{m^4 \tau^2 \chi^l}{e}, f \cdot FFI \rightarrow \frac{m^6 \chi^f \chi^l}{e^2} \right\}$$

$$\text{Out[60]} = -\frac{e^2 FFI^\mu (\chi^{f^2} - \chi^{l^2} (\xi^2 \rho^2 + \tau^2 + 1))}{2 m^4 \chi^f \chi^{l^3}} - \frac{e \tau^2 FDI^\mu}{m^2 \chi^l} - \frac{e \xi \rho^2 FI^\mu}{m^2 \chi^l} + \frac{\chi^f l^\mu}{\chi^l}$$

(pf)
 -project f on l
 -contract, reexpress via ρ_n, τ_n ,
 also using subsp1τ1pl and scalar product (lp) calculated before

```
In[61]:= fpexpansion[μ_] =
  Collect[ExpandVector[p, f, μ], {pv[μ], Fpv[μ], FDpv[μ], FFpv[μ]}, Simplify]
pf = Simplify[Contract[pv[μ] × fv[μ] /. {fv[μ] → flexpansion[μ]}] /. subsp1τ1pl /.
  {Contract[pv[μ] × lv[μ]] → lp}]
fp = pf;
```

$$\text{Out[61]} = \frac{FFp^\mu (e^4 (f \cdot FDp)^2 + e^4 (f \cdot Fp)^2 + e^2 m^8 (\chi p^2 - \chi^f))}{2 m^{12} \chi^f \chi p^3} - \frac{e^2 FDp^\mu (f \cdot FDp)}{m^6 \chi p^2} - \frac{e^2 Fp^\mu (f \cdot Fp)}{m^6 \chi p^2} + \frac{\chi^f p^\mu}{\chi p}$$

$$\text{Out[62]} = \frac{m^2 (\chi^{f^2} (\xi^2 \rho^2 + \tau^2 + 1) + \chi^f (2 \tau^1 \tau^2 \chi p - 2 \xi^2 \rho^1 \rho^2 \chi p) + \chi p^2 (\xi^2 \rho^2 + \tau^2 + 1))}{2 \chi^f \chi p}$$

(qf)
 -project q on p (calculated above)
 -project f on l (calculated above)
 -contract, reexpress via ρ_n, τ_n ,
 -the result will contain (lp),
 which we calculated above and substitute here;
 -In the last step,
 we remove all terms containing $(F)^{n>2}$ and $(F^* F)$ as they are zero

```

In[64]:= qf = Simplify[
  Contract[qv[μ] × fv[μ] /. {qv[μ] → qpexpansion[μ], fv[μ] → flexpansion[μ]}] /.
    subsρ1τ1qp /. subsρ1τ1pl /. subsρ1τ1lp /. {Contract[pv[μ] × lv[μ] →
      lp} /. {Pair[Momentum[FFp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[FFp, D], Momentum[FDl, D]] → 0,
      Pair[Momentum[FDp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[FFl, D]] → 0,
      Pair[Momentum[FFp, D], Momentum[Ff, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[FDl, D]] → 0,
      Pair[Momentum[FDp, D], Momentum[Ff, D]] → 0,
      Pair[Momentum[Fp, D], Momentum[Ff, D]] → av2 kp kl,
      Pair[Momentum[FDp, D], Momentum[FDl, D]] → av2 kp kl} /.
    ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}]
fq = qf;

```

$$\text{Out[64]= } \frac{m^2 (\xi^2 (\rho^2 \chi p \chi q - \rho^1 \chi f (\chi^1 + \chi q))^2 + \tau^1{}^2 \chi f^2 (\chi^1 + \chi q)^2 + 2 \tau^1 \tau^2 \chi f \chi p \chi q (\chi^1 + \chi q) + \chi p^2 ((\tau^2{}^2 + 1) \chi q^2 + \chi f^2))}{2 \chi f \chi p^2 \chi q}$$

All scalar products of the form

(aq), (al), ...

can be reduced to only two : (ap) or (af) plus invariant terms,

This is achieved again by projecting 4 –

momenta on the basis spanned by F and p or f

$$q^\mu = C_1 p^\mu + C_2 (Fp)^\mu + C_3 (F^* p)^\mu + C_4 (FFp)^\mu$$

If done correctly,

all terms containing (ap) and (af) in variable Expr will cancel ,
and the expression will depend only on gauge – invariant parameters

The result is stored in Expr4

```

In[66]:= aqsubs = {Pair[Momentum[a, D], Momentum[q, D]] → Contract[av[μ] × qpexpansion[μ]] /.
    ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}}
alsubs = {Pair[Momentum[a, D], Momentum[l, D]] → Contract[av[μ] × lpexpansion[μ]] /.
    ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}}
agsubs = {Pair[Momentum[a, D], Momentum[g, D]] → Contract[av[μ] × glexpansion[μ]] /.
    alsubs /. ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}}
afsubs = {Pair[Momentum[a, D], Momentum[f, D]] → Contract[av[μ] × flxpansion[μ]] /.
    alsubs /. ktochiSubs[p, q, f, g, l] /. {av2 → -m^2 ξ^2 / e^2}}

```

$$\text{Out[66]} = \left\{ a \cdot q \rightarrow \frac{\chi q (a \cdot p)}{\chi p} - \frac{m^2 \xi^2 \rho 1 \chi 1}{e \chi p} \right\}$$

$$\text{Out[67]} = \left\{ a \cdot l \rightarrow \frac{\chi 1 \left(m^2 \chi p (a \cdot p) + \frac{m^4 \xi^2 \rho 1 \chi p}{e} \right)}{m^2 \chi p^2} \right\}$$

$$\text{Out[68]} = \left\{ a \cdot g \rightarrow \frac{\chi g \left(m^2 \chi p (a \cdot p) + \frac{m^4 \xi^2 \rho 1 \chi p}{e} \right)}{m^2 \chi p^2} + \frac{m^2 \xi^2 \rho 2}{e} \right\}$$

$$\text{Out[69]} = \left\{ a \cdot f \rightarrow \frac{\chi f \left(m^2 \chi p (a \cdot p) + \frac{m^4 \xi^2 \rho 1 \chi p}{e} \right)}{m^2 \chi p^2} - \frac{m^2 \xi^2 \rho 2}{e} \right\}$$

```

In[70]:= Expr3 /. aqsubs /. agsubs /. afsubs /. {Pair[Momentum[p, D], Momentum[q, D]] → pq,
    Pair[Momentum[p, D], Momentum[f, D]] → pf,
    Pair[Momentum[p, D], Momentum[g, D]] → pg,
    Pair[Momentum[q, D], Momentum[f, D]] → qf,
    Pair[Momentum[q, D], Momentum[g, D]] → qg,
    Pair[Momentum[f, D], Momentum[g, D]] → fg};
Expand[%];
Expr4 = SortTermsSimplify[Contract[%] /. ktochiSubs[p, q, f, g] /.
    {av2 → -m^2 ξ^2 / e^2} (* /. {χq→χp-χl, χg→χl-χg} *) ];
% /. {D → 4}

```

$$\begin{aligned} \text{Out[73]} = & \frac{16 A2 B2 C0 D0 m^4 \chi f \chi g \xi^4}{\chi p \chi q} + \\ & A1 B2 \left(\frac{8 C1 D0 m^4 (\chi f - \chi g) (\chi p + \chi q) \xi^4}{\chi p \chi q} + \frac{8 C0 D0 m^4 (\rho 2 (\chi f - \chi g) \chi p (\chi p + \chi q) + 2 \rho 1 \chi f \chi g (\chi 1 + \chi p + \chi q)) \xi^4}{\chi p^2 \chi q} \right) + \\ & A2 B1 \left(\frac{8 C0 D1 m^4 (\chi f - \chi g) (\chi p + \chi q) \xi^4}{\chi p \chi q} + \frac{8 C0 D0 m^4 (\rho 2 (\chi f - \chi g) \chi p (\chi p + \chi q) + 2 \rho 1 \chi f \chi g (\chi 1 + \chi p + \chi q)) \xi^4}{\chi p^2 \chi q} \right) + \\ & A0 B2 \left(-16 C2 D0 \xi^4 m^4 + \frac{8 C1 D0 \xi^4 (\rho 1 (\chi f - \chi g) (\chi 1 + 2 \chi q) - 4 \rho 2 \chi p \chi q) m^4}{\chi p \chi q} + \frac{1}{\chi f \chi g \chi p^2 \chi q} \right. \\ & \quad 8 C0 D0 \xi^2 ((2 \rho 1^2 \chi f^2 (\chi 1 + \chi q) \chi g^2 + \rho 1 \rho 2 \chi f (\chi f - \chi g) \chi p (\chi 1 + 2 \chi q) \chi g + \rho 2^2 (\chi f^2 + \chi g^2 - \chi 1^2) \chi p^2 \chi q) \xi^2 + \\ & \quad 2 \tau 1^2 \chi f^2 \chi g^2 (\chi 1 + \chi q) - \tau 1 \tau 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi 1 + 2 \chi q) + \\ & \quad \left. \chi p ((\chi g^2 - \chi 1^2) \chi p \chi q \tau 2^2 - 2 \chi f \chi g \chi p \chi q + \chi f^2 (\chi p \chi q \tau 2^2 + 2 \chi g^2))) m^4 \right) + \end{aligned}$$

$$\begin{aligned}
& \text{A2 B0} \left(-16 \text{C0 D2 } \xi^4 m^4 + \frac{8 \text{C0 D1 } \xi^4 (\rho 1 (\chi f - \chi g) (\chi l + 2 \chi q) - 4 \rho 2 \chi p \chi q) m^4}{\chi p \chi q} + \frac{1}{\chi f \chi g \chi p^2 \chi q} \right. \\
& \quad 8 \text{C0 D0 } \xi^2 ((2 \rho 1^2 \chi f^2 (\chi l + \chi q) \chi g^2 + \rho 1 \rho 2 \chi f (\chi f - \chi g) \chi p (\chi l + 2 \chi q) \chi g + \rho 2^2 (\chi f^2 + \chi g^2 - \chi l^2) \chi p^2 \chi q) \xi^2 + \\
& \quad 2 \tau 1^2 \chi f^2 \chi g^2 (\chi l + \chi q) - \tau 1 \tau 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi l + 2 \chi q) + \\
& \quad \left. \chi p ((\chi g^2 - \chi l^2) \chi p \chi q \tau 2^2 - 2 \chi f \chi g \chi p \chi q + \chi f^2 (\chi p \chi q \tau 2^2 + 2 \chi g^2))) m^4 \right) + \\
& \text{A1 B1} \left(\frac{4 \text{C1 D1 } \xi^4 ((2 \chi p^2 + 2 \chi q^2) \chi f^2 - 8 \chi g \chi p \chi q \chi f + \chi g^2 (2 \chi p^2 + 2 \chi q^2)) m^4}{\chi f \chi g \chi p \chi q} + \frac{1}{\chi f \chi g \chi p^2 \chi q} \right. \\
& \quad 4 \text{C1 D0 } \xi^4 (2 \rho 1 \chi f (\chi f - \chi g) \chi g (\chi p + \chi q) (\chi l + \chi p + \chi q) + \\
& \quad \rho 2 \chi p ((2 \chi p^2 + 2 \chi q^2) \chi f^2 - 8 \chi g \chi p \chi q \chi f + \chi g^2 (2 \chi p^2 + 2 \chi q^2))) m^4 + \\
& \quad \frac{1}{\chi f \chi g \chi p^2 \chi q} 4 \text{C0 D1 } \xi^4 (2 \rho 1 \chi f (\chi f - \chi g) \chi g (\chi p + \chi q) (\chi l + \chi p + \chi q) + \\
& \quad \rho 2 \chi p ((2 \chi p^2 + 2 \chi q^2) \chi f^2 - 8 \chi g \chi p \chi q \chi f + \chi g^2 (2 \chi p^2 + 2 \chi q^2))) m^4 + \frac{1}{\chi f \chi g \chi p^3 \chi q} \\
& \quad 4 \text{C0 D0 } \xi^2 ((\rho 2^2 (4 \chi p \chi q \chi l^2 + 2 \chi f^2 (\chi p - \chi q)^2 + 2 \chi g^2 (\chi p - \chi q)^2 - 16 \chi f \chi g \chi p \chi q) \chi p^2 + \\
& \quad 4 \rho 1 \rho 2 \chi f (\chi f - \chi g) \chi g (\chi p + \chi q) (\chi l + \chi p + \chi q) \chi p + 4 \rho 1^2 \chi f^2 \chi g^2 (\chi l + \chi p + \chi q)^2) \xi^2 + \\
& \quad 4 \tau 1^2 \chi f^2 \chi g^2 (\chi l - \chi p + \chi q)^2 + 4 \tau 1 \tau 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi p - \chi q) (\chi l - \chi p + \chi q) + \\
& \quad \chi p^2 ((-4 \chi p \chi q \tau 2^2 + (2 \tau 2^2 + 2) \chi p^2 + (2 \tau 2^2 + 2) \chi q^2) \chi f^2 + 2 \chi g (2 \chi p^2 + 2 \chi q^2) \chi f + \\
& \quad \left. 4 \tau 2^2 \chi l^2 \chi p \chi q + \chi g^2 (-4 \chi p \chi q \tau 2^2 + (2 \tau 2^2 + 2) \chi p^2 + (2 \tau 2^2 + 2) \chi q^2))) m^4 \right) + \\
& \text{A1 B0} \left(-\frac{8 \text{D2 } \xi^4 (\text{C1 } (\chi f - \chi g) \chi p (\chi p + \chi q) + \text{C0 } \rho 2 (\chi f - \chi g) \chi p (\chi p + \chi q) + 2 \text{C0 } \rho 1 \chi f \chi g (\chi l + \chi p + \chi q)) m^4}{\chi f \chi g \chi p} + \right. \\
& \quad \frac{1}{\chi f \chi g \chi p \chi q} 4 \text{C1 D1 } \xi^4 \\
& \quad (\rho 1 ((2 \chi l \chi p + 2 \chi q (\chi p + \chi q)) \chi f^2 + 2 \chi g (-2 \chi l \chi q - 2 (\chi p + \chi q) \chi q) \chi f + \chi g^2 (2 \chi l \chi p + 2 \chi q (\chi p + \chi q))) - \\
& \quad 4 \rho 2 (\chi f - \chi g) \chi p \chi q (\chi p + \chi q)) m^4 + \frac{1}{\chi f^2 \chi g^2 \chi p^2 \chi q} \\
& \quad 4 \text{C0 D1 } \xi^2 ((2 \rho 1^2 \chi f^2 (\chi f - \chi g) (\chi l + \chi p + \chi q) (\chi l + 2 \chi q) \chi g^2 + \rho 1 \rho 2 \chi f \chi p ((2 \chi l \chi p + 2 \chi q (\chi p + \chi q)) \chi f^2 + \\
& \quad 2 \chi g (-6 \chi l \chi q - 6 (\chi p + \chi q) \chi q) \chi f + \chi g^2 (2 \chi l \chi p + 2 \chi q (\chi p + \chi q))) \chi g - \\
& \quad \rho 2^2 (\chi f - \chi g) (\chi f^2 + 6 \chi g \chi f + \chi g^2 - \chi l^2) \chi p^2 \chi q (\chi p + \chi q)) \xi^2 + \\
& \quad 2 \tau 1^2 \chi f^2 (\chi f - \chi g) \chi g^2 \chi l (\chi l - \chi p + \chi q) - \tau 2^2 (\chi f - \chi g) (\chi f^2 + 2 \chi g \chi f + \chi g^2 - \chi l^2) \chi p^2 \chi q (\chi p + \chi q) + \\
& \quad 2 \tau 1 \tau 2 \chi f \chi g \chi p ((\chi l \chi p + \chi q (\chi p + \chi q)) \chi f^2 + 2 \chi g \chi q (\chi l + \chi p + \chi q) \chi f + \chi g^2 (\chi l \chi p + \chi q (\chi p + \chi q))) m^4 + \\
& \quad \frac{1}{\chi f^2 \chi g^2 \chi p^2 \chi q} 4 \text{C1 D0 } \xi^2 ((2 \rho 1^2 \chi f^2 (\chi f - \chi g) (\chi l + \chi q) (\chi p + \chi q) \chi g^2 + \rho 1 \rho 2 \chi f \chi p ((2 \chi l \chi p + 2 \chi q (\chi p + \chi q)) \\
& \quad \chi f^2 + 2 \chi g (-2 \chi l \chi q - 2 (\chi p + \chi q) \chi q) \chi f + \chi g^2 (2 \chi l \chi p + 2 \chi q (\chi p + \chi q))) \chi g + \\
& \quad \rho 2^2 (\chi f - \chi g) (\chi f^2 + \chi g^2 - \chi l^2) \chi p^2 \chi q (\chi p + \chi q)) \xi^2 + 2 \tau 1^2 \chi f^2 (\chi f - \chi g) \chi g^2 (\chi l + \chi q) (\chi p + \chi q) + \\
& \quad (\chi f - \chi g) \chi p (\chi p + \chi q) ((\chi g^2 - \chi l^2) \chi p \chi q \tau 2^2 - 2 \chi f \chi g \chi p \chi q + \chi f^2 (\chi p \chi q \tau 2^2 + 2 \chi g^2)) - \\
& \quad 2 \tau 1 \tau 2 \chi f \chi g \chi p ((\chi l \chi p + \chi q (\chi p + \chi q)) \chi f^2 - 2 \chi g \chi q (\chi l + \chi p + \chi q) \chi f + \chi g^2 (\chi l \chi p + \chi q (\chi p + \chi q))) \\
& \quad m^4 + \frac{1}{\chi f \chi g \chi p^3 \chi q} 4 \text{C0 D0 } \xi^2 \\
& \quad ((4 \chi f^2 \chi g^2 (\chi l + \chi q) (\chi l + \chi p + \chi q) \rho 1^3 + 2 \rho 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi l^2 + 2 (\chi p + 2 \chi q) \chi l + 3 \chi q (\chi p + \chi q)) \rho 1^2 + \\
& \quad \rho 2^2 \chi p^2 (2 (\chi l \chi p + \chi q (\chi p + \chi q)) \chi f^2 - 8 \chi g \chi q (\chi l + \chi p + \chi q) \chi f + 2 \chi g^2 (\chi l \chi p + \chi q (\chi p + \chi q))) \rho 1 -
\end{aligned}$$

$$\begin{aligned}
& 2\rho^2{}^3(\chi f - \chi g)\chi p^3\chi q(\chi p + \chi q))\xi^2 + 2\rho^2\chi p(\chi f(\chi f - \chi g)\chi g(\chi l^2 + 2\chi q\chi l + \chi q(\chi p + \chi q))\tau l^2 + \\
& 4\tau^2\chi f\chi g\chi p\chi q(\chi l + \chi p + \chi q)\tau l + (\chi f - \chi g)\chi p(\chi p + \chi q)(\chi f\chi g - (\tau^2 + 1)\chi p\chi q)) + \\
& \rho l(4\tau l^2\chi f^2(\chi l + \chi q)(\chi l + \chi p + \chi q)\chi g^2 - 4\tau l\tau^2\chi f(\chi f - \chi g)\chi p(\chi l + \chi q)(\chi p + \chi q)\chi g + \\
& \chi p((4(\chi l + \chi p + \chi q)\chi g^2 + \chi p((2\tau^2 + 2)\chi l\chi p + 2(\tau^2 + 1)\chi q(\chi p + \chi q)))\chi f^2 + \\
& 4\chi g\chi l\chi p(\chi p - \chi q)\chi f + \chi g^2\chi p((2\tau^2 + 2)\chi l\chi p + 2(\tau^2 + 1)\chi q(\chi p + \chi q))))m^4) + \\
\text{A0 B1} & \left(-\frac{8\text{C2 D1}\xi^4(\chi f - \chi g)(\chi p + \chi q)m^4}{\chi f\chi g} - \frac{8\text{C2 D0}\xi^4(\rho^2(\chi f - \chi g)\chi p(\chi p + \chi q) + 2\rho l\chi f\chi g(\chi l + \chi p + \chi q))m^4}{\chi f\chi g\chi p} + \right. \\
& \frac{1}{\chi f\chi g\chi p\chi q}4\text{C1 D1}\xi^4 \\
& (\rho l((2\chi l\chi p + 2\chi q(\chi p + \chi q))\chi f^2 + 2\chi g(-2\chi l\chi q - 2(\chi p + \chi q)\chi q)\chi f + \chi g^2(2\chi l\chi p + 2\chi q(\chi p + \chi q))) - \\
& 4\rho^2(\chi f - \chi g)\chi p\chi q(\chi p + \chi q))m^4 + \frac{1}{\chi f^2\chi g^2\chi p^2\chi q} \\
& 4\text{C1 D0}\xi^2((2\rho l^2\chi f^2(\chi f - \chi g)(\chi l + \chi p + \chi q)(\chi l + 2\chi q)\chi g^2 + \rho l\rho^2\chi f\chi p((2\chi l\chi p + 2\chi q(\chi p + \chi q))\chi f^2 + \\
& 2\chi g(-6\chi l\chi q - 6(\chi p + \chi q)\chi q)\chi f + \chi g^2(2\chi l\chi p + 2\chi q(\chi p + \chi q)))\chi g - \\
& \rho^2(\chi f - \chi g)(\chi f^2 + 6\chi g\chi f + \chi g^2 - \chi l^2)\chi p^2\chi q(\chi p + \chi q))\xi^2 + \\
& 2\tau l^2\chi f^2(\chi f - \chi g)\chi g^2\chi l(\chi l - \chi p + \chi q) - \tau^2(\chi f - \chi g)(\chi f^2 + 2\chi g\chi f + \chi g^2 - \chi l^2)\chi p^2\chi q(\chi p + \chi q) + \\
& 2\tau l\tau^2\chi f\chi g\chi p((\chi l\chi p + \chi q(\chi p + \chi q))\chi f^2 + 2\chi g\chi q(\chi l + \chi p + \chi q)\chi f + \chi g^2(\chi l\chi p + \chi q(\chi p + \chi q))))m^4 + \\
& \frac{1}{\chi f^2\chi g^2\chi p^2\chi q}4\text{C0 D1}\xi^2((2\rho l^2\chi f^2(\chi f - \chi g)(\chi l + \chi q)(\chi p + \chi q)\chi g^2 + \rho l\rho^2\chi f\chi p((2\chi l\chi p + 2\chi q(\chi p + \chi q)) \\
& \chi f^2 + 2\chi g(-2\chi l\chi q - 2(\chi p + \chi q)\chi q)\chi f + \chi g^2(2\chi l\chi p + 2\chi q(\chi p + \chi q)))\chi g + \\
& \rho^2(\chi f - \chi g)(\chi f^2 + \chi g^2 - \chi l^2)\chi p^2\chi q(\chi p + \chi q))\xi^2 + 2\tau l^2\chi f^2(\chi f - \chi g)\chi g^2(\chi l + \chi q)(\chi p + \chi q) + \\
& (\chi f - \chi g)\chi p(\chi p + \chi q)((\chi g^2 - \chi l^2)\chi p\chi q\tau^2 - 2\chi f\chi g\chi p\chi q + \chi f^2(\chi p\chi q\tau^2 + 2\chi g^2)) - \\
& 2\tau l\tau^2\chi f\chi g\chi p((\chi l\chi p + \chi q(\chi p + \chi q))\chi f^2 - 2\chi g\chi q(\chi l + \chi p + \chi q)\chi f + \chi g^2(\chi l\chi p + \chi q(\chi p + \chi q)))) \\
& m^4 + \frac{1}{\chi f\chi g\chi p^3\chi q}4\text{C0 D0}\xi^2 \\
& ((4\chi f^2\chi g^2(\chi l + \chi q)(\chi l + \chi p + \chi q)\rho l^3 + 2\rho^2\chi f(\chi f - \chi g)\chi g\chi p(\chi l^2 + 2(\chi p + 2\chi q)\chi l + 3\chi q(\chi p + \chi q))\rho l^2 + \\
& \rho^2\chi p^2(2(\chi l\chi p + \chi q(\chi p + \chi q))\chi f^2 - 8\chi g\chi q(\chi l + \chi p + \chi q)\chi f + 2\chi g^2(\chi l\chi p + \chi q(\chi p + \chi q)))\rho l - \\
& 2\rho^2{}^3(\chi f - \chi g)\chi p^3\chi q(\chi p + \chi q))\xi^2 + 2\rho^2\chi p(\chi f(\chi f - \chi g)\chi g(\chi l^2 + 2\chi q\chi l + \chi q(\chi p + \chi q))\tau l^2 + \\
& 4\tau^2\chi f\chi g\chi p\chi q(\chi l + \chi p + \chi q)\tau l + (\chi f - \chi g)\chi p(\chi p + \chi q)(\chi f\chi g - (\tau^2 + 1)\chi p\chi q)) + \\
& \rho l(4\tau l^2\chi f^2(\chi l + \chi q)(\chi l + \chi p + \chi q)\chi g^2 - 4\tau l\tau^2\chi f(\chi f - \chi g)\chi p(\chi l + \chi q)(\chi p + \chi q)\chi g + \\
& \chi p((4(\chi l + \chi p + \chi q)\chi g^2 + \chi p((2\tau^2 + 2)\chi l\chi p + 2(\tau^2 + 1)\chi q(\chi p + \chi q)))\chi f^2 + \\
& 4\chi g\chi l\chi p(\chi p - \chi q)\chi f + \chi g^2\chi p((2\tau^2 + 2)\chi l\chi p + 2(\tau^2 + 1)\chi q(\chi p + \chi q))))m^4) + \\
\text{A0 B0} & \left(\frac{16\text{C2 D2}\xi^4\chi p\chi qm^4}{\chi f\chi g} - \frac{8\text{C2 D1}\xi^4(\rho l(\chi f - \chi g)(\chi l + 2\chi q) - 4\rho^2\chi p\chi q)m^4}{\chi f\chi g} - \right. \\
& \frac{1}{\chi f^2\chi g^2\chi p} \\
& 8\text{C2 D0}\xi^2((2\rho l^2\chi f^2(\chi l + \chi q)\chi g^2 + \rho l\rho^2\chi f(\chi f - \chi g)\chi p(\chi l + 2\chi q)\chi g + \rho^2(\chi f^2 + \chi g^2 - \chi l^2)\chi p^2\chi q)\xi^2 + \\
& 2\tau l^2\chi f^2\chi g^2(\chi l + \chi q) - \tau l\tau^2\chi f(\chi f - \chi g)\chi g\chi p(\chi l + 2\chi q) + \\
& \chi p((\chi g^2 - \chi l^2)\chi p\chi q\tau^2 - 2\chi f\chi g\chi p\chi q + \chi f^2(\chi p\chi q\tau^2 + 2\chi g^2)))m^4 - \\
& \frac{1}{\chi f^2\chi g^2\chi p}8\text{D2}\xi^2(\text{C1}\chi f\chi g\chi p(\rho l(\chi f - \chi g)(\chi l + 2\chi q) - 4\rho^2\chi p\chi q)\xi^2 +
\end{aligned}$$

$$\begin{aligned}
& C0 \left((2\rho 1^2 \chi f^2 (\chi l + \chi q) \chi g^2 + \rho 1 \rho 2 \chi f (\chi f - \chi g) \chi p (\chi l + 2\chi q) \chi g + \rho 2^2 (\chi f^2 + \chi g^2 - \chi l^2) \chi p^2 \chi q) \xi^2 + \right. \\
& \quad 2\tau 1^2 \chi f^2 \chi g^2 (\chi l + \chi q) - \tau 1 \tau 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi l + 2\chi q) + \\
& \quad \left. \chi p ((\chi g^2 - \chi l^2) \chi p \chi q \tau 2^2 - 2\chi f \chi g \chi p \chi q + \chi f^2 (\chi p \chi q \tau 2^2 + 2\chi g^2)) \right) m^4 + \\
& \frac{1}{\chi f^2 \chi g^2 \chi p \chi q} 4 C1 D1 \xi^2 (2\chi g \chi p^2 \chi f^3 + 2\chi g \chi q^2 \chi f^3 - 4\chi g \chi p \chi q \chi f^3 + 4\tau 2^2 \chi p^2 \chi q^2 \chi f^2 + \\
& \quad 2\chi g^3 \chi p^2 \chi f + 2\chi g^3 \chi q^2 \chi f + 8\tau 2^2 \chi g \chi p^2 \chi q^2 \chi f - 4\chi g^3 \chi p \chi q \chi f + \\
& \quad \tau 1^2 \chi g ((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + 2\chi g (4\chi q^2 + 4\chi l \chi q) \chi f + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \chi f + \\
& \quad 4\tau 2^2 \chi g^2 \chi p^2 \chi q^2 - 4\tau 2^2 \chi l^2 \chi p^2 \chi q^2 + \\
& \quad \xi^2 (\chi f \chi g ((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + 2\chi g (-4\chi q^2 - 4\chi l \chi q) \chi f + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \rho 1^2 - \\
& \quad 8\rho 2 \chi f (\chi f - \chi g) \chi g \chi p \chi q (\chi l + 2\chi q) \rho 1 + 4\rho 2^2 (\chi f^2 + 6\chi g \chi f + \chi g^2 - \chi l^2) \chi p^2 \chi q^2) m^4 + \\
& \frac{1}{\chi f \chi g \chi p^2 \chi q} 4 C1 D0 \xi^2 ((2\chi f (\chi f - \chi g) \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \rho 1^3 + \\
& \quad \rho 2 \chi p ((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + 2\chi g (-8\chi q^2 - 8\chi l \chi q) \chi f + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \rho 1^2 - \\
& \quad 6\rho 2^2 (\chi f - \chi g) \chi p^2 \chi q (\chi l + 2\chi q) \rho 1 + 8\rho 2^3 \chi p^3 \chi q^2) \xi^2 + \\
& \quad \rho 2 \chi p (((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \tau 1^2 + \\
& \quad 4\tau 2 (\chi f - \chi g) \chi p \chi q (\chi l + 2\chi q) \tau 1 + 2\chi g^2 \chi p^2 + 2\chi f^2 (\chi p - \chi q)^2 + \\
& \quad 2\chi g^2 \chi q^2 + 8\tau 2^2 \chi p^2 \chi q^2 + 8\chi p^2 \chi q^2 - 4\chi g^2 \chi p \chi q - 8\chi f \chi g \chi p \chi q) + \\
& \quad 2\rho 1 (\chi f (\chi f - \chi g) \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \tau 1^2 + 8\tau 2 \chi f \chi g \chi p \chi q (\chi l + \chi q) \tau 1 + \\
& \quad (\chi f - \chi g) (\chi f \chi g (\chi p^2 + \chi q (\chi l + \chi q)) - (\tau 2^2 + 1) \chi p^2 \chi q (\chi l + 2\chi q))) m^4 + \\
& \frac{1}{\chi f \chi g \chi p^2 \chi q} 4 C0 D1 \xi^2 ((2\chi f (\chi f - \chi g) \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \rho 1^3 + \\
& \quad \rho 2 \chi p ((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + 2\chi g (-8\chi q^2 - 8\chi l \chi q) \chi f + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \rho 1^2 - \\
& \quad 6\rho 2^2 (\chi f - \chi g) \chi p^2 \chi q (\chi l + 2\chi q) \rho 1 + 8\rho 2^3 \chi p^3 \chi q^2) \xi^2 + \\
& \quad \rho 2 \chi p (((2\chi l^2 + 4\chi q \chi l + 4\chi q^2) \chi f^2 + \chi g^2 (2\chi l^2 + 4\chi q \chi l + 4\chi q^2)) \tau 1^2 + \\
& \quad 4\tau 2 (\chi f - \chi g) \chi p \chi q (\chi l + 2\chi q) \tau 1 + 2\chi g^2 \chi p^2 + 2\chi f^2 (\chi p - \chi q)^2 + \\
& \quad 2\chi g^2 \chi q^2 + 8\tau 2^2 \chi p^2 \chi q^2 + 8\chi p^2 \chi q^2 - 4\chi g^2 \chi p \chi q - 8\chi f \chi g \chi p \chi q) + \\
& \quad 2\rho 1 (\chi f (\chi f - \chi g) \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \tau 1^2 + 8\tau 2 \chi f \chi g \chi p \chi q (\chi l + \chi q) \tau 1 + \\
& \quad (\chi f - \chi g) (\chi f \chi g (\chi p^2 + \chi q (\chi l + \chi q)) - (\tau 2^2 + 1) \chi p^2 \chi q (\chi l + 2\chi q))) m^4 + \\
& \frac{1}{\chi f \chi g \chi p^3 \chi q} 4 C0 D0 ((4\chi f^2 \chi g^2 (\chi l + \chi q)^2 \rho 1^4 + 4\rho 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \rho 1^3 + \\
& \quad \rho 2^2 \chi p^2 (2(\chi l^2 + 2\chi q \chi l + 2\chi q^2) \chi f^2 - 16\chi g \chi q (\chi l + \chi q) \chi f + 2\chi g^2 (\chi l^2 + 2\chi q \chi l + 2\chi q^2)) \rho 1^2 - \\
& \quad 4\rho 2^3 (\chi f - \chi g) \chi p^3 \chi q (\chi l + 2\chi q) \rho 1 + 4\rho 2^4 \chi p^4 \chi q^2) \xi^4 + \\
& \quad (((4((2\tau 1^2 + 1)\chi l^2 + 2(2\chi q \tau 1^2 + \chi q)\chi l + \chi p^2 + 2\tau 1^2 \chi q^2 + \chi q^2) \chi g^2 - 4\tau 1 \tau 2 \chi p \\
& \quad (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \chi g + \chi p^2 ((2\tau 2^2 + 2)\chi l^2 + 4(\tau 2^2 + 1)\chi q \chi l + 4(\tau 2^2 + 1)\chi q^2)) \\
& \quad \chi f^2 + 2\chi g \chi p (2\chi p \chi l^2 + 2\tau 1 \tau 2 \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2)) \chi f + \\
& \quad \chi g^2 \chi p^2 ((2\tau 2^2 + 2)\chi l^2 + 4(\tau 2^2 + 1)\chi q \chi l + 4(\tau 2^2 + 1)\chi q^2)) \rho 1^2 + \\
& \quad 4\rho 2 \chi p (\chi f (\chi f - \chi g) \chi g (\chi l^2 + 3\chi q \chi l + 2\chi q^2) \tau 1^2 + 8\tau 2 \chi f \chi g \chi p \chi q (\chi l + \chi q) \tau 1 + \\
& \quad (\chi f - \chi g) (\chi f \chi g (\chi p^2 + \chi q (\chi l + \chi q)) - (\tau 2^2 + 1) \chi p^2 \chi q (\chi l + 2\chi q))) \rho 1 + \\
& \quad \rho 2^2 \chi p^2 ((2(\chi l^2 + 2\chi q \chi l + 2\chi q^2) \chi f^2 + 2\chi g^2 (\chi l^2 + 2\chi q \chi l + 2\chi q^2)) \tau 1^2 + \\
& \quad 4\tau 2 (\chi f - \chi g) \chi p \chi q (\chi l + 2\chi q) \tau 1 + 2\chi f^2 \chi p^2 + 2\chi g^2 \chi p^2 + \\
& \quad 2\chi f^2 \chi q^2 + 2\chi g^2 \chi q^2 + 8\tau 2^2 \chi p^2 \chi q^2 + 8\chi p^2 \chi q^2 - 4\chi l^2 \chi p \chi q)) \xi^2 + \\
& \quad 4\tau 1^4 \chi f^2 \chi g^2 (\chi l + \chi q)^2 - 4\tau 1^3 \tau 2 \chi f (\chi f - \chi g) \chi g \chi p (\chi l^2 + 3\chi q \chi l + 2\chi q^2) + \\
& \quad \chi p^2 (4\chi p^2 \chi q^2 \tau 2^4 + 8\chi p^2 \chi q^2 \tau 2^2 - 4\chi l^2 \chi p \chi q \tau 2^2 + 4\chi p^2 \chi q^2 + \\
& \quad 2\chi f \chi g (2\chi p^2 - 8\chi q \chi p + 2\chi q^2) + \chi g^2 ((2\tau 2^2 + 2)\chi p^2 - 4\chi q \chi p + (2\tau 2^2 + 2)\chi q^2) + \\
& \quad \chi f^2 (4\chi g^2 + (2\tau 2^2 + 2)\chi p^2 + (2\tau 2^2 + 2)\chi q^2 - 4\chi p \chi q)) +
\end{aligned}$$

$$\begin{aligned}
& 4 \tau_1 \tau_2 (\chi_f - \chi_g) \chi_p ((\tau^2 + 1) \chi_p^2 \chi_q (\chi_l + 2 \chi_q) - \chi_f \chi_g (\chi_p^2 + \chi_q (\chi_l + \chi_q))) + \\
& \tau_l^2 ((4 (\chi_l^2 + 2 \chi_q \chi_l + \chi_p^2 + \chi_q^2) \chi_g^2 + \chi_p^2 ((2 \tau^2 + 2) \chi_l^2 + 4 (\tau^2 + 1) \chi_q \chi_l + 4 (\tau^2 + 1) \chi_q^2)) \chi_f^2 + \\
& 2 \chi_g \chi_p^2 (-8 \chi_q^2 \tau^2 - 8 \chi_l \chi_q \tau^2 + 2 \chi_l^2) \chi_f + \\
& \chi_g^2 \chi_p^2 ((2 \tau^2 + 2) \chi_l^2 + 4 (\tau^2 + 1) \chi_q \chi_l + 4 (\tau^2 + 1) \chi_q^2))) m^4 \Big)
\end{aligned}$$

Check

No terms linear in a, as expected

```
In[74]:= Collect[Coefficient[Expr4, Pair[Momentum[a, D], Momentum[p, D]]], e]
```

```
Out[74]= 0
```

Finally, in this step, recall that

$$l = p + s k - q$$

$$l' = p + t k - q$$

Namely,

$$l^2 = 2 m^2 + 2 s [(kp) - (kq)] - 2 (pq)$$

$$(l')^2 = 2 m^2 + 2 t [(kp) - (kq)] - 2 (pq)$$

so we need to substitute (pq) here as well

$$\frac{1}{(l^2 + i\epsilon)(l'^2 + i\epsilon)} \text{ enters Coeff, therefore we rewrite it}$$

Note that we still did not pass to

new variables in the integral measure ,

we will introduce the Jacobian determinant in the following steps

```
In[75]:= Diffs4 = Diffs3;
```

```
Coeff4 = Simplify[
```

```
  Coeff3 /. {l1v2 -> 2 m^2 + 2 s (kp - kq) - 2 pq, l2v2 -> 2 m^2 + 2 t (kp - kq) - 2 pq} /. 
```

```
  ktochiSubs[p, q, f, g, l]]
```

```
Out[76]= (a^2 \xi^3 \chi p^2 \chi q^2)/(64 \pi^2 g0 L \phi m^2 p0 q0 \chi f (m^2 (\xi^3 \rho l^2 \chi l^2 + \xi (\tau l^2 \chi l^2 + (\chi p - \chi q)^2) + 2 s \chi p \chi q (\chi q - \chi p)) - i \epsilon \xi \chi p \chi q)
  (i \epsilon \xi \chi p \chi q + m^2 (\xi^3 \rho l^2 \chi l^2 + \xi (\tau l^2 \chi l^2 + (\chi p - \chi q)^2) + 2 t \chi p \chi q (\chi q - \chi p))))
```

Step 6: Recovering Airy functions

In this step, we substitute the explicit form of An,

Bn, Cn, Dn and pass to invariant variables in the process

Recall some definitions introduced at the beginning before Step 1

$$A_0(s, q, p) = \int \frac{d\phi}{2\pi} e^{is\phi} e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} = \frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \text{Ai}(y)$$

$$A_1(s, q, p) = -i \frac{dA_0}{ds} = \int \frac{d\phi}{2\pi} e^{is\phi} \phi e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} = \frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \left\{ \frac{\alpha}{8\beta} \text{Ai}(y) - i \frac{1}{(4\beta)^{1/3}} \text{Ai}'(y) \right\}$$

$$A_2(s, q, p) = (-i)^2 \frac{d^2 A_0}{ds^2} = \int \frac{d\phi}{2\pi} e^{is\phi} \phi^2 e^{-i\left(\frac{\alpha}{2}\phi^2 - \frac{4\beta}{3}\phi^3\right)} = \frac{\text{Exp}\left[i\left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2}\right)\right]}{(4\beta)^{1/3}} \left\{ \left[2\left(\frac{\alpha}{8\beta}\right)^2 - \frac{s}{4\beta} \right] \text{Ai}(y) - i \frac{\alpha}{(4\beta)^{4/3}} \text{Ai}'(y) \right\}$$

$$y = y(s, q, p) = \frac{1}{[4\beta(q, p)]^{1/3}} \left[s - \frac{\alpha^2(q, p)}{16\beta(q, p)} \right]$$

$$B_n = A_n^*$$

Note that each term in Expr4 is

proportional to multiplication of four symbols

$A_k B_l C_m D_n$,

where $k, l, m, n = 0, 1$,

2. Each represents a function A_n or B_n , in particular,

$$A_n \rightarrow A_n[s, q, p]$$

$$B_n \rightarrow B_n[t, q, p]$$

$$C_n \rightarrow A_n[\tilde{s} - s, f, -g]$$

$$D_n \rightarrow B_n[\tilde{s} - t, f, -g]$$

In effect, after writing functions A_n and B_n in Expr explicitly, we will obtain a linear combination of Ai and Ai' function tetrads, each having a different argument

We perform the substitution in the following order

1. Substitute A_n, B_n in place of A_n, B_n, C_n, D_n in Expr and Coeff

2. Functions A_n, B_n depend on momenta through $y[s, q, p]$,

$\alpha[q, p]$, and $\beta[q, p]$. Therefore,

we rewrite all these functions explicitly in terms of $\rho_{1,2}$, $\tau_{1,2}$, χ

3. Substitute y , α , β functions

4. Simplify the resulting expression and collect terms

In what follows, we first (Step 6.1) define

some symbols that contain a list of substitutions,
and then (Step 6.2) apply these substitutions to
Expr4 and Coeff4 obtained in the previous step

Step 6.1: Preparations: define substitutions $A_n \rightarrow A_i(y)$, $A_i'(y)$

We recover the explicit form of A_n and B_n by using pre -
defined substitutions

As mentioned (see the comment right after title ' Step 6 : ... '),
each A_n takes the form

$$A_n(s, q, p) = \frac{\text{Exp}\left[i \left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2} \right)\right]}{(4\beta)^{1/3}} * R_n[A_i(y), A_i'(y)],$$

where R_n is a linear combination
of the Airy functions (similarly for B_n)

As the prefactor here does not depend on n ,
it is convenient to keep the overall factor in Coeff. We substitute A_n ,
 B_n , C_n , D_n , appearing in Expr, by the corresponding expressions for R_n

SubsAnBack is the list of substitutions of symbols A_n , B_n , C_n ,
 D_n with A_n and B_n in the form of functions of their arguments

AnsubsNoFactor is introduced in definitions.nb

It contains the substitutions of A_n and B_n in a functional form with
corresponding explicit expressions for $R_n[A_i(y), A_i'(y)]$,
namely, linear combinations of Airy functions

Four factors $1/(4\beta)^{1/3}$ (one per each of A_n , B_n , C_n , and D_n)
are collected in a common factor named **AnFactor**

The sum of four exponentials, $i \left(\frac{\alpha s}{8\beta} - \frac{\alpha^3}{192\beta^2} \right)$, etc,

will be introduced in a separate variable in what follows

Note that Airy functions are expressed using Mathematica's `AiryAi` and `AiryAiPrime`, which obey usual Airy function properties

```
In[77]:= SubsAnBack = {A0 -> A0[s, q, p], A1 -> A1[s, q, p], A2 -> A2[s, q, p],
  B0 -> B0[t, q, p], B1 -> B1[t, q, p], B2 -> B2[t, q, p],
  C0 -> A0[-s + stilde, f, -g],
  C1 -> A1[-s + stilde, f, -g], C2 -> A2[-s + stilde, f, -g],
  D0 -> B0[stilde - t, f, -g], D1 -> B1[stilde - t, f, -g], D2 -> B2[stilde - t, f, -g]}
```

```
Out[77]= {A0 -> A0(s, q, p), A1 -> A1(s, q, p), A2 -> A2(s, q, p), B0 -> B0(t, q, p), B1 -> B1(t, q, p),
  B2 -> B2(t, q, p), C0 -> A0(stilde - s, f, -g), C1 -> A1(stilde - s, f, -g), C2 -> A2(stilde - s, f, -g),
  D0 -> B0(stilde - t, f, -g), D1 -> B1(stilde - t, f, -g), D2 -> B2(stilde - t, f, -g)}
```

```
In[78]:= AnSubsNoFactor (*print the list of substitutions, we use them later*)
AnFactor = 1 / (4 β[q, p] 4 β[f, -g] * 4 β[q, p] 4 β[f, -g]) ^ (1 / 3)
```

```
Out[78]= {A0(s_, q_, p_) -> Ai(y(s, q, p)), B0(s_, q_, p_) -> Ai(y(s, q, p)),
  A1(s_, q_, p_) -> \frac{\alpha(q, p) Ai(y(s, q, p))}{8 \beta(q, p)} - \frac{i Ai'(y(s, q, p))}{2^{2/3} \sqrt[3]{\beta(q, p)}}, B1(s_, q_, p_) -> \frac{\alpha(q, p) Ai(y(s, q, p))}{8 \beta(q, p)} + \frac{i Ai'(y(s, q, p))}{2^{2/3} \sqrt[3]{\beta(q, p)}},
  A2(s_, q_, p_) -> Ai(y(s, q, p)) \left( \frac{\alpha(q, p)^2}{64 \beta(q, p)^2} - \frac{y(s, q, p)}{2 \sqrt[3]{2} \beta(q, p)^{2/3}} \right) - \frac{i \alpha(q, p) Ai'(y(s, q, p))}{4 \times 2^{2/3} \beta(q, p)^{4/3}},
  B2(s_, q_, p_) -> Ai(y(s, q, p)) \left( \frac{\alpha(q, p)^2}{64 \beta(q, p)^2} - \frac{y(s, q, p)}{2 \sqrt[3]{2} \beta(q, p)^{2/3}} \right) + \frac{i \alpha(q, p) Ai'(y(s, q, p))}{4 \times 2^{2/3} \beta(q, p)^{4/3}} \}
```

```
Out[79]= \frac{1}{4 \times 2^{2/3} \sqrt[3]{\beta(f, -g)^2 \beta(q, p)^2}}
```

Explicit expressions for y , α , β in terms of invariant variables

Recall the definitions

$$y[s, q, p] = (4 \beta[q, p])^{2/3} \left[\frac{s}{4 \beta[q, p]} - \left(\frac{\alpha[q, p]}{8 \beta[q, p]} \right)^2 \right]$$

Similarly, we can write

$$y[t, q, p]$$

$$y[\tilde{s} - s, f, -g]$$

$$y[\tilde{s} - t, f, -g]$$

$$\alpha(q, p) = -e \left(\frac{ap}{kp} - \frac{aq}{kq} \right)$$

$$\beta(q, p) = \frac{e^2 a^2}{8} \left(\frac{1}{kp} - \frac{1}{kq} \right)$$

and analogously for $\alpha(f, -g)$, $\beta(f, -g)$

Let us rewrite $\tilde{s}$ (entering C_n and D_n), which we defined above as

$$\tilde{s} = \frac{\xi}{m^2 \chi_f} (pq + pg - qg - m^2)$$

in terms of the invariants. We already calculated scalar products (pq) , (pg) , (qg) , so we reuse these results

```
In[80]:= (*The definition of  $\tilde{s}$ *)
Simplify[( $\xi / m^2 / \chi_f$ ) (pq + pg - qg - m^2)];
Expand[% /. { $\chi_l \rightarrow \chi_p - \chi_q$ }];
Collect[%, {1 /  $\chi_f / \chi_g$ , 1 /  $\chi_f / \chi_p / \chi_q$ ,  $\rho_1$ ,  $\tau_1$ ,  $\rho_2$ ,  $\tau_2$ }, Simplify]
stildeDef = Simplify[% /. { $\chi_p - \chi_q \rightarrow \chi_l$ ,  $\chi_g - \chi_p + \chi_q \rightarrow -\chi_f$ }]
```

$$\begin{aligned} \text{Out[82]} = & -\frac{\xi^3 \rho_1^2 (\chi_p - \chi_q) (\chi_g - \chi_p + \chi_q)}{2 \chi_f \chi_p \chi_q} + \frac{\frac{1}{2} \xi^3 \rho_2^2 (\chi_p - \chi_q) + \frac{1}{2} \xi \tau_2^2 (\chi_p - \chi_q) + \frac{1}{2} \xi (\chi_p - \chi_q)}{\chi_f \chi_g} - \\ & \frac{\xi \tau_1^2 (\chi_p - \chi_q) (\chi_g - \chi_p + \chi_q)}{2 \chi_f \chi_p \chi_q} - \frac{\xi (\chi_p - \chi_q) (\chi_g - \chi_p + \chi_q)}{2 \chi_f \chi_p \chi_q} \\ \text{Out[83]} = & \frac{\xi \chi_l (\chi_f \chi_g (\xi^2 \rho_1^2 + \tau_1^2 + 1) + \chi_p \chi_q (\xi^2 \rho_2^2 + \tau_2^2 + 1))}{2 \chi_f \chi_g \chi_p \chi_q} \end{aligned}$$

Now, we find the arguments for the Airy functions

Commands **yArg**, **αdef**, and **βdef** are introduced in definitions.nb
They are used to write the Airy function arguments explicitly

```
In[84]:= (*example*)
yArg[s, q, p]
Expand[%]
```

$$\begin{aligned} \text{Out[84]} = & 2 \sqrt[3]{2} \beta(q, p)^{2/3} \left(\frac{s}{4 \beta(q, p)} - \frac{\alpha(q, p)^2}{64 \beta(q, p)^2} \right) \\ \text{Out[85]} = & \frac{s}{2^{2/3} \sqrt[3]{\beta(q, p)}} - \frac{\alpha(q, p)^2}{16 \times 2^{2/3} \beta(q, p)^{4/3}} \end{aligned}$$

Now, we rewrite α and β in terms of invariants by replacing scalar products (ap) , (af) , etc, which we identified earlier

```
In[86]:=  $\alpha$ def[q, p] /. {Pair[Momentum[a, D], Momentum[q, D]]  $\rightarrow$  av[ $\mu$ ]  $\times$  qv[ $\mu$ ]}
 $\alpha$ qpInv =
  Simplify[Contract[% /. {qv[ $\mu$ ]  $\rightarrow$  qpexpansion[ $\mu$ ]}] /. ktochiSubs[p, q, f, g, l] /.
    {av2  $\rightarrow$  -m^2  $\xi$ ^2 / e^2}] /. {xp - xq  $\rightarrow$  x $l$ , xq - xp  $\rightarrow$  -x $l$ }
 $\beta$ def[q, p] /. {Pair[Momentum[a, D], Momentum[q, D]]  $\rightarrow$  av[ $\mu$ ]  $\times$  qv[ $\mu$ ]}
 $\beta$ qpInv =
  Simplify[Contract[% /. {qv[ $\mu$ ]  $\rightarrow$  qpexpansion[ $\mu$ ]}] /. ktochiSubs[p, q, f, g, l] /.
    {av2  $\rightarrow$  -m^2  $\xi$ ^2 / e^2}] /. {xp - xq  $\rightarrow$  x $l$ , xq - xp  $\rightarrow$  -x $l$ }
```

$$\text{Out[86]} = -e \left(\frac{a \cdot p}{k \cdot p} - \frac{a^\mu q^\mu}{k \cdot q} \right)$$

$$\text{Out[87]} = -\frac{\xi^3 \rho_1 \chi^1}{\chi p \chi q}$$

$$\text{Out[88]} = \frac{1}{8} a^2 e^2 \left(\frac{1}{k \cdot p} - \frac{1}{k \cdot q} \right)$$

$$\text{Out[89]} = \frac{\xi^3 \chi^1}{8 \chi p \chi q}$$

```
In[90]:=  $\alpha$ def[f, -g] /. {Pair[Momentum[a, D], Momentum[g, D]]  $\rightarrow$  av[ $\mu$ ]  $\times$  gv[ $\mu$ ]}
 $\alpha$ fgInv =
  Simplify[Contract[% /. {gv[ $\mu$ ]  $\rightarrow$  gfexpansion[ $\mu$ ]}] /. ktochiSubs[p, q, f, g, l] /.
    {av2  $\rightarrow$  -m^2  $\xi$ ^2 / e^2}] /. {xf + xg  $\rightarrow$  x $l$ }
 $\beta$ def[f, -g] /. {Pair[Momentum[a, D], Momentum[g, D]]  $\rightarrow$  av[ $\mu$ ]  $\times$  gv[ $\mu$ ]}
 $\beta$ fgInv =
  Simplify[Contract[% /. {gv[ $\mu$ ]  $\rightarrow$  gfexpansion[ $\mu$ ]}] /. ktochiSubs[p, q, f, g, l] /.
    {av2  $\rightarrow$  -m^2  $\xi$ ^2 / e^2}] /. {xf + xg  $\rightarrow$  x $l$ }
```

$$\text{Out[90]} = -e \left(\frac{a^\mu g^\mu}{g \cdot k} - \frac{a \cdot f}{f \cdot k} \right)$$

$$\text{Out[91]} = -\frac{\xi^3 \rho_2 \chi^1}{\chi f \chi g}$$

$$\text{Out[92]} = \frac{1}{8} a^2 e^2 \left(-\frac{1}{f \cdot k} - \frac{1}{g \cdot k} \right)$$

$$\text{Out[93]} = \frac{\xi^3 \chi^1}{8 \chi f \chi g}$$

With this, we can write the Airy function arguments explicitly

Notations

ysqp == y (s, q, p)

ytqp == y (t, q, p)

In[94]:= `yArg[s, q, p] /. {α[q, p] → αqpInv, β[q, p] → βqpInv};`

$$ysqp = \frac{1}{\xi} \left(\frac{\chi^l}{2 \chi p \chi q} \right)^{-1/3} \text{Simplify} \left[\text{Expand}[\%] \left(\frac{\xi^3 \chi^l}{2 \chi p \chi q} \right)^{1/3} \right]$$

`yArg[t, q, p] /. {α[q, p] → αqpInv, β[q, p] → βqpInv};`

$$ytqp = \frac{1}{\xi} \left(\frac{\chi^l}{2 \chi p \chi q} \right)^{-1/3} \text{Simplify} \left[\text{Expand}[\%] \left(\frac{\xi^3 \chi^l}{2 \chi p \chi q} \right)^{1/3} \right]$$

Out[95]=
$$\frac{\sqrt[3]{2} \left(s - \frac{\xi^3 \rho^2 \chi^l}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi p \chi q}}}$$

Out[97]=
$$\frac{\sqrt[3]{2} \left(t - \frac{\xi^3 \rho^2 \chi^l}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi p \chi q}}}$$

Notations

`ysfg` == $y(s, f, -g)$ (used only in an intermediate step)

`ystsf` == $y(\tilde{s} - s, f, -g)$

`ytfg` == $y(t, f, -g)$ (used only in an intermediate step)

`ysttf` == $y(\tilde{s} - t, f, -g)$

In[98]:= `yArg[s, f, -g] /. {α[f, -g] → αfgInv, β[f, -g] → βfgInv};`

$$ysfg = \frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \text{Expand} \left[\% \left(\frac{\xi^3 \chi^l}{2 \chi^f \chi^g} \right)^{1/3} \right]$$

`ystsfg =`

$$\frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \text{Collect} \left[\text{Expand} \left[\left(ysfg / \left(\frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \right) \right) /. \{s \rightarrow \text{stildeDef} - s\} \right], \{ \chi^p, \chi^q, \chi^f, \chi^g \}, \text{Simplify} \right]$$

`yArg[t, f, -g] /. {α[f, -g] → αfgInv, β[f, -g] → βfgInv};`

$$ytfg = \frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \text{Expand} \left[\% \left(\frac{\xi^3 \chi^l}{2 \chi^f \chi^g} \right)^{1/3} \right]$$

`ysttfg =`

$$\frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \text{Collect} \left[\text{Expand} \left[\left(ytfg / \left(\frac{1}{\xi} \left(\frac{\chi^l}{2 \chi^f \chi^g} \right)^{-1/3} \right) \right) /. \{t \rightarrow \text{stildeDef} - t\} \right], \{ \chi^p, \chi^q, \chi^f, \chi^g \}, \text{Simplify} \right]$$

Out[99]=
$$\frac{\sqrt[3]{2} \left(s - \frac{\xi^3 \rho^2 \chi^l}{2 \chi^f \chi^g} \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi^f \chi^g}}}$$

Out[100]=
$$\frac{\sqrt[3]{2} \left(\frac{\xi \chi^l (\xi^2 \rho^2 + \tau^2 + 1)}{2 \chi^p \chi^q} + \frac{\xi (\tau^2 + 1) \chi^l}{2 \chi^f \chi^g} - s \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi^f \chi^g}}}$$

Out[102]=
$$\frac{\sqrt[3]{2} \left(t - \frac{\xi^3 \rho^2 \chi^l}{2 \chi^f \chi^g} \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi^f \chi^g}}}$$

Out[103]=
$$\frac{\sqrt[3]{2} \left(\frac{\xi \chi^l (\xi^2 \rho^2 + \tau^2 + 1)}{2 \chi^p \chi^q} + \frac{\xi (\tau^2 + 1) \chi^l}{2 \chi^f \chi^g} - t \right)}{\xi \sqrt[3]{\frac{\chi^l}{\chi^f \chi^g}}}$$

The resulting list for α ,
 β and y variables : **Subsαβ**, **SubsyArgExplicit**

SubsyArgsInAi is used to make the output expressions more compact
 by replacing the Airy function arguments with shorthands

`ys1 == y[s, q, p]`

`yt1 == y[t, q, p]`

`ys2 == y[-s + stilde, f, -g]`

`yt2 == y[stilde - t, f, -g]`

In[104]:= **Subs** $\alpha\beta = \{\alpha[q, p] \rightarrow \alpha qpInv, \beta[q, p] \rightarrow \beta qpInv, \alpha[f, -g] \rightarrow \alpha fgInv, \beta[f, -g] \rightarrow \beta fgInv\}$

SubsyArgExplicit = $\{y[s, q, p] \rightarrow ysqp, y[t, q, p] \rightarrow ytpq,$

$y[-s + stilde, f, -g] \rightarrow ystsfg, y[stilde - t, f, -g] \rightarrow ysttfg\}$

SubsyArgsInAi =

$\{AiryAi[y[s, q, p]] \rightarrow AiryAi[ys1], AiryAi[y[t, q, p]] \rightarrow AiryAi[yt1],$

$AiryAi[y[-s + stilde, f, -g]] \rightarrow AiryAi[ys2],$

$AiryAi[y[stilde - t, f, -g]] \rightarrow AiryAi[yt2],$

$AiryAiPrime[y[s, q, p]] \rightarrow AiryAiPrime[ys1],$

$AiryAiPrime[y[t, q, p]] \rightarrow AiryAiPrime[yt1],$

$AiryAiPrime[y[-s + stilde, f, -g]] \rightarrow AiryAiPrime[ys2],$

$AiryAiPrime[y[stilde - t, f, -g]] \rightarrow AiryAiPrime[yt2]\}$

$$\text{Out[104]} = \left\{ \alpha(q, p) \rightarrow -\frac{\xi^3 \rho^1 \chi^1}{\chi p \chi q}, \beta(q, p) \rightarrow \frac{\xi^3 \chi^1}{8 \chi p \chi q}, \alpha(f, -g) \rightarrow -\frac{\xi^3 \rho^2 \chi^1}{\chi f \chi g}, \beta(f, -g) \rightarrow \frac{\xi^3 \chi^1}{8 \chi f \chi g} \right\}$$

$$\text{Out[105]} = \left\{ y(s, q, p) \rightarrow \frac{\sqrt[3]{2} \left(s - \frac{\xi^3 \rho^1 \chi^1}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi p \chi q}}}, y(t, q, p) \rightarrow \frac{\sqrt[3]{2} \left(t - \frac{\xi^3 \rho^1 \chi^1}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi p \chi q}}}, \right.$$

$$\left. y(stilde - s, f, -g) \rightarrow \frac{\sqrt[3]{2} \left(\frac{\xi \chi^1 (\xi^2 \rho^1 \chi^1 + \tau^1 \chi^1)}{2 \chi p \chi q} + \frac{\xi (\tau^2 + 1) \chi^1}{2 \chi f \chi g} - s \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi f \chi g}}}, y(stilde - t, f, -g) \rightarrow \frac{\sqrt[3]{2} \left(\frac{\xi \chi^1 (\xi^2 \rho^1 \chi^1 + \tau^1 \chi^1)}{2 \chi p \chi q} + \frac{\xi (\tau^2 + 1) \chi^1}{2 \chi f \chi g} - t \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi f \chi g}}} \right\}$$

Out[106]= **Ai**($y(s, q, p)$) $\rightarrow$ **Ai**(ys1), **Ai**($y(t, q, p)$) $\rightarrow$ **Ai**(yt1), **Ai**($y(stilde - s, f, -g)$) $\rightarrow$ **Ai**(ys2),

Ai($y(stilde - t, f, -g)$) $\rightarrow$ **Ai**(yt2), **Ai'**($y(s, q, p)$) $\rightarrow$ **Ai'**(ys1), **Ai'**($y(t, q, p)$) $\rightarrow$ **Ai'**(yt1),

Ai'($y(stilde - s, f, -g)$) $\rightarrow$ **Ai'**(ys2), **Ai'**($y(stilde - t, f, -g)$) $\rightarrow$ **Ai'**(yt2)}

Some default assumptions used in command **Simplify** below

In[107]:= **assump** $\xi\chi = \{\xi > 0, \chi p > 0, \chi q > 0, \chi f > 0, \chi g > 0\}$

assumpAll = $\{u > 0, v > 4, \chi p > 0, \chi q > 0, \chi g > 0, \chi f > 0, \chi f > 0, \xi > 0\}$

\$Assumptions = **assumpAll**;

Out[107]= $\{\xi > 0, \chi p > 0, \chi q > 0, \chi f > 0, \chi g > 0\}$

Out[108]= $\{u > 0, v > 4, \chi p > 0, \chi q > 0, \chi g > 0, \chi f > 0, \chi f > 0, \xi > 0\}$

```
In[110]:= (*a small test, all the substitutions defined above are applied one by one*)
A2 B2 C2 D2 /. SubsAnBack /. AnsubsNoFactor /. SubsyArgsInAi
% /. Subsaβ /. SubsyArgExplicit
```

$$\begin{aligned} \text{Out[110]} = & \left(\text{Ai}(ys1) \left(\frac{\alpha(q, p)^2}{64 \beta(q, p)^2} - \frac{y(s, q, p)}{2 \sqrt[3]{2} \beta(q, p)^{2/3}} \right) - \frac{i \text{Ai}'(ys1) \alpha(q, p)}{4 \times 2^{2/3} \beta(q, p)^{4/3}} \right) \\ & \left(\text{Ai}(yt1) \left(\frac{\alpha(q, p)^2}{64 \beta(q, p)^2} - \frac{y(t, q, p)}{2 \sqrt[3]{2} \beta(q, p)^{2/3}} \right) + \frac{i \text{Ai}'(yt1) \alpha(q, p)}{4 \times 2^{2/3} \beta(q, p)^{4/3}} \right) \\ & \left(\text{Ai}(ys2) \left(\frac{\alpha(f, -g)^2}{64 \beta(f, -g)^2} - \frac{y(\text{stilde} - s, f, -g)}{2 \sqrt[3]{2} \beta(f, -g)^{2/3}} \right) - \frac{i \text{Ai}'(ys2) \alpha(f, -g)}{4 \times 2^{2/3} \beta(f, -g)^{4/3}} \right) \\ & \left(\text{Ai}(yt2) \left(\frac{\alpha(f, -g)^2}{64 \beta(f, -g)^2} - \frac{y(\text{stilde} - t, f, -g)}{2 \sqrt[3]{2} \beta(f, -g)^{2/3}} \right) + \frac{i \text{Ai}'(yt2) \alpha(f, -g)}{4 \times 2^{2/3} \beta(f, -g)^{4/3}} \right) \\ \text{Out[111]} = & \left(\text{Ai}(ys1) \left(\rho^2 - \frac{2 \left(s - \frac{\xi^3 \rho^2 \chi^1}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi p \chi q} \left(\frac{\xi^3 \chi^1}{\chi p \chi q} \right)^{2/3}}} \right) + \frac{2 i \sqrt[3]{2} \xi^3 \rho^1 \chi^1 \text{Ai}'(ys1)}{\chi p \chi q \left(\frac{\xi^3 \chi^1}{\chi p \chi q} \right)^{4/3}} \right) \\ & \left(\text{Ai}(yt1) \left(\rho^2 - \frac{2 \left(t - \frac{\xi^3 \rho^2 \chi^1}{2 \chi p \chi q} \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi p \chi q} \left(\frac{\xi^3 \chi^1}{\chi p \chi q} \right)^{2/3}}} \right) - \frac{2 i \sqrt[3]{2} \xi^3 \rho^1 \chi^1 \text{Ai}'(yt1)}{\chi p \chi q \left(\frac{\xi^3 \chi^1}{\chi p \chi q} \right)^{4/3}} \right) \\ & \left(\text{Ai}(ys2) \left(\rho^2 - \frac{2 \left(\frac{\xi \chi^1 (\xi^2 \rho^2 + \tau^2 + 1)}{2 \chi p \chi q} + \frac{\xi (\tau^2 + 1) \chi^1}{2 \chi f \chi g} - s \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi f \chi g} \left(\frac{\xi^3 \chi^1}{\chi f \chi g} \right)^{2/3}}} \right) + \frac{2 i \sqrt[3]{2} \xi^3 \rho^2 \chi^1 \text{Ai}'(ys2)}{\chi f \chi g \left(\frac{\xi^3 \chi^1}{\chi f \chi g} \right)^{4/3}} \right) \\ & \left(\text{Ai}(yt2) \left(\rho^2 - \frac{2 \left(\frac{\xi \chi^1 (\xi^2 \rho^2 + \tau^2 + 1)}{2 \chi p \chi q} + \frac{\xi (\tau^2 + 1) \chi^1}{2 \chi f \chi g} - t \right)}{\xi \sqrt[3]{\frac{\chi^1}{\chi f \chi g} \left(\frac{\xi^3 \chi^1}{\chi f \chi g} \right)^{2/3}}} \right) - \frac{2 i \sqrt[3]{2} \xi^3 \rho^2 \chi^1 \text{Ai}'(yt2)}{\chi f \chi g \left(\frac{\xi^3 \chi^1}{\chi f \chi g} \right)^{4/3}} \right) \end{aligned}$$

Step 6.2: Substitutions and simplification of resulting expressions

Now we use the introduced substitutions to express An, Bn, Cn, Dn explicitly through invariant variables

First, let us calculate

Phase5 –

the sum of all exponentials (with i omitted) of the form $\left(\frac{\alpha s}{8 \beta} - \frac{\alpha^3}{192 \beta^2} \right)$

```
In[112]:= Simplify[(AnPhase[s, q, p] +
  AnPhase[st - s, f, -g] - AnPhase[t, q, p] - AnPhase[st - t, f, -g])]
Phase5 = Simplify[% /. {st → stildeDef} /. Subsaβ, Assumptions → assumptξχ]
```

$$\text{Out[112]} = \frac{(s-t)(\beta(f, -g)\alpha(q, p) - \alpha(f, -g)\beta(q, p))}{8\beta(f, -g)\beta(q, p)}$$

$$\text{Out[113]} = -((\rho_1 - \rho_2)(s - t))$$

Performing the actual substitutions as described, to obtain Expr5

Note that AnFactor is introduced in variable Expr and not in Coeff

Expr5 is lengthy and is not printed for this reason

```
In[114]:= Diffs5 = Diffs4;
Coeff5 = Coeff4;
Expr4 * AnFactor;
Expand[%] /. SubsAnBack /. AnsubsNoFactor /. SubsyArgsInAi /. Subsaβ /.
  SubsyArgExplicit;
Expr5 = Expand[%];
```

In order to simplify the result,
let us now pass to new variables in the integral measure

$$u = \frac{\chi_l}{\chi_q}, \quad 0 < u < \infty$$

$$v^2 = \frac{\chi_l^2}{\chi_g \chi_f}, \quad 4 < v < \infty$$

$$\chi_g = \frac{\chi_l}{2} \left(1 + \sigma \sqrt{1 - \frac{4}{v}} \right), \quad \chi_f = \frac{\chi_l}{2} \left(1 - \sigma \sqrt{1 - \frac{4}{v}} \right),$$

$\sigma = \pm 1$ are two branches, arising in this variable change;

The final result is given by summation over σ

In effect,

combined with variable changes introduced above $(\rho_{1,2}, \tau_{1,2})$,

the overall integral measure transforms as

$$d^3g d^3q \rightarrow |J_1^{-1}| |J_2^{-1}| d\rho_1 d\tau_1 du d\rho_2 d\tau_2 dv,$$

with Jacobian determinants given by

$$|J_1^{-1}| = q_0 m^2 \xi \frac{u^2}{(1+u)^3}$$

$$|J_2^{-1}| = g_0 m^2 \xi \frac{\chi_l}{\chi_g} \frac{1}{v \sqrt{v(v-4)}}$$

$$\text{In[119]:= } \text{jac1} = q_0 m^2 \xi \frac{u^2}{(1+u)^3}$$

$$\text{jac2} = g_0 m^2 \xi \frac{\chi^l}{\chi g} \frac{1}{v \sqrt{v-4}}$$

$$\text{Out[119]= } \frac{m^2 \xi q_0 u^2}{(u+1)^3}$$

$$\text{Out[120]= } \frac{g_0 m^2 \xi \chi^l}{v \sqrt{v-4} v \chi g}$$

$$\text{In[121]:= } \text{subsu} = \{\chi q \rightarrow \chi p / (1+u), \chi^l \rightarrow \chi p u / (1+u)\}$$

$$\text{subsv}\sigma = \{\chi g \rightarrow (1/2) \chi^l (1 + \sigma \sqrt{1-4/v}), \chi^f \rightarrow (1/2) \chi^l (1 - \sigma \sqrt{1-4/v})\}$$

$$\text{Out[121]= } \left\{ \chi q \rightarrow \frac{\chi p}{u+1}, \chi^l \rightarrow \frac{u \chi p}{u+1} \right\}$$

$$\text{Out[122]= } \left\{ \chi g \rightarrow \frac{1}{2} \chi^l \left(\sigma \sqrt{1 - \frac{4}{v}} + 1 \right), \chi^f \rightarrow \frac{1}{2} \chi^l \left(1 - \sigma \sqrt{1 - \frac{4}{v}} \right) \right\}$$

$$\text{In[123]:= } (*\text{It is straightforward to see that the following result depends only on } \sigma^2, \text{ so we can just set } \sigma \rightarrow 1*)$$

$$\text{Coeff5} * \text{jac1} * \text{jac2};$$

$$\% /. \{\sigma \rightarrow 1\};$$

$$\text{Coeff6} = \text{Simplify}[\% /. \text{subsv}\sigma /. \text{subsu}]$$

$$\text{Diffs6} = \text{Diffs5} /. \{d3q \rightarrow d\rho_1 d\tau_1 du, d3g \rightarrow d\rho_2 d\tau_2 dv\}$$

$$\text{Phase6} = \text{Phase5}$$

$$\text{Out[125]= } (a^2 m^2 \xi^5 u) / \left(16 \pi^2 L \phi p_0 v \sqrt{v-4} v \chi p \left(\sigma \sqrt{\frac{v-4}{v}} - 1 \right) \left(\sigma \sqrt{\frac{v-4}{v}} + 1 \right) \right. \\ \left. (i \epsilon \xi (u+1) - m^2 u (\xi u (\xi^2 \rho_1^2 + \tau_1^2 + 1) - 2 s \chi p)) (i \epsilon \xi (u+1) + m^2 u (\xi u (\xi^2 \rho_1^2 + \tau_1^2 + 1) - 2 t \chi p)) \right)$$

$$\text{Out[126]= } d\rho_1 d\rho_2 ds dt d\tau_1 d\tau_2 du dv$$

$$\text{Out[127]= } -((\rho_1 - \rho_2)(s - t))$$

Summation over two σ branches

$$\text{In[128]:= } \text{Expr6}\sigma = \text{Expr5} /. \text{subsv}\sigma /. \text{subsu};$$

$$\text{Expr6} = (\text{Expr6}\sigma /. \{\sigma \rightarrow 1\}) + (\text{Expr6}\sigma /. \{\sigma \rightarrow -1\});$$

In the next step, we collect terms in Expr -- all possible combinations of four Airy functions and their derivatives -- and simplify them

```

In[130]:= Collect4Airy = {
    AiryAi[ys1] AiryAi[ys2] AiryAi[yt1] AiryAi[yt2],
    AiryAi'[ys1] AiryAi[ys2] AiryAi[yt1] AiryAi[yt2],
    AiryAi[ys1] AiryAi'[ys2] AiryAi[yt1] AiryAi[yt2],
    AiryAi[ys1] AiryAi[ys2] AiryAi'[yt1] AiryAi[yt2],
    AiryAi[ys1] AiryAi[ys2] AiryAi[yt1] AiryAi'[yt2],

    AiryAi'[ys1] AiryAi'[ys2] AiryAi[yt1] AiryAi[yt2],
    AiryAi'[ys1] AiryAi[ys2] AiryAi'[yt1] AiryAi[yt2],
    AiryAi'[ys1] AiryAi[ys2] AiryAi[yt1] AiryAi'[yt2],
    AiryAi[ys1] AiryAi'[ys2] AiryAi'[yt1] AiryAi[yt2],
    AiryAi[ys1] AiryAi'[ys2] AiryAi[yt1] AiryAi'[yt2],
    AiryAi[ys1] AiryAi[ys2] AiryAi'[yt1] AiryAi'[yt2],

    AiryAi'[ys1] AiryAi'[ys2] AiryAi'[yt1] AiryAi[yt2],
    AiryAi'[ys1] AiryAi'[ys2] AiryAi[yt1] AiryAi'[yt2],
    AiryAi'[ys1] AiryAi[ys2] AiryAi'[yt1] AiryAi'[yt2],
    AiryAi[ys1] AiryAi'[ys2] AiryAi'[yt1] AiryAi'[yt2],
    AiryAi'[ys1] AiryAi'[ys2] AiryAi'[yt1] AiryAi'[yt2]
}

Out[130]:= {Ai(ys1) Ai(ys2) Ai(yt1) Ai(yt2), Ai'(ys1) Ai(ys2) Ai(yt1) Ai(yt2),
    Ai(ys1) Ai'(ys2) Ai(yt1) Ai(yt2), Ai(ys1) Ai(ys2) Ai'(yt1) Ai(yt2), Ai(ys1) Ai(ys2) Ai(yt1) Ai'(yt2),
    Ai'(ys1) Ai'(ys2) Ai(yt1) Ai(yt2), Ai'(ys1) Ai(ys2) Ai'(yt1) Ai(yt2), Ai'(ys1) Ai(ys2) Ai(yt1) Ai'(yt2),
    Ai(ys1) Ai'(ys2) Ai'(yt1) Ai(yt2), Ai(ys1) Ai'(ys2) Ai(yt1) Ai'(yt2), Ai(ys1) Ai(ys2) Ai'(yt1) Ai'(yt2),
    Ai'(ys1) Ai'(ys2) Ai'(yt1) Ai(yt2), Ai'(ys1) Ai'(ys2) Ai(yt1) Ai'(yt2), Ai'(ys1) Ai(ys2) Ai'(yt1) Ai'(yt2),
    Ai(ys1) Ai'(ys2) Ai'(yt1) Ai'(yt2), Ai'(ys1) Ai'(ys2) Ai'(yt1) Ai'(yt2)}

```

In[131]:= Simplify[

Expr6 /. {D → 4},

Assumptions → {u > 0, v > 4, xp > 0, xq > 0, xg > 0, xf > 0, χf > 0, ξ > 0}];

Expr7 = PowerExpand[Collect[Expand[%], Collect4Airy, Simplify]];

Diffs7 = Diffs6

Coeff7 = Coeff6

Phase7 = Phase6

Expr7

Out[133]:= dp1 dp2 ds dt dτ1 dτ2 du dv

$$\text{Out[134]} = (\alpha^2 m^2 \xi^5 u) \left/ \left(16 \pi^2 L \phi p_0 v \sqrt{(v-4)v} \chi p \left(\sigma \sqrt{\frac{v-4}{v}} - 1 \right) \left(\sigma \sqrt{\frac{v-4}{v}} + 1 \right) \right. \right. \right. \\ \left. \left. \left. (\text{i} \epsilon \xi (u+1) - m^2 u (\xi u (\xi^2 \rho_1^2 + \tau_1^2 + 1) - 2 s \chi p)) (\text{i} \epsilon \xi (u+1) + m^2 u (\xi u (\xi^2 \rho_1^2 + \tau_1^2 + 1) - 2 t \chi p)) \right) \right)$$

Out[135]:= -((ρ1 - ρ2) (s - t))

$$\text{Out[136]} = \frac{1}{(u+1)^{5/3} v^{5/3} \xi^6} 32 \sqrt[3]{2} \chi p^{4/3} \\ \left(-8 v \tau_1^2 \tau_2^2 \xi^2 + 2 v^2 \tau_1^2 (\tau_2^2 + 1) \xi^2 + u^2 ((\tau_1^2 + 1) (\tau_2^2 + 1) v^2 - 2 (\tau_1^2 + 1) \tau_2^2 v + 8 (\xi^2 \rho_1^2 + \tau_1^2 + 1)^2) \xi^2 - \right. \\ \left. 2 u (4 v \xi \tau_2^2 \tau_1^2 - v^2 \xi (\tau_2^2 + 1) \tau_1^2 + 8 (s+t) (\xi^2 \rho_1^2 + \tau_1^2 + 1) \chi p) \xi + 32 s t \chi p^2 \right) \text{Ai}(ys_1) \text{Ai}(ys_2) \text{Ai}(yt_1) \\ \text{Ai}(yt_2) m^4 + \frac{1}{u^{2/3} (u+1)^{8/3} v^{5/3} \xi^4} 64 \tau_1 \tau_2 (u^{5/3} (u+1)^{2/3} (u+2) \chi p^{2/3} v^{5/3} + 2 u^{2/3} (u+1)^{2/3} \chi p^{2/3} v^{5/3} - \\ 2 ((u+1)^{2/3} v^{2/3} \chi p^{2/3} u^{8/3} + 4 (u+1)^{2/3} v^{2/3} \chi p^{2/3} u^{5/3} + 4 (u+1)^{2/3} v^{2/3} \chi p^{2/3} u^{2/3})) \chi p^{4/3} \text{Ai}(yt_1) \text{Ai}(yt_2) \\ \text{Ai}'(ys_1) \text{Ai}'(ys_2) m^4 + \frac{64 ((u^2 + 2 u + 2) v (\tau_2^2 + 1) - 2 u^2 \tau_2^2) \chi p^2 \text{Ai}(ys_2) \text{Ai}(yt_2) \text{Ai}'(ys_1) \text{Ai}'(yt_1) m^4}{u^{2/3} (u+1)^{5/3} v^{2/3} \xi^4} + \\ \frac{64 ((v-2) u^2 + 2 v u + 2 v) \tau_1 \tau_2 \chi p^2 \text{Ai}(ys_1) \text{Ai}(yt_2) \text{Ai}'(ys_2) \text{Ai}'(yt_1) m^4}{(u+1)^2 v \xi^4} + \\ \frac{64 ((v-2) u^2 + 2 v u + 2 v) \tau_1 \tau_2 \chi p^2 \text{Ai}(ys_2) \text{Ai}(yt_1) \text{Ai}'(ys_1) \text{Ai}'(yt_2) m^4}{(u+1)^2 v \xi^4} - \\ \frac{1}{u^2 (u+1)^{8/3} v^{8/3} \xi^4} 64 (-\sqrt[3]{u+1} v^{7/3} \chi p^{2/3} u^{14/3} + 2 \sqrt[3]{u+1} v^{4/3} \chi p^{2/3} u^{14/3} + \tau_1^2 \\ (-\sqrt[3]{u+1} v^{7/3} \chi p^{2/3} u^{14/3} + 2 \sqrt[3]{u+1} v^{4/3} \chi p^{2/3} u^{14/3} - 2 \sqrt[3]{u+1} v^{7/3} \chi p^{2/3} u^{11/3} - 2 \sqrt[3]{u+1} v^{7/3} \chi p^{2/3} u^{8/3})) \\ \chi p^{4/3} \text{Ai}(ys_1) \text{Ai}(yt_1) \text{Ai}'(ys_2) \text{Ai}'(yt_2) m^4 + \\ \frac{64 ((v-2) u^2 + 2 (v-4) u + 2 (v-4)) \tau_1 \tau_2 \chi p^2 \text{Ai}(ys_1) \text{Ai}(ys_2) \text{Ai}'(yt_1) \text{Ai}'(yt_2) m^4}{(u+1)^2 v \xi^4} - \\ \frac{1}{u^{2/3} (u+1)^{8/3} v^{8/3} \xi^4} 64 \times 2^{2/3} \chi p^{4/3} \\ (-\sqrt[3]{u+1} v^{7/3} \chi p^{4/3} u^{8/3} + 2 \sqrt[3]{u+1} v^{4/3} \chi p^{4/3} u^{8/3} - 2 \sqrt[3]{u+1} v^{7/3} \chi p^{4/3} u^{5/3} + 8 \sqrt[3]{u+1} v^{4/3} \chi p^{4/3} u^{5/3} - \\ 2 \sqrt[3]{u+1} v^{7/3} \chi p^{4/3} u^{2/3} + 8 \sqrt[3]{u+1} v^{4/3} \chi p^{4/3} u^{2/3}) \text{Ai}'(ys_1) \text{Ai}'(ys_2) \text{Ai}'(yt_1) \text{Ai}'(yt_2) m^4$$

At this stage, we observe that Expr7 contains a number of terms that are proportional to ρ_1^n , $n = 1, 2, 3, 4$. In what follows,

we will carry out integrals over $\rho_{1,2}$. Apparently, these terms can be removed by changing variables s and t , namely, by shifting them. This simplifies the integration greatly.

As another observation, we do not get any such terms proportional to powers of ρ_2

```
In[137]:= Coefficient[Expr7, {ρ1, ρ1^2, ρ1^3, ρ1^4}];
Coefficient[%, AiryAi[ys1] AiryAi[ys2] AiryAi[yt1] AiryAi[yt2]]
Out[138]= {0, - $\frac{512 \sqrt{2} m^4 s u \chi p^{7/3}}{\xi^3 (u+1)^{5/3} v^{5/3}} - \frac{512 \sqrt{2} m^4 t u \chi p^{7/3}}{\xi^3 (u+1)^{5/3} v^{5/3}} + \frac{512 \sqrt{2} m^4 \tau l^2 u^2 \chi p^{4/3}}{\xi^2 (u+1)^{5/3} v^{5/3}} + \frac{512 \sqrt{2} m^4 u^2 \chi p^{4/3}}{\xi^2 (u+1)^{5/3} v^{5/3}},$ 
0,  $\frac{256 \sqrt{2} m^4 u^2 \chi p^{4/3}}{(u+1)^{5/3} v^{5/3}}$ }
```

```
In[139]:= Coefficient[Expr7, {ρ2, ρ2^2, ρ2^3, ρ2^4}]
Out[139]= {0, 0, 0, 0}
```

Step 7: Shift of s and t and intermediate results

We shift s and t variables to simplify Coeff7 and remove the explicit dependence on τi and ρi in the denominator

Note that we use the property

$$i \in * X = i \in \text{for } X > 0$$

```
In[140]:= shift = {s → s +  $\frac{u \xi}{2 \chi p} (1 + \tau l^2 + \xi^2 \rho l^2)$ , t → t +  $\frac{u \xi}{2 \chi p} (1 + \tau l^2 + \xi^2 \rho l^2)$ };
factor8 = (8 m^2 u χ p / ξ^3 / (1 + u))^2;
Coeff8 = Simplify[Coeff7 * factor8 /. {ie → 0} /. shift] /.
{1 / s → 1 / (s + iε), 1 / t → 1 / (t - iε)}
Diffs8 = Diffs7;
Phase8 = Phase7;
```

$$\text{Out[142]} = -\frac{\alpha^2 m^2 u}{\pi^2 L \phi \xi p_0 (u+1)^2 \sqrt{(v-4)v} \chi p (i\epsilon + s) (t - i\epsilon) ((\sigma^2 - 1)v - 4\sigma^2)}$$

In[145]:= **Expand**[**Simplify**[**Expr7** / **factor8** /. **shift**] /. { $\left(\sqrt{\frac{v-4}{v}} - 1 \right) \left(\sqrt{\frac{v-4}{v}} + 1 \right) \rightarrow -\frac{4}{v} \}$];

Simplify[**PowerExpand**[%],

Assumptions → {**u** > 0, **v** > 4, **χp** > 0, **χq** > 0, **χg** > 0, **χf** > 0, **χf** > 0, **ξ** > 0}];

Expr8 = **Collect**[**Expand**[**PowerExpand**[%]], **Collect4Airy**, **Simplify**]

$$\begin{aligned} \text{Out[147]} = & \frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \text{Ai}(ys1) \text{Ai}(ys2) \text{Ai}(yt1) \text{Ai}(yt2) (32 s t \chi p^2 + \\ & \xi^2 (\tau_1^2 + 1) u^2 v (-2 \tau_2^2 + \tau_2^2 v + v) + 2 \xi^2 \tau_1^2 u v (-4 \tau_2^2 + \tau_2^2 v + v) + 2 \xi^2 \tau_1^2 (\tau_2^2 + 1) v^2 - 8 \xi^2 \tau_1^2 \tau_2^2 v) + \\ & \frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v-2) + 2 u (v-4) + 2 (v-4)) \text{Ai}'(ys1) \text{Ai}'(ys2) \text{Ai}'(yt1) \text{Ai}'(yt2)}{u^2 \sqrt[3]{(u+1) v^4}} + \\ & \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u (v-4) + 2 (v-4)) \text{Ai}'(ys1) \text{Ai}'(ys2) \text{Ai}(yt1) \text{Ai}(yt2)}{u^2 v} + \\ & \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u v + 2 v) \text{Ai}(ys1) \text{Ai}'(ys2) \text{Ai}'(yt1) \text{Ai}(yt2)}{u^2 v} + \\ & \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u v + 2 v) \text{Ai}'(ys1) \text{Ai}(ys2) \text{Ai}(yt1) \text{Ai}'(yt2)}{u^2 v} + \\ & \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u (v-4) + 2 (v-4)) \text{Ai}(ys1) \text{Ai}(ys2) \text{Ai}'(yt1) \text{Ai}'(yt2)}{u^2 v} + \\ & \frac{\xi^2 \sqrt[3]{u+1} \text{Ai}'(ys1) \text{Ai}(ys2) \text{Ai}'(yt1) \text{Ai}(yt2) ((\tau_2^2 + 1) (u^2 + 2 u + 2) v - 2 \tau_2^2 u^2)}{u^{8/3} v^{2/3}} + \\ & \frac{\xi^2 \text{Ai}(ys1) \text{Ai}'(ys2) \text{Ai}(yt1) \text{Ai}'(yt2) ((\tau_1^2 + 1) u^2 (v-2) + 2 \tau_1^2 u v + 2 \tau_1^2 v)}{\sqrt[3]{u^4 (u+1) v^4}} \end{aligned}$$

In[148]:= **(*must be all zeros*)**

Coefficient[**Expr8**, {**ρ1**, **ρ1**^2, **ρ1**^3, **ρ1**^4}]

Out[148]:= {0, 0, 0, 0}

Let us collect the intermediate result for Expr, expressed term by term (collected for different Airy function combinations)

In[149]:= **Coefficient**[**Expr8**, **AiryAi**[**ys1**] **AiryAi**[**ys2**] **AiryAi**[**yt1**] **AiryAi**[**yt2**]] /.
{**τ1** → **τ₁**, **τ2** → **τ₂**}

$$\text{Out[149]} = \frac{\sqrt[3]{u+1} (32 s t \chi p^2 + \xi^2 (\tau_1^2 + 1) u^2 v (-2 \tau_2^2 + \tau_2^2 v + v) + 2 \xi^2 \tau_1^2 u v (-4 \tau_2^2 + \tau_2^2 v + v) + 2 \xi^2 \tau_1^2 (\tau_2^2 + 1) v^2 - 8 \xi^2 \tau_1^2 \tau_2^2 v)}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}}$$

In[150]:= **(Coefficient**[**Expr8**, **AiryAi**[**ys1**] **AiryAi**[**ys2**] **AiryAiPrime**[**yt1**] **AiryAiPrime**[**yt2**]] /.
{**τ1** → **τ₁**, **τ2** → **τ₂**}) * **AiryAi**[**ys1**] **AiryAi**[**ys2**] **AiryAiPrime**[**yt1**] **AiryAiPrime**[**yt2**]

$$\text{Out[150]} = \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u (v-4) + 2 (v-4)) \text{Ai}(ys1) \text{Ai}(ys2) \text{Ai}'(yt1) \text{Ai}'(yt2)}{u^2 v}$$

In[151]:= **Coefficient**[Expr8, **AiryAiPrime**[ys1] **AiryAiPrime**[ys2] **AiryAi**[yt1] **AiryAi**[yt2]] /.
 $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[151]} = \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u (v-4) + 2 (v-4))}{u^2 v}$$

In[152]:= **Coefficient**[Expr8, **AiryAiPrime**[ys1] **AiryAi**[ys2] **AiryAiPrime**[yt1] **AiryAi**[yt2]] /.
 $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[152]} = \frac{\xi^2 \sqrt[3]{u+1} ((\tau_2^2 + 1)(u^2 + 2 u + 2)v - 2 \tau_2^2 u^2)}{u^{8/3} v^{2/3}}$$

In[153]:= **Coefficient**[Expr8, **AiryAi**[ys1] **AiryAiPrime**[ys2] **AiryAi**[yt1] **AiryAiPrime**[yt2]] /.
 $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[153]} = \frac{\xi^2 ((\tau_1^2 + 1)u^2 (v-2) + 2 \tau_1^2 u v + 2 \tau_1^2 v)}{\sqrt[3]{u^4 (u+1) v^4}}$$

In[154]:= **Coefficient**[Expr8, **AiryAiPrime**[ys1] **AiryAi**[ys2] **AiryAi**[yt1] **AiryAiPrime**[yt2]] /.
 $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[154]} = \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u v + 2 v)}{u^2 v}$$

In[155]:= **Coefficient**[Expr8, **AiryAi**[ys1] **AiryAiPrime**[ys2] **AiryAiPrime**[yt1] **AiryAi**[yt2]] /.
 $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[155]} = \frac{\xi^2 \tau_1 \tau_2 (u^2 (v-2) + 2 u v + 2 v)}{u^2 v}$$

In[156]:= **Coefficient**[Expr8, **AiryAiPrime**[ys1] **AiryAiPrime**[ys2]
AiryAiPrime[yt1] **AiryAiPrime**[yt2]] /. $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[156]} = \frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v-2) + 2 u (v-4) + 2 (v-4))}{u^2 \sqrt[3]{(u+1) v^4}}$$

In[157]:= **Coefficient**[Expr8, **AiryAiPrime**[ys1] **AiryAiPrime**[ys2]
AiryAiPrime[yt1] **AiryAiPrime**[yt2]] /. $\{\tau_1 \rightarrow \tau_1, \tau_2 \rightarrow \tau_2\}$

$$\text{Out[157]} = \frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v-2) + 2 u (v-4) + 2 (v-4))}{u^2 \sqrt[3]{(u+1) v^4}}$$

Explicit expressions for the y -
arguments of the Airy functions after the shift
They depend on τ_1^2 , but are independent of ρ_1

```
In[158]:= ys1new = Simplify[ysqp /. subsvσ /. subsu /. shift] /. {σ^2 → 1}
ys2new = Simplify[ystsfg /. subsvσ /. subsu /. shift] /. {σ^2 → 1}
yt1new = Simplify[ytqp /. subsvσ /. subsu /. shift] /. {σ^2 → 1}
yt2new = Simplify[ysttfg /. subsvσ /. subsu /. shift] /. {σ^2 → 1}
```

$$\text{Out[158]} = \frac{2 s \chi p + \xi (\tau^2 + 1) u}{2^{2/3} \xi \sqrt[3]{u \chi p^2}}$$

$$\text{Out[159]} = -\frac{4 s u \chi p - 2 \xi (\tau^2 + 1) (u + 1) v}{2 \times 2^{2/3} \xi u \chi p \sqrt[3]{\frac{u v + v}{u \chi p}}}$$

$$\text{Out[160]} = \frac{2 t \chi p + \xi (\tau^2 + 1) u}{2^{2/3} \xi \sqrt[3]{u \chi p^2}}$$

$$\text{Out[161]} = -\frac{4 t u \chi p - 2 \xi (\tau^2 + 1) (u + 1) v}{2 \times 2^{2/3} \xi u \chi p \sqrt[3]{\frac{u v + v}{u \chi p}}}$$

The integral over τ_1 is symmetric from $-\infty$ to $+\infty$;
 The Airy functions are even functions of τ_1 , hence,
 all terms proportional to τ_1 in an odd power give zero upon integration;

We remove such terms in this step

```
In[162]:= Expr9 = Simplify[(Expr8 + (Expr8 /. {τ1 → -τ1})) / 2]
Diffs9 = Diffs8
Coeff9 = Coeff8
Phase9 = Phase8
```

$$\begin{aligned} \text{Out[162]} = & \frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \text{Ai}(ys1) \text{Ai}(ys2) \text{Ai}(yt1) \text{Ai}(yt2) (32 s t \chi p^2 + \\ & \xi^2 (\tau^2 + 1) u^2 v (-2 \tau^2 + \tau^2 v + v) + 2 \xi^2 \tau^2 u v (-4 \tau^2 + \tau^2 v + v) + 2 \xi^2 \tau^2 (\tau^2 + 1) v^2 - 8 \xi^2 \tau^2 \tau^2 v) + \\ & \frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v-2) + 2 u (v-4) + 2 (v-4)) \text{Ai}'(ys1) \text{Ai}'(ys2) \text{Ai}'(yt1) \text{Ai}'(yt2)}{u^2 \sqrt[3]{(u+1) v^4}} + \\ & \frac{\xi^2 \sqrt[3]{u+1} \text{Ai}'(ys1) \text{Ai}(ys2) \text{Ai}'(yt1) \text{Ai}(yt2) ((\tau^2 + 1) (u^2 + 2 u + 2) v - 2 \tau^2 u^2)}{u^{8/3} v^{2/3}} + \\ & \frac{\xi^2 \text{Ai}(ys1) \text{Ai}'(ys2) \text{Ai}(yt1) \text{Ai}'(yt2) ((\tau^2 + 1) u^2 (v-2) + 2 \tau^2 u v + 2 \tau^2 v)}{\sqrt[3]{u^4 (u+1) v^4}} \end{aligned}$$

$$\text{Out[163]} = d\rho_1 d\rho_2 ds dt d\tau_1 d\tau_2 du dv$$

$$\text{Out[164]} = -\frac{\alpha^2 m^2 u}{\pi^2 L \phi \xi p_0 (u+1)^2 \sqrt{(v-4) v} \chi p (i\epsilon + s) (t - i\epsilon) ((\sigma^2 - 1) v - 4 \sigma^2)}$$

$$\text{Out[165]} = -((\rho_1 - \rho_2) (s - t))$$

Step 8: Integration over ρ_1, ρ_2, s and t ; one- and two-step contributions

Integrals over $d\rho_1 d\rho_2 ds dt$ are considered together and are regularised as described in section 6.1 in [arXiv 2605.12479] (eqs. (6.36) , (6.37))

The procedure amounts to

$$\int d\rho_1 d\rho_2 ds dt \text{Coeff}_9 * \text{Expr}_9 * \text{Exp}[-i (\rho_1 - \rho_2) (s - t)] \rightarrow I_{1st} + I_{2st}$$

where

$$I_{1st} = 2 \pi L_\phi \int \frac{ds}{s^2} \text{Coeff}_{1st} * (G[s] - G[-s] - 2 G[0])$$

$$I_{2st} = 2 \pi^2 L_\phi^2 \text{Coeff}_{2st} * G[0]$$

correspond to one - and two - step contributions, respectively

factor10 is an overall factor which we transfer from the Expr to Coeff

We introduce G_s and split Expr into two contributions, Expr_{1st} and Expr_{2st} , which we track separately

Phase is zero in this step, so we drop the corresponding variable

```
In[166]:= factor10 = 8 m^4;
Gs = Expr9 / factor10 /. {t -> s} /. {yt1 -> ys1, yt2 -> ys2}
Phase10 = Phase9 /. {t -> s}

Out[167]= 
$$\frac{1}{8 m^4} \left( \frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \text{Ai}(ys1)^2 \text{Ai}(ys2)^2 (32 s^2 \chi p^2 + \xi^2 (\tau^2 + 1) u^2 v (-2 \tau^2 + \tau^2 v + v) + \right.$$


$$2 \xi^2 \tau^2 u v (-4 \tau^2 + \tau^2 v + v) + 2 \xi^2 \tau^2 (\tau^2 + 1) v^2 - 8 \xi^2 \tau^2 \tau^2 v) +$$


$$\frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v - 2) + 2 u (v - 4) + 2 (v - 4)) \text{Ai}'(ys1)^2 \text{Ai}'(ys2)^2}{u^2 \sqrt[3]{(u+1) v^4}} +$$


$$\frac{\xi^2 \sqrt[3]{u+1} \text{Ai}'(ys1)^2 \text{Ai}(ys2)^2 ((\tau^2 + 1) (u^2 + 2 u + 2) v - 2 \tau^2 u^2)}{u^{8/3} v^{2/3}} +$$


$$\left. \frac{\xi^2 \text{Ai}(ys1)^2 \text{Ai}'(ys2)^2 ((\tau^2 + 1) u^2 (v - 2) + 2 \tau^2 u v + 2 \tau^2 v)}{\sqrt[3]{u^4 (u+1) v^4}} \right)$$

```

Out[168]= 0

One - step contribution

$ys1$ and $ys2$ are $y_{1,2} (s)$ arguments, introduced above
 $yms1$ and $yms1$ are the same arguments with $s \rightarrow -s$

y01 and y02 are given by $y_{1,2}(0)$

```
In[169]:= Diffss101st = Diffss9 /. {dt → 1, dρ1 → 1, dρ2 → 1}
Coeff101st = Simplify[Coeff9 * 2 π L ϕ * factor10 /. {t → s} /. {ie → 0}]
Expr101st =
  Gs + (Gs /. {s → -s, ys1 → yms1, ys2 → yms2}) - 2 (Gs /. {s → 0, ys1 → y01, ys2 → y02})
Out[169]= ds dτ1 dτ2 du dv
```

$$\text{Out[170]} = -\frac{16 \alpha^2 m^6 u}{\pi \xi p_0 s^2 (u+1)^2 \sqrt{(v-4)v} \chi p ((\sigma^2 - 1)v - 4\sigma^2)}$$

$$\text{Out[171]} = -\frac{1}{4m^4} \left(\frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \right. \\
\begin{aligned} & (-8 v \tau_1^2 \tau_2^2 \xi^2 + 2 v^2 \tau_1^2 (\tau_2^2 + 1) \xi^2 + 2 u v \tau_1^2 (v \tau_2^2 - 4 \tau_2^2 + v) \xi^2 + u^2 v (\tau_1^2 + 1) (v \tau_2^2 - 2 \tau_2^2 + v) \xi^2) \\ & \text{Ai}(y_{01})^2 \text{Ai}(y_{02})^2 + \frac{\sqrt[3]{u+1} \xi^2 ((u^2 + 2u + 2)v (\tau_2^2 + 1) - 2u^2 \tau_2^2) \text{Ai}'(y_{01})^2 \text{Ai}(y_{02})^2}{u^{8/3} v^{2/3}} + \\ & \frac{\xi^2 ((v-2)(\tau_1^2 + 1)u^2 + 2 v \tau_1^2 u + 2 v \tau_1^2) \text{Ai}(y_{01})^2 \text{Ai}'(y_{02})^2}{\sqrt[3]{u^4 (u+1) v^4}} + \\ & \left. \frac{2^{2/3} ((v-2)u^2 + 2(v-4)u + 2(v-4)) \xi^2 \chi p^{2/3} \text{Ai}'(y_{01})^2 \text{Ai}'(y_{02})^2}{u^2 \sqrt[3]{(u+1) v^4}} \right) + \\ & \frac{1}{8m^4} \left(\frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \right. \\ & \begin{aligned} & (-8 v \tau_1^2 \tau_2^2 \xi^2 + 2 v^2 \tau_1^2 (\tau_2^2 + 1) \xi^2 + 2 u v \tau_1^2 (v \tau_2^2 - 4 \tau_2^2 + v) \xi^2 + \\ & u^2 v (\tau_1^2 + 1) (v \tau_2^2 - 2 \tau_2^2 + v) \xi^2 + 32 s^2 \chi p^2) \text{Ai}(y_{ms1})^2 \text{Ai}(y_{ms2})^2 + \\ & \frac{\sqrt[3]{u+1} \xi^2 ((u^2 + 2u + 2)v (\tau_2^2 + 1) - 2u^2 \tau_2^2) \text{Ai}'(y_{ms1})^2 \text{Ai}(y_{ms2})^2}{u^{8/3} v^{2/3}} + \\ & \frac{\xi^2 ((v-2)(\tau_1^2 + 1)u^2 + 2 v \tau_1^2 u + 2 v \tau_1^2) \text{Ai}(y_{ms1})^2 \text{Ai}'(y_{ms2})^2}{\sqrt[3]{u^4 (u+1) v^4}} + \\ & \left. \frac{2^{2/3} ((v-2)u^2 + 2(v-4)u + 2(v-4)) \xi^2 \chi p^{2/3} \text{Ai}'(y_{ms1})^2 \text{Ai}'(y_{ms2})^2}{u^2 \sqrt[3]{(u+1) v^4}} \right) + \\ & \frac{1}{8m^4} \left(\frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \right. \\ & \begin{aligned} & (-8 v \tau_1^2 \tau_2^2 \xi^2 + 2 v^2 \tau_1^2 (\tau_2^2 + 1) \xi^2 + 2 u v \tau_1^2 (v \tau_2^2 - 4 \tau_2^2 + v) \xi^2 + \\ & u^2 v (\tau_1^2 + 1) (v \tau_2^2 - 2 \tau_2^2 + v) \xi^2 + 32 s^2 \chi p^2) \text{Ai}(y_{s1})^2 \text{Ai}(y_{s2})^2 + \\ & \frac{\sqrt[3]{u+1} \xi^2 ((u^2 + 2u + 2)v (\tau_2^2 + 1) - 2u^2 \tau_2^2) \text{Ai}'(y_{s1})^2 \text{Ai}(y_{s2})^2}{u^{8/3} v^{2/3}} + \\ & \frac{\xi^2 ((v-2)(\tau_1^2 + 1)u^2 + 2 v \tau_1^2 u + 2 v \tau_1^2) \text{Ai}(y_{s1})^2 \text{Ai}'(y_{s2})^2}{\sqrt[3]{u^4 (u+1) v^4}} + \\ & \left. \frac{2^{2/3} ((v-2)u^2 + 2(v-4)u + 2(v-4)) \xi^2 \chi p^{2/3} \text{Ai}'(y_{s1})^2 \text{Ai}'(y_{s2})^2}{u^2 \sqrt[3]{(u+1) v^4}} \right) \end{aligned} \end{aligned}$$

Two – step contribution

Note that the integral over s is carried out in this term

```
In[172]:= Diffsl02 st = Diffs9 /. {ds → 1, dt → 1, dp1 → 1, dp2 → 1}
Coeffl02 st = Simplify[Coeff9 * 2 π^2 Lφ^2 * factor10 /. {t → 1, s → 1} /. {ie → 0}]
Exprl02 st = Gs /. {s → 0, ys1 → y01, ys2 → y02}
```

```
Out[172]= dr1 dτ2 du dv
```

$$\text{Out[173]} = -\frac{16 \alpha^2 L \phi m^6 u}{\xi p_0 (u+1)^2 \sqrt{(v-4)v} \chi p ((\sigma^2-1)v - 4\sigma^2)}$$

$$\begin{aligned} \text{Out[174]} = & \frac{1}{8 m^4} \left(\frac{2^{2/3} \xi^2 \chi p^{2/3} (u^2 (v-2) + 2u(v-4) + 2(v-4)) \text{Ai}'(y_{01})^2 \text{Ai}'(y_{02})^2}{u^2 \sqrt[3]{(u+1)v^4}} + \frac{1}{2^{2/3} u^2 v^{5/3} \chi p^{2/3}} \sqrt[3]{u+1} \text{Ai}(y_{01})^2 \text{Ai}(y_{02})^2 \right. \\ & \frac{(\xi^2 (\tau^2 + 1) u^2 v (-2\tau^2 + \tau^2 v + v) + 2\xi^2 \tau^2 u v (-4\tau^2 + \tau^2 v + v) + 2\xi^2 \tau^2 (\tau^2 + 1) v^2 - 8\xi^2 \tau^2 \tau^2 v) +}{u^{8/3} v^{2/3}} \\ & \left. \frac{\xi^2 \sqrt[3]{u+1} \text{Ai}'(y_{01})^2 \text{Ai}(y_{02})^2 ((\tau^2 + 1)(u^2 + 2u + 2)v - 2\tau^2 u^2)}{u^{8/3} v^{2/3}} + \right. \\ & \left. \frac{\xi^2 \text{Ai}(y_{01})^2 \text{Ai}'(y_{02})^2 ((\tau^2 + 1)u^2 (v-2) + 2\tau^2 u v + 2\tau^2 v)}{\sqrt[3]{u^4 (u+1)v^4}} \right) \end{aligned}$$

Step 9: Integration over τ

Integrate over τ

$$\int d\tau \tau^{2k} \text{Ai}^2(a + x \tau^2) = \frac{\Gamma(k+1/2)}{2^{2k/3+1} \pi^{1/2} x^{k+1/2}} \text{Ai}_{k+1}(2^{2/3} a)$$

$$\int d\tau \text{Ai}^2(a + x \tau^2) = \frac{1}{2 x^{1/2}} \text{Ai}_1(2^{2/3} a)$$

$$\int d\tau \tau^2 \text{Ai}^2(a + x \tau^2) =$$

$$\frac{1}{4 \times 2^{2/3} x^{3/2}} \text{Ai}_2(2^{2/3} a) = \frac{1}{4 \times 2^{2/3} x^{3/2}} [-\text{Ai}'(2^{2/3} a) - 2^{2/3} a \text{Ai}_1(2^{2/3} a)]$$

$$\int d\tau \text{Ai}'^2(a + x \tau^2) = \frac{1}{4 \times 2^{2/3} x^{1/2}} [-3 \text{Ai}'(2^{2/3} a) - 2^{2/3} a \text{Ai}_1(2^{2/3} a)]$$

$$\text{Ai}_2(y) = -\text{Ai}'(y) - y \text{Ai}_1(y)$$

$ys_i = a_i + x_i \tau_i$, see a_i and x_i below

We will perform the integration for G_s

and express the overall result via integrated G_s

Explicit representation for the Airy function arguments

```
In[175]:= Expand[PowerExpand[ys1new]]
a1expl = (u / 2 / xp) ^ (2 / 3) Expand[Simplify[(% / (u / 2 / xp) ^ (2 / 3) /. {tau1 -> 0})]]
x1expl = Coefficient[%%, tau1^2]
```

$$\text{Out[175]} = \frac{\sqrt[3]{2} s \sqrt[3]{xp}}{\xi \sqrt[3]{u}} + \frac{\tau^2 u^{2/3}}{2^{2/3} xp^{2/3}} + \frac{u^{2/3}}{2^{2/3} xp^{2/3}}$$

$$\text{Out[176]} = \frac{\left(\frac{u}{xp}\right)^{2/3} \left(\frac{2 s xp}{\xi u} + 1\right)}{2^{2/3}}$$

$$\text{Out[177]} = \frac{u^{2/3}}{2^{2/3} xp^{2/3}}$$

```
In[178]:= 2 ^ (2 / 3) \left( \frac{(u + 1) v}{4 u xp} \right)^{2/3} *
```

```
Collect[Expand[PowerExpand[ys2new / 2 ^ (2 / 3) / \left( \frac{(u + 1) v}{4 u xp} \right)^{2/3}]], tau2, Simplify]
```

```
a2expl = % /. {tau2 -> 0}
x2expl = Coefficient[%%, tau2^2]
```

$$\text{Out[178]} = \frac{\left(\frac{(u+1)v}{u xp}\right)^{2/3} \left(-\frac{2 s u xp}{\xi u v + \xi v} + \tau^2 + 1\right)}{2^{2/3}}$$

$$\text{Out[179]} = \frac{\left(\frac{(u+1)v}{u xp}\right)^{2/3} \left(1 - \frac{2 s u xp}{\xi u v + \xi v}\right)}{2^{2/3}}$$

$$\text{Out[180]} = \frac{\left(\frac{(u+1)v}{u xp}\right)^{2/3}}{2^{2/3}}$$

```
In[181]:= 2 ^ (2 / 3) Simplify[a1expl, Assumptions -> assumpAll]
Simplify[% /. {s -> 0}, Assumptions -> assumpAll]
2 ^ (2 / 3) \left( \frac{(u + 1) v}{4 u xp} \right)^{2/3} Simplify[a2expl / \left( \frac{(u + 1) v}{4 u xp} \right)^{2/3}, Assumptions -> assumpAll]
Simplify[% /. {s -> 0}, Assumptions -> assumpAll]
```

$$\text{Out[181]} = \left(\frac{u}{xp}\right)^{2/3} \left(\frac{2 s xp}{\xi u} + 1\right)$$

$$\text{Out[182]} = \left(\frac{u}{xp}\right)^{2/3}$$

$$\text{Out[183]} = \left(\frac{(u+1)v}{u xp}\right)^{2/3} \left(1 - \frac{2 s u xp}{\xi u v + \xi v}\right)$$

$$\text{Out[184]} = \left(\frac{(u+1)v}{u xp}\right)^{2/3}$$

The following expression is used later for collecting terms


```
In[185]:= collectAirya = {AiryAiPrime[22/3 a1] AiryAiPrime[22/3 a2],
  Ai1[22/3 a1] Ai1[22/3 a2],
  AiryAiPrime[22/3 a1] Ai1[22/3 a2],
  Ai1[22/3 a1] AiryAiPrime[22/3 a2]};
```

Integration of Gs over $\tau_{1,2}$ is
implemented in this step via substitutions in Gs

Result is stored in GsIntegrated

```
In[186]:= Collect[Gs, {τ1, AiryAi[ys1]2, AiryAiPrime[ys1]2}, Simplify] /.
  {τ12 AiryAi[ys1]2 →
     $\frac{1}{4 \times 2^{2/3} x \exp^{3/2}} (-\text{AiryAiPrime}[2^{2/3} a1] - 2^{2/3} a1 \exp^{1/2} \text{Ai}_1[2^{2/3} a1])$  } /.
  {AiryAi[ys1]2 →  $\frac{1}{2 x \exp^{1/2}} \text{Ai}_1[2^{2/3} a1]$ , AiryAiPrime[ys1]2 →
     $\frac{1}{4 \times 2^{2/3} x \exp^{1/2}} (-3 \text{AiryAiPrime}[2^{2/3} a1] - 2^{2/3} a1 \exp^{1/2} \text{Ai}_1[2^{2/3} a1])$  };
Expand[%];
Collect[%, {τ2, AiryAi[ys2]2, AiryAiPrime[ys2]2}, Simplify] /.
  {τ22 AiryAi[ys2]2 →
     $\frac{1}{4 \times 2^{2/3} x2 \exp^{3/2}} (-\text{AiryAiPrime}[2^{2/3} a2] - 2^{2/3} a2 \exp^{1/2} \text{Ai}_1[2^{2/3} a2])$  } /.
  {AiryAi[ys2]2 →  $\frac{1}{2 x2 \exp^{1/2}} \text{Ai}_1[2^{2/3} a2]$ , AiryAiPrime[ys2]2 →
     $\frac{1}{4 \times 2^{2/3} x2 \exp^{1/2}} (-3 \text{AiryAiPrime}[2^{2/3} a2] - 2^{2/3} a2 \exp^{1/2} \text{Ai}_1[2^{2/3} a2])$  };
PowerExpand[%];
GsIntegrated = Collect[Expand[%],
  collectAirya, Simplify]
```

```
Out[190]:= 
$$-\frac{1}{8 m^4 u^3 (u+1) v^2} \text{Ai}_1(2^{2/3} a1) \text{Ai}_1(2^{2/3} a2) (u v (\xi^2 + 2 s^2 \chi p^2 + 4 \xi s \chi p) - 12 s^2 u \chi p^2 +$$


$$u^3 (\xi + s \chi p) (v (\xi + s \chi p) - 2 s \chi p) + 2 u^2 (v (\xi + s \chi p)^2 - s \chi p (\xi + 6 s \chi p)) + 2 \xi s v \chi p) -$$


$$\frac{\xi (u^2 + 2 u + 2) \chi p^{2/3} \text{Ai}'(2^{2/3} a1) \text{Ai}_1(2^{2/3} a2) (v (\xi + s u \chi p + \xi u) - 2 s u \chi p)}{8 m^4 u^{8/3} (u+1) v^2} +$$


$$\frac{\xi (v - 2) \chi p^{2/3} \text{Ai}_1(2^{2/3} a1) \text{Ai}'(2^{2/3} a2) (u^2 (\xi + s \chi p) + u (\xi + 2 s \chi p) + 2 s \chi p)}{8 m^4 \sqrt[3]{u^7} (u+1)^2 v^5} +$$


$$\frac{\xi^2 \chi p^{4/3} \text{Ai}'(2^{2/3} a1) \text{Ai}'(2^{2/3} a2) (u^2 (v - 2) + u (2 v - 5) + 2 v - 5)}{8 m^4 u^2 (u+1)^{2/3} v^{5/3}}$$

```

One step term ;
Expressed via an integral over s

(equivalent of variable T in [arXiv 2605.12479] in Eq. (6.36))

```
In[191]:= Diffs111st = Diffs101st /. {dτ1 → 1, dτ2 → 1}
Coeff111st = Coeff101st
Expr111st = (GsIntegrated /. {a1 → a1expl, a2 → a2expl}) +
  (GsIntegrated /. {a1 → a1expl, a2 → a2expl} /. {s → -s}) -
  2 (GsIntegrated /. {a1 → a1expl, a2 → a2expl} /. {s → 0})
Out[191]= ds du dv
```

$$\text{Out[192]} = -\frac{16 \alpha^2 m^6 u}{\pi \xi p_0 s^2 (u+1)^2 \sqrt{(v-4)v} \chi_P((\sigma^2-1)v-4\sigma^2)}$$

Out[193]=

$$\begin{aligned}
& \frac{((v-2)u^2 + (2v-5)u + 2v-5)\chi p^{4/3} \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(\frac{2s\chi p}{u\xi} + 1\right)\right) \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(1 - \frac{2su\chi p}{uv\xi + v\xi}\right)\right) \xi^2}{8m^4 u^2 (u+1)^{2/3} v^{5/3}} + \\
& \frac{((v-2)u^2 + (2v-5)u + 2v-5)\chi p^{4/3} \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(1 - \frac{2s\chi p}{u\xi}\right)\right) \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(\frac{2su\chi p}{uv\xi + v\xi} + 1\right)\right) \xi^2}{8m^4 u^2 (u+1)^{2/3} v^{5/3}} + \\
& \frac{(v-2)\chi p^{2/3} ((\xi - s\chi p)u^2 + (\xi - 2s\chi p)u - 2s\chi p) \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(\frac{2su\chi p}{uv\xi + v\xi} + 1\right)\right) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(1 - \frac{2s\chi p}{u\xi}\right)\right) \xi}{8m^4 \sqrt[3]{u^7 (u+1)^2 v^5}} + \\
& \frac{(v-2)\chi p^{2/3} ((\xi + s\chi p)u^2 + (\xi + 2s\chi p)u + 2s\chi p) \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(1 - \frac{2su\chi p}{uv\xi + v\xi}\right)\right) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(\frac{2s\chi p}{u\xi} + 1\right)\right) \xi}{8m^4 \sqrt[3]{u^7 (u+1)^2 v^5}} - \\
& \frac{(u^2 + 2u + 2)\chi p^{2/3} (v(u\xi + \xi + su\chi p) - 2su\chi p) \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(\frac{2s\chi p}{u\xi} + 1\right)\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(1 - \frac{2su\chi p}{uv\xi + v\xi}\right)\right) \xi}{8m^4 u^{8/3} (u+1)v^2} - \\
& \frac{(u^2 + 2u + 2)\chi p^{2/3} (2su\chi p + v(u\xi + \xi - su\chi p)) \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(1 - \frac{2s\chi p}{u\xi}\right)\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(\frac{2su\chi p}{uv\xi + v\xi} + 1\right)\right) \xi}{8m^4 u^{8/3} (u+1)v^2} - \\
& 2 \left(\frac{((v-2)u^2 + (2v-5)u + 2v-5)\chi p^{4/3} \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \xi^2}{8m^4 u^2 (u+1)^{2/3} v^{5/3}} + \right. \\
& \frac{(v-2)(\xi u^2 + \xi u)\chi p^{2/3} \operatorname{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \xi}{8m^4 \sqrt[3]{u^7 (u+1)^2 v^5}} - \\
& \frac{(u^2 + 2u + 2)(u\xi + \xi)\chi p^{2/3} \operatorname{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \xi}{8m^4 u^{8/3} (u+1)v} - \\
& \left. \frac{(v\xi^2 u^3 + 2v\xi^2 u^2 + v\xi^2 u) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right)}{8m^4 u^3 (u+1)v^2} \right) - \frac{1}{8m^4 u^3 (u+1)v^2} \\
& ((\xi + s\chi p)(v(\xi + s\chi p) - 2s\chi p)u^3 + 2(v(\xi + s\chi p)^2 - s\chi p(\xi + 6s\chi p))u^2 - 12s^2\chi p^2 u + \\
& v(\xi^2 + 4s\chi p\xi + 2s^2\chi p^2)u + 2sv\xi\chi p) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(\frac{2s\chi p}{u\xi} + 1\right)\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(1 - \frac{2su\chi p}{uv\xi + v\xi}\right)\right) - \\
& \frac{1}{8m^4 u^3 (u+1)v^2} ((\xi - s\chi p)(2s\chi p + v(\xi - s\chi p))u^3 + 2(v(\xi - s\chi p)^2 + s\chi p(\xi - 6s\chi p))u^2 - 12s^2\chi p^2 u + \\
& v(\xi^2 - 4s\chi p\xi + 2s^2\chi p^2)u - 2sv\xi\chi p) \operatorname{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3} \left(1 - \frac{2s\chi p}{u\xi}\right)\right) \operatorname{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3} \left(\frac{2su\chi p}{uv\xi + v\xi} + 1\right)\right)
\end{aligned}$$

Two - step term

```

In[194]:= Diffss112 st = Diffss102 st /. {dτ1 → 1, dτ2 → 1}
Coeff112 st = Coeff102 st
Expr112 st = GsIntegrated /. {a1 → a1expl, a2 → a2expl} /. {s → 0}
Out[194]= du dv

```

$$\begin{aligned}
\text{Out[195]} &= -\frac{16 \alpha^2 L \phi m^6 u}{\xi p_0 (u+1)^2 \sqrt{(v-4)v} \chi p ((\sigma^2 - 1)v - 4 \sigma^2)} \\
\text{Out[196]} &= -\frac{\xi (u^2 + 2u + 2) \chi p^{2/3} (\xi + \xi u) \text{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{8 m^4 u^{8/3} (u+1)v} + \\
&\quad \frac{\xi (v-2) \chi p^{2/3} (\xi u^2 + \xi u) \text{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \text{Ai}'\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right) - \text{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) (\xi^2 u^3 v + 2 \xi^2 u^2 v + \xi^2 u v) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{8 m^4 \sqrt[3]{u^7} (u+1)^2 v^5} - \frac{\text{Ai}_1\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) (\xi^2 u^3 v + 2 \xi^2 u^2 v + \xi^2 u v) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{8 m^4 u^3 (u+1)v^2} + \\
&\quad \frac{\xi^2 \chi p^{4/3} (u^2 (v-2) + u (2v-5) + 2v-5) \text{Ai}'\left(\left(\frac{u}{\chi p}\right)^{2/3}\right) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{8 m^4 u^2 (u+1)^{2/3} v^{5/3}}
\end{aligned}$$

Step 10: Rewriting the expressions to shape them into final result

We pass to variable $\lambda = s \frac{2 m^2 u \chi_p}{\xi (1+u)}$,
which is the virtuality of the intermediate photon

```

In[197]:= a1expl
targ = u ^ (2 / 3) / χp ^ (2 / 3) (1 + (1 + u) / u ^ 2 * λ / m ^ 2)
Simplify[2 ^ (2 / 3) * %% == targ]
stoλ = Solve[%, s][[1]]
ξarg = Simplify[2 ^ (2 / 3) a2expl /. stoλ]
jacstoλ = D[s /. stoλ, λ]

```

$$\text{Out[197]} = \frac{\left(\frac{u}{\chi p}\right)^{2/3} \left(\frac{2 s \chi p}{\xi u} + 1\right)}{2^{2/3}}$$

$$\text{Out[198]} = \frac{u^{2/3} \left(\frac{\lambda (u+1)}{m^2 u^2} + 1\right)}{\chi p^{2/3}}$$

$$\text{Out[199]} = 2 m s u \chi p = \frac{\lambda \xi (u+1)}{m}$$

$$\text{Out[200]} = \left\{ s \rightarrow \frac{\lambda \xi + \lambda \xi u}{2 m^2 u \chi p} \right\}$$

$$\text{Out[201]} = \frac{(m^2 v - \lambda) \left(\frac{(u+1)v}{u \chi p}\right)^{2/3}}{m^2 v}$$

$$\text{Out[202]} = \frac{\xi + \xi u}{2 m^2 u \chi p}$$

factor12 is an overall factor, which we transfer from Expr to Coeff

$\phi\lambda$ is the new function resulting from
passing from s to λ in G_s and diving it by factor12

The Airy function arguments are now

$$t = 2^{2/3} a_1$$

$$\xi = 2^{2/3} a_2$$

```
In[203]:= factor12 = (u + 1) ξ^2 / (8 u^2 v m^4);
φλ = Collect[GsIntegrated / factor12 /. stoλ, collectAirya, Simplify] /.
{a1 → t / 2^(2/3), a2 → ξ / 2^(2/3)}
```

$$\begin{aligned} \text{Out[204]} = & -\frac{(u^2 + 2u + 2) \text{Ai}_1(\xi) \text{Ai}'_1(t) \left(\frac{\chi p}{u}\right)^{2/3} (2m^2 v + \lambda(v-2))}{2m^2 (u+1)v} + \frac{(v-2) \text{Ai}_1(t) \text{Ai}'_1(\xi) (2m^2 u^2 + \lambda(u^2 + 2u + 2)) \left(\frac{\chi p}{u v + v}\right)^{2/3}}{2m^2 u^{4/3}} + \\ & \frac{\text{Ai}_1(\xi) \text{Ai}_1(t) (-4m^4 u^2 v - 4\lambda m^2 (u^2(v-1) + uv + v) + \lambda^2 (-u^2(v-2) - 2u(v-6) - 2(v-6)))}{4m^4 u^2 v} + \\ & \frac{\chi p^{4/3} \text{Ai}'_1(\xi) \text{Ai}'_1(t) (u^2(v-2) + u(2v-5) + 2v-5)}{(u+1)^{5/3} v^{2/3}} \end{aligned}$$

```
In[205]:= Diffs12_1st = Diffs11_1st /. {ds → dλ}
Coeff12_1st = Simplify[(Coeff11_1st * jacstoλ * factor12) /. stoλ]
Expr12_1st = (φλ /. {t → targ, ξ → ξarg}) +
(φλ /. {t → targ, ξ → ξarg} /. {λ → -λ}) - 2 (φλ /. {t → targ, ξ → ξarg} /. {λ → 0})
```

Out[205]= $d\lambda du dv$

$$\text{Out[206]} = -\frac{4\alpha^2 m^4}{\pi \lambda^2 p_0 (u+1)^2 v \sqrt{(v-4)v} ((\sigma^2 - 1)v - 4\sigma^2)}$$

$$\begin{aligned}
& \text{Out[207]=} \frac{((v-2)u^2 + (2v-5)u + 2v-5) \text{Ai}'\left(\frac{(vm^2+\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \text{Ai}'\left(\frac{u^{2/3}\left(1-\frac{(u+1)\lambda}{m^2 u^2}\right)}{\chi p^{2/3}}\right) \chi p^{4/3}}{(u+1)^{5/3} v^{2/3}} + \\
& \frac{((v-2)u^2 + (2v-5)u + 2v-5) \text{Ai}'\left(\frac{(m^2 v-\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \text{Ai}'\left(\frac{u^{2/3}\left(\frac{(u+1)\lambda}{m^2 u^2}+1\right)}{\chi p^{2/3}}\right) \chi p^{4/3}}{(u+1)^{5/3} v^{2/3}} - \\
& \frac{(u^2 + 2u + 2)(2vm^2 + (v-2)\lambda)\left(\frac{\chi p}{u}\right)^{2/3} \text{Ai}'\left(\frac{u^{2/3}\left(\frac{(u+1)\lambda}{m^2 u^2}+1\right)}{\chi p^{2/3}}\right) \text{Ai}_1\left(\frac{(m^2 v-\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right)}{2m^2(u+1)v} - \\
& \frac{(u^2 + 2u + 2)(2m^2 v - (v-2)\lambda)\left(\frac{\chi p}{u}\right)^{2/3} \text{Ai}'\left(\frac{u^{2/3}\left(1-\frac{(u+1)\lambda}{m^2 u^2}\right)}{\chi p^{2/3}}\right) \text{Ai}_1\left(\frac{(vm^2+\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right)}{2m^2(u+1)v} - \\
& 2 \left(\frac{((v-2)u^2 + (2v-5)u + 2v-5) \text{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \text{Ai}'\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \chi p^{4/3}}{(u+1)^{5/3} v^{2/3}} - \frac{(u^2 + 2u + 2)\left(\frac{\chi p}{u}\right)^{2/3} \text{Ai}'\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \text{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right)}{u+1} + \right. \\
& \left. u^{2/3}(v-2)\left(\frac{\chi p}{uv+v}\right)^{2/3} \text{Ai}'\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \text{Ai}_1\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) - \text{Ai}_1\left(\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}\right) \text{Ai}_1\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \right) + \\
& \frac{(v-2)(2m^2 u^2 - (u^2 + 2u + 2)\lambda)\left(\frac{\chi p}{uv+v}\right)^{2/3} \text{Ai}'\left(\frac{(vm^2+\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \text{Ai}_1\left(\frac{u^{2/3}\left(1-\frac{(u+1)\lambda}{m^2 u^2}\right)}{\chi p^{2/3}}\right)}{2m^2 u^{4/3}} + \frac{1}{4m^4 u^2 v} \\
& (-4u^2 v m^4 + 4((v-1)u^2 + vu + v)\lambda m^2 + (-(v-2)u^2) - 2(v-6)u - 2(v-6)\lambda^2) \text{Ai}_1\left(\frac{(vm^2+\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \\
& \text{Ai}_1\left(\frac{u^{2/3}\left(1-\frac{(u+1)\lambda}{m^2 u^2}\right)}{\chi p^{2/3}}\right) + \frac{(v-2)(2m^2 u^2 + (u^2 + 2u + 2)\lambda)\left(\frac{\chi p}{uv+v}\right)^{2/3} \text{Ai}'\left(\frac{(m^2 v-\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \text{Ai}_1\left(\frac{u^{2/3}\left(\frac{(u+1)\lambda}{m^2 u^2}+1\right)}{\chi p^{2/3}}\right)}{2m^2 u^{4/3}} + \\
& \frac{1}{4m^4 u^2 v} (-4u^2 v m^4 - 4((v-1)u^2 + vu + v)\lambda m^2 + (-(v-2)u^2) - 2(v-6)u - 2(v-6)\lambda^2) \\
& \text{Ai}_1\left(\frac{(m^2 v-\lambda)\left(\frac{(u+1)v}{u\chi p}\right)^{2/3}}{m^2 v}\right) \text{Ai}_1\left(\frac{u^{2/3}\left(\frac{(u+1)\lambda}{m^2 u^2}+1\right)}{\chi p^{2/3}}\right)
\end{aligned}$$

```
In[208]:= Diff12st = Diff11st
          Coeff12st = Factor[Simplify[(Coeff11st * factor12) /. stoλ]]
          Expr12st = (φλ /. {t → targ, ξ → ξarg} /. {λ → 0})
```

```
Out[208]= du dv
```

$$\text{Out[209]} = -\frac{2 \alpha^2 L \phi m^2 \xi}{p_0 u (u+1) v \sqrt{(v-4)v} \chi p (-4 \sigma^2 + \sigma^2 v - v)}$$

$$\begin{aligned} \text{Out[210]} = & -\text{Ai}_1\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right) + u^{2/3} (v-2) \text{Ai}_1\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \left(\frac{\chi p}{u v + v}\right)^{2/3} \text{Ai}'\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right) - \\ & \frac{(u^2 + 2 u + 2) \left(\frac{\chi p}{u}\right)^{2/3} \text{Ai}'\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \text{Ai}_1\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{u+1} + \frac{\chi p^{4/3} (u^2 (v-2) + u (2 v - 5) + 2 v - 5) \text{Ai}'\left(\frac{u^{2/3}}{\chi p^{2/3}}\right) \text{Ai}'\left(\left(\frac{(u+1)v}{u \chi p}\right)^{2/3}\right)}{(u+1)^{5/3} v^{2/3}} \end{aligned}$$

The last two cells are our final result,
which can be found in [arXiv 2605.12479]

For the one – step contribution

($\phi_{\text{tree}}(\lambda)$ is equivalent of $\phi\lambda$ used in this notebook), and $\chi = \chi p$

$$W_{\text{dir}}^{1\text{st}} = \frac{\alpha^2 m^4}{\pi p^0} \int_0^\infty du \int_4^\infty dv \int_0^\infty \frac{d\lambda}{\lambda^2} \frac{\phi_{\text{tree}}(\lambda) + \phi_{\text{tree}}(-\lambda) - 2\phi_{\text{tree}}(0)}{(1+u)^2 v^{3/2} \sqrt{v-4}}, \quad (6.54)$$

with

$$\begin{aligned} \phi_{\text{tree}}(\lambda) = & \chi^{4/3} \frac{u^2(v-2) + (1+u)(2v-5)}{(1+u)^{5/3} v^{2/3}} \text{Ai}'(t) \text{Ai}'(\zeta) \\ & + \left[\frac{\chi u}{(1+u)v} \right]^{2/3} \left(1 + \frac{\lambda}{m^2} \frac{u^2 + 2u + 2}{2u^2} \right) (v-2) \text{Ai}_1(t) \text{Ai}'(\zeta) \\ & - \left(\frac{\chi}{u} \right)^{2/3} \frac{u^2 + 2u + 2}{1+u} \left(1 + \frac{\lambda}{m^2} \frac{v-2}{2v} \right) \text{Ai}'(t) \text{Ai}_1(\zeta) \\ & - \left[1 + \frac{\lambda}{m^2} \left(\frac{u^2 + u + 1}{u^2} - \frac{1}{v} \right) \right. \\ & \quad \left. + \frac{\lambda^2}{m^4} \frac{u^2(v-2) + 2(1+u)(v-6)}{4u^2 v} \right] \text{Ai}_1(t) \text{Ai}_1(\zeta). \end{aligned} \quad (6.55)$$

In[211]:=

$$W_{\text{dir}}^{1\text{st}} = \frac{\alpha^2 m^4}{\pi p^0} \int_0^\infty du \int_4^\infty dv \int_0^\infty \frac{d\lambda}{\lambda^2} \frac{\phi_{\text{tree}}(\lambda) + \phi_{\text{tree}}(-\lambda) - 2\phi_{\text{tree}}(0)}{(1+u)^2 v^{3/2} \sqrt{v-4}}, \quad (6.54)$$

with

$$\begin{aligned} \phi_{\text{tree}}(\lambda) = & \chi^{4/3} \frac{u^2(v-2) + (1+u)(2v-5)}{(1+u)^{5/3} v^{2/3}} \text{Ai}'(t) \text{Ai}'(\zeta) \\ & + \left[\frac{\chi u}{(1+u)v} \right]^{2/3} \left(1 + \frac{\lambda}{m^2} \frac{u^2 + 2u + 2}{2u^2} \right) (v-2) \text{Ai}_1(t) \text{Ai}'(\zeta) \\ & - \left(\frac{\chi}{u} \right)^{2/3} \frac{u^2 + 2u + 2}{1+u} \left(1 + \frac{\lambda}{m^2} \frac{v-2}{2v} \right) \text{Ai}'(t) \text{Ai}_1(\zeta) \\ & - \left[1 + \frac{\lambda}{m^2} \left(\frac{u^2 + u + 1}{u^2} - \frac{1}{v} \right) \right. \\ & \quad \left. + \frac{\lambda^2}{m^4} \frac{u^2(v-2) + 2(1+u)(v-6)}{4u^2 v} \right] \text{Ai}_1(t) \text{Ai}_1(\zeta). \end{aligned} \quad (6.55)$$

Out[211]=

The two –

step result obtained here is equivalent to the following expression

$$W_{\text{dir},2\text{-loop}}^{2\text{st}}(\chi) = \frac{\alpha^2 m^2 \theta(\chi)}{2p^0 \chi} \xi L_\varphi \int_0^\infty du \int_4^\infty dv \frac{\phi(0)}{u(1+u)v^{3/2} \sqrt{v-4}}, \quad (6.53)$$

In[212]:=

$$W_{\text{dir},2\text{-loop}}^{2\text{st}}(\chi) = \frac{\alpha^2 m^2 \theta(\chi)}{2p^0 \chi} \xi L_\varphi \int_0^\infty du \int_4^\infty dv \frac{\phi(0)}{u(1+u)v^{3/2} \sqrt{v-4}}, \quad (6.53)$$

Out[212]=

Recall the defitinions

$$\mathbf{p} + \mathbf{s} \mathbf{k} = \mathbf{q} + \mathbf{l}$$

$$\chi_{\mathbf{p}} = \chi_{\mathbf{q}} + \chi_{\mathbf{l}}$$

$$\rho_{\mathbf{l}} = \frac{\mathbf{e}(\mathbf{q} \mathbf{F} \mathbf{p})}{\xi m^4 \chi_{\mathbf{l}}} = - \frac{\mathbf{e}(\mathbf{l} \mathbf{F} \mathbf{p})}{\xi m^4 \chi_{\mathbf{l}}} = \frac{\mathbf{e}(\mathbf{q} \mathbf{F} \mathbf{l})}{\xi m^4 \chi_{\mathbf{l}}}$$

$$\tau_{\mathbf{l}} = - \frac{\mathbf{e}(\mathbf{q} \mathbf{F}^* \mathbf{p})}{m^4 \chi_{\mathbf{l}}} = \frac{\mathbf{e}(\mathbf{l} \mathbf{F}^* \mathbf{p})}{m^4 \chi_{\mathbf{l}}} = - \frac{\mathbf{e}(\mathbf{q} \mathbf{F}^* \mathbf{l})}{m^4 \chi_{\mathbf{l}}}$$

$$\mathbf{k} + \mathbf{l} = \mathbf{f} + \mathbf{g}$$

$$\chi_l = \chi_f + \chi_g$$

$$\rho_2 = -\frac{e(gFf)}{\xi m^4 \chi_l} = -\frac{e(gFl)}{\xi m^4 \chi_l}$$

$$\tau_2 = -\frac{e(gF^* f)}{m^4 \chi_l} = -\frac{e(gF^* l)}{m^4 \chi_l}$$