#### 1

## **Assignment 1**

**AI1110**: Probability and Random Variables Indian Institute of Technology Hyderabad

### Arsh Srivastava AI22BTECH11003

#### Question 12.13.3.14:

If A and B are two events such that  $A \subset B$  and  $P(B) \neq 0$ , then which of the following is correct?

### **Options:**

- 1)  $P(A \mid B) = \frac{P(B)}{P(A)}$
- 2)  $P(A \mid B) < P(A)$
- 3)  $P(A \mid B) \ge P(A)$
- 4) None of these

# Answer: (3) Solution:

As given in the question,

$$A \subset B \Rightarrow AB = A \tag{1}$$

$$\therefore P(A \mid B) = \frac{P(AB)}{P(B)} \tag{2}$$

$$\Rightarrow P(A|B) = \frac{P(A)}{P(B)} \tag{3}$$

$$P(B) \neq 0 \tag{4}$$

$$\Rightarrow 0 < P(B) \le 1 \Rightarrow \frac{1}{P(B)} \ge 1 \tag{5}$$

:. Using equation (1),

$$P(A|B) = P(A) \times \frac{1}{P(B)} \ge P(A) \tag{6}$$

$$\therefore P(A \mid B) \ge P(A)$$

Hence, the correct option is (3).