NSW Education Standards Authority

2020 HIGHER SCHOOL CERTIFICATE EXAMINATION

Mathematics Advanced
Mathematics Extension 1
Mathematics Extension 2

REFERENCE SHEET

Measurement

Length

$$l = \frac{\theta}{360} \times 2\pi r$$

Area

$$A = \frac{\theta}{360} \times \pi r^2$$

$$A = \frac{h}{2}(a + b)$$

Surface area

$$A = 2\pi r^2 + 2\pi rh$$

$$A = 4\pi r^2$$

Volume

$$V = \frac{1}{3}Ah$$

$$V = \frac{4}{3}\pi r^3$$

Functions

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For $ax^3 + bx^2 + cx + d = 0$:

$$\alpha + \beta + \gamma = -\frac{b}{a}$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = \frac{c}{a}$$

and $\alpha\beta\gamma = -\frac{d}{a}$

Relations

$$(x - h)^2 + (y - k)^2 = r^2$$

Financial Mathematics

$$A = P(1 + r)^n$$

Sequences and series

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2}[2a + (n - 1)d] = \frac{n}{2}(a + l)$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(1 - r^n)}{1 - r} = \frac{a(r^n - 1)}{r - 1},\ r \neq 1$$

$$S = \frac{a}{1 - r},\ |r| < 1$$

Logarithmic and Exponential Functions

$$\log_a a^x = x = a^{\log_a x}$$

$$\log_a x = \frac{\log_b x}{\log_b a}$$

$$a^x = e^{x \ln a}$$

Trigonometric Functions

$$\sin A = \frac{\text{opp}}{\text{hyp}},\quad \cos A = \frac{\text{adj}}{\text{hyp}},\quad \tan A = \frac{\text{opp}}{\text{adj}}$$

$$A = \frac{1}{2}ab\sin C$$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$



$$c^2 = a^2 + b^2 - 2ab\cos C$$

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$l = r\theta$$

$$A = \frac{1}{2}r^2\theta$$



Trigonometric identities

$$\sec A = \frac{1}{\cos A},\ \cos A \neq 0$$

$$\operatorname{cosec} A = \frac{1}{\sin A},\ \sin A \neq 0$$

$$\cot A = \frac{\cos A}{\sin A},\ \sin A \neq 0$$

$$\cos^2 x + \sin^2 x = 1$$

Compound angles

$$\sin(A + B) = \sin A\cos B + \cos A\sin B$$

$$\cos(A + B) = \cos A\cos B - \sin A\sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A\tan B}$$

If $t = \tan\frac{A}{2}$ then $\sin A = \frac{2t}{1 + t^2}$

$$\cos A = \frac{1 - t^2}{1 + t^2}$$

$$\tan A = \frac{2t}{1 - t^2}$$

$$\cos A\cos B = \frac{1}{2}[\cos(A - B) + \cos(A + B)]$$

$$\sin A\sin B = \frac{1}{2}[\cos(A - B) - \cos(A + B)]$$

$$\sin A\cos B = \frac{1}{2}[\sin(A + B) + \sin(A - B)]$$

$$\cos A\sin B = \frac{1}{2}[\sin(A + B) - \sin(A - B)]$$

$$\sin^2 nx = \frac{1}{2}(1 - \cos 2nx)$$

$$\cos^2 nx = \frac{1}{2}(1 + \cos 2nx)$$

Statistical Analysis

$$z = \frac{x - \mu}{\sigma}$$

An outlier is a score
less than $Q_1 - 1.5 \times IQR$
or
more than $Q_3 + 1.5 \times IQR$

Normal distribution



- approximately 68% of scores have z-scores between −1 and 1
- approximately 95% of scores have z-scores between −2 and 2
- approximately 99.7% of scores have z-scores between −3 and 3

$$E(X) = \mu$$

$$\operatorname{Var}(X) = E[(X - \mu)^2] = E(X^2) - \mu^2$$

Probability

$$P(A \cap B) = P(A)P(B)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)},\ P(B) \neq 0$$

Continuous random variables

$$P(X \leq r) = \int_a^r f(x)\,dx$$

$$P(a < X < b) = \int_a^b f(x)\,dx$$

Binomial distribution

$$P(X = r) = {}^nC_r p^r (1 - p)^{n-r}$$

$$X \sim \operatorname{Bin}(n, p)$$

$$\Rightarrow\ P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x},\ x = 0, 1, \ldots, n$$

$$E(X) = np$$

$$\operatorname{Var}(X) = np(1 - p)$$

Differential Calculus

Function	Derivative
$y = f(x)^n$	$\frac{dy}{dx} = nf'(x)[f(x)]^{n-1}$
$y = uv$	$\frac{dy}{dx} = u\frac{dv}{dx} + v\frac{du}{dx}$
$y = g(u)$ where $u = f(x)$	$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
$y = \frac{u}{v}$	$\frac{dy}{dx} = \frac{v\frac{du}{dx} - u\frac{dv}{dx}}{v^2}$
$y = \sin f(x)$	$\frac{dy}{dx} = f'(x)\cos f(x)$
$y = \cos f(x)$	$\frac{dy}{dx} = -f'(x)\sin f(x)$
$y = \tan f(x)$	$\frac{dy}{dx} = f'(x)\sec^2 f(x)$
$y = e^{f(x)}$	$\frac{dy}{dx} = f'(x)e^{f(x)}$
$y = \ln f(x)$	$\frac{dy}{dx} = \frac{f'(x)}{f(x)}$
$y = a^{f(x)}$	$\frac{dy}{dx} = (\ln a) f'(x) a^{f(x)}$
$y = \log_a f(x)$	$\frac{dy}{dx} = \frac{f'(x)}{(\ln a) f(x)}$
$y = \sin^{-1} f(x)$	$\frac{dy}{dx} = \frac{f'(x)}{\sqrt{1-[f(x)]^2}}$
$y = \cos^{-1} f(x)$	$\frac{dy}{dx} = -\frac{f'(x)}{\sqrt{1-[f(x)]^2}}$
$y = \tan^{-1} f(x)$	$\frac{dy}{dx} = \frac{f'(x)}{1+[f(x)]^2}$

Integral Calculus

$$\int f'(x)[f(x)]^n dx = \frac{1}{n+1}[f(x)]^{n+1} + c$$

where $n \neq -1$

$$\int f'(x)\sin f(x)\,dx = -\cos f(x) + c$$

$$\int f'(x)\cos f(x)\,dx = \sin f(x) + c$$

$$\int f'(x)\sec^2 f(x)\,dx = \tan f(x) + c$$

$$\int f'(x)e^{f(x)}dx = e^{f(x)} + c$$

$$\int \frac{f'(x)}{f(x)}dx = \ln|f(x)| + c$$

$$\int f'(x)a^{f(x)}dx = \frac{a^{f(x)}}{\ln a} + c$$

$$\int \frac{f'(x)}{\sqrt{a^2 - [f(x)]^2}}dx = \sin^{-1}\frac{f(x)}{a} + c$$

$$\int \frac{f'(x)}{a^2 + [f(x)]^2}dx = \frac{1}{a}\tan^{-1}\frac{f(x)}{a} + c$$

$$\int u\frac{dv}{dx}dx = uv - \int v\frac{du}{dx}dx$$

$$\int_a^b f(x)\,dx$$

$$\approx \frac{b-a}{2n}\left\{f(a) + f(b) + 2\left[f(x_1) + \cdots + f(x_{n-1})\right]\right\}$$

where $a = x_0$ and $b = x_n$

Combinatorics

$${}^nP_r = \frac{n!}{(n-r)!}$$

$$\binom{n}{r} = {}^nC_r = \frac{n!}{r!\,(n-r)!}$$

$$(x+a)^n = x^n + \binom{n}{1}x^{n-1}a + \cdots + \binom{n}{r}x^{n-r}a^r + \cdots + a^n$$

Vectors

$$|\underset{\sim}{u}| = |x\underset{\sim}{i} + y\underset{\sim}{j}| = \sqrt{x^2 + y^2}$$

$$\underset{\sim}{u} \cdot \underset{\sim}{v} = |\underset{\sim}{u}|\,|\underset{\sim}{v}|\cos\theta = x_1x_2 + y_1y_2,$$

where $\underset{\sim}{u} = x_1\underset{\sim}{i} + y_1\underset{\sim}{j}$

and $\underset{\sim}{v} = x_2\underset{\sim}{i} + y_2\underset{\sim}{j}$

$$\underset{\sim}{r} = \underset{\sim}{a} + \lambda\underset{\sim}{b}$$

Complex Numbers

$$z = a + ib = r(\cos\theta + i\sin\theta)$$
$$= re^{i\theta}$$

$$[r(\cos\theta + i\sin\theta)]^n = r^n(\cos n\theta + i\sin n\theta)$$
$$= r^n e^{in\theta}$$

Mechanics

$$\frac{d^2x}{dt^2} = \frac{dv}{dt} = v\frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$$

$$x = a\cos(nt + \alpha) + c$$

$$x = a\sin(nt + \alpha) + c$$

$$\ddot{x} = -n^2(x - c)$$

© NSW Education Standards Authority