

[1] "Plasma"

Exact binomial test

data: `sum(df.good$Is.Latent)` and `length(df.good$Is.Latent)`

number of successes = 83, number of trials = 183, p-value = 0.9084

alternative hypothesis: true probability of success is greater than 0.5

95 percent confidence interval:

0.3911924 1.0000000

sample estimates:

probability of success

0.4535519

Generalized linear mixed model fit by maximum likelihood (Laplace
Approximation) [glmerMod]
Family: binomial (logit)
Formula: ls.Latent ~ (1 | Patient)
Data: df.good

AIC	BIC	logLik	deviance	df.resid
256.1	262.5	-126.1	252.1	181

Scaled residuals:

Min	1Q	Median	3Q	Max
-0.911	-0.911	-0.911	1.098	1.098

Random effects:

Groups	Name	Variance	Std.Dev.
Patient	(Intercept)	1.741e-14	1.319e-07

Number of obs: 183, groups: Patient, 7

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-0.1863	0.1485	-1.255	0.21


[1] "AIC: 256.110353848107, null AIC: 254.110353848107"

[2] "p-value: 1"


[3] "mean: 0.453551912568308"

EXPLORATORY DATA ANALYSIS


histogram of resid(bin.glme.test) Density of resid(bin.glme.test)




Boxplot of resid(bin.glme.test) Q-Q Plot of resid(bin.glme.test)




2658




13333




13334




7265




13336



35566



825



Overall

