

[1] "Mixed"

Exact binomial test

data: `sum(df.good$Is.Latent)` and `length(df.good$Is.Latent)`

number of successes = 746, number of trials = 1317, p-value = $7.88e-07$

alternative hypothesis: true probability of success is greater than 0.5

95 percent confidence interval:

0.5435041 1.0000000

sample estimates:

probability of success

0.5664389

Generalized linear mixed model fit by maximum likelihood (Laplace
Approximation) [glmerMod]
Family: binomial (logit)
Formula: ls.Latent ~ (1 | Patient)
Data: df.good

AIC	BIC	logLik	deviance	df.resid
1619.2	1629.5	-807.6	1615.2	1315

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.0304	-0.8924	0.4150	0.8101	2.0604

Random effects:

Groups	Name	Variance	Std.Dev.
Patient	(Intercept)	1.228	1.108

Number of obs: 1317, groups: Patient, 14

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	0.3217	0.3047	1.056	0.291


[1] "AIC: 1619.16129557413, null AIC: 1804.42714946063"

[2] "p-value: 0"


[3] "mean: 0.57974432304816"

EXPLORATORY DATA ANALYSIS


histogram of resid(bin.glme.test) Density of resid(bin.glme.test)




Boxplot of resid(bin.glme.test) Q-Q Plot of resid(bin.glme.test)




10138




13889




16616




16617




16618




16619




34393




34399




34410




34411




820




821



822



824



Overall

