

[1] ""

## Exact binomial test

data: `sum(df.good$Is.Latent)` and `length(df.good$Is.Latent)`

number of successes = 47, number of trials = 101, p-value = 0.7869

alternative hypothesis: true probability of success is greater than 0.5

95 percent confidence interval:

0.3803851 1.0000000

sample estimates:

probability of success

0.4653465

Generalized linear mixed model fit by maximum likelihood (Laplace  
Approximation) [glmerMod]  
Family: binomial ( logit )  
Formula: ls.Latent ~ (1 | Patient)  
Data: df.good

AIC	BIC	logLik	deviance	df.resid
137.4	142.6	-66.7	133.4	99

Scaled residuals:

Min	1Q	Median	3Q	Max
-1.2434	-0.8512	-0.5408	0.8042	1.8491

Random effects:

Groups	Name	Variance	Std.Dev.
Patient	(Intercept)	0.5127	0.716

Number of obs: 101, groups: Patient, 4

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z )
(Intercept)	-0.3311	0.4260	-0.777	0.437


[1] "AIC: 137.385953638275, null AIC: 141.530192812247"

[2] "p-value: 0.0131840875879025"


[3] "mean: 0.417969444746623"

# EXPLORATORY DATA ANALYSIS


histogram of resid(bin.glme.test) Density of resid(bin.glme.test)




Boxplot of resid(bin.glme.test) Q-Q Plot of resid(bin.glme.test)




2658



13333




13334





13336



# Overall

