

Exact binomial test

data: `sum(df.good$Is.Latent)` and `length(df.good$Is.Latent)`

number of successes = 104, number of trials = 190, p-value = 0.1087

alternative hypothesis: true probability of success is greater than 0.5

95 percent confidence interval:

0.4850982 1.0000000

sample estimates:

probability of success

0.5473684

Generalized linear mixed model fit by maximum likelihood (Laplace
Approximation) [glmerMod]
Family: binomial (logit)
Formula: ls.Latent ~ (1 | Patient)
Data: df.good

AIC	BIC	logLik	deviance	df.resid
265.7	272.2	-130.8	261.7	188

Scaled residuals:

Min	1Q	Median	3Q	Max
-1.0997	-1.0997	0.9093	0.9093	0.9093

Random effects:

Groups	Name	Variance	Std.Dev.
	Patient (Intercept)	0	0

Number of obs: 190, groups: Patient, 8

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	0.1900	0.1458	1.304	0.192


[1] "AIC: 265.688105443976 null AIC: 263.688105443976"

[2] "p-value: 0.999999999999886"


[3] "mean: 0.547368421052514 (0.476111253958456, 0.616735524503627)"

EXPLORATORY DATA ANALYSIS


histogram of resid(bin.glme.test) Density of resid(bin.glme.test)




Boxplot of resid(bin.glme.test) Q-Q Plot of resid(bin.glme.test)




2658




13333




13334




7265




1336




7263



35566



825



Overall

