

# Multi-isotope generator calculation

Artem Lebedev<sup>1\*</sup>

<sup>1\*</sup>Department of Chemistry and Chemical Biology, McMaster  
University, Hamilton, ON, Canada.

Corresponding author(s). E-mail(s): [artem@indiechemistry.com](mailto:artem@indiechemistry.com);

## Abstract

A brief explanation of the math behind solving arbitrarily complex isotope generator.

## 1 Introduction

Isotope generators are the backbone of medical isotope production, accounting for well over 90% of all the nuclear medicine procedures. The bulk of this volume comes from the use of Mo-99/Tc-99m generator, but other parent-daughter pairs have been gaining popularity recently, namely Ge-68/Ga-68, Ra-225/Ac-225 and a few others. Theoretical calculations of the generator state and predictions of the generator output rely on classic Bateman equations[1]. These equations, derived in 1908, allow to predict the present state of the generator, that is the concentration of all isotopes, given the initial state of the generator. While this is technically sufficient, practical considerations of harvesting schedules, generator shelf-life and economical viability add a new layer of complexity. Additionally, computational implementations of Bateman equations are known to suffer from a series of drawbacks [2]. Over the years, a few approaches to solving the Reutherford's system of equations have been proposed [3]

This paper describes the logic behind a computer code designed to simulate generator states and output under various operational conditions.

## 2 Two solutions to the same equations

Let's consider the most important isotope generator in nuclear medicine: Tc generator, that is still a staple of the clinical practice, used for millions upon millions of SPECT scans every year, all around the world. Mo-99 decays two ways: ~87.6% to Tc-99m

(branching ratio  $b$ ) and  $\sim 12.4\%$  directly to (virtually) stable Tc-99, which does not factor into the generator considerations. From here, a conventional chemist's thinking can easily lead us to the following system of equations:

$$\begin{cases} \frac{d[{}^{99}\text{Mo}]}{dt} = -\lambda_{\text{Mo}}[{}^{99}\text{Mo}] \\ \frac{d[{}^{99m}\text{Tc}]}{dt} = b\lambda_{\text{Mo}}[{}^{99}\text{Mo}] - \lambda_{\text{Tc}}[{}^{99m}\text{Tc}] \end{cases} \quad (1)$$

where decay constants are  $\lambda_{\text{Mo}} = \ln 2/65.94h$ ,  $\lambda_{\text{Tc}} = \ln 2/6.01h$  and branching ratio  $b = 0.876$

What this is saying is that the rate of accumulation of  ${}^{99}\text{Mo}$  is negatively proportional to its concentration, i.e. the more  ${}^{99}\text{Mo}$  we have in solution the slower it accumulates, or what is the same, the faster it decays. This is essentially a definition of the radioactive decay. For the  ${}^{99m}\text{Tc}$  the picture is more complex, but is still intuitive: its rate of accumulation is *negatively* proportional to its own concentration  $[{}^{99m}\text{Tc}]$ , but also *positively* proportional to the parent's concentration  $[{}^{99}\text{Mo}]$ . Also, let's not forget that the  $[{}^{99}\text{Mo}]$  is scaled to the branching ratio in the second equation. Of course, the question that concerns the chemists the most is "What is in the beaker right now?", and the quest to answering this question can lead us down two surprisingly divergent paths. The more intuitive one was explored by Bateman and bears its name, while the second one relies on matrix exponential.

## 2.1 Bateman equations and their shortcomings

In 1910 Bateman proposed an analytical solution to arbitrarily long sequential decay chain

## 2.2 Matrix exponential: break your brains

Now notice that system (1) can be re-written in matrix form. Just a simple change in notation, nothing more:

$$\frac{d}{dt} \begin{bmatrix} [{}^{99}\text{Mo}] \\ [{}^{99m}\text{Tc}] \end{bmatrix} = \begin{pmatrix} -\lambda_{\text{Mo}} & 0 \\ b\lambda_{\text{Mo}} & -\lambda_{\text{Tc}} \end{pmatrix} \begin{bmatrix} [{}^{99}\text{Mo}] \\ [{}^{99m}\text{Tc}] \end{bmatrix} \quad (2)$$

Only the subdiagonal entry picks up the factor  $b$ . Diagonalizing exactly as before (eigenvalues still  $-\lambda_1, -\lambda_2$  since those live on the diagonal regardless of  $b$ ):

$$\begin{aligned} \mathbf{e}^{\mathbf{A}t} &= \begin{pmatrix} e^{-\lambda_1 t} & 0 \\ \frac{b\lambda_1}{\lambda_2 - \lambda_1} \left( e^{-\lambda_1 t} - e^{-\lambda_2 t} \right) & e^{-\lambda_2 t} \end{pmatrix} \end{aligned}$$

So Bateman becomes:

$$\mathbf{N}(t) = \mathbf{N}_0 \begin{pmatrix} e^{-\lambda_1 t} & 0 \\ \frac{b\lambda_1}{\lambda_2 - \lambda_1} \left( e^{-\lambda_1 t} - e^{-\lambda_2 t} \right) & e^{-\lambda_2 t} \end{pmatrix}$$

— literally just the un-branched formula times  $b$ . This makes sense: branching ratio only ever multiplies the *\*production\** term, never the decay-loss term, so it

factors out linearly and leaves the transient/secular equilibrium math (the classic 6-hour Mo-99/Tc-99m generator “milking” curve) otherwise unchanged.

**\*\*Where it stops being this simple\*\***

If you extend the chain further (e.g. tracking Tc-99 ground state buildup too, or handling isotopes with 3+ branches like your Ac-225 chain has at Bi-213  $\rightarrow$  Tl-209/Po-213), you get multiple daughters fed from one parent. That’s not a single chain anymore —  $A$  becomes a genuine tree structure, and you’d want one column feeding multiple rows (each weighted by its own  $b_i$ , with  $\sum b_i = 1$ ) rather than a simple bidiagonal matrix. The matrix exponential machinery still works fine, it’s just no longer tridiagonal/bidiagonal, so ‘expm’ is basically mandatory rather than a convenience at that point.

**Acknowledgements.** Author thanks Ian Horwath for a stimulating discussion and Andrew Knight for the feedback on the web app version of these computations.

## Declarations

- Funding: The work was entirely self-funded
- Conflict of interest: The author is an employee of BWXT Medical, Ltd.
- Data availability: The article and the app code are available on GitHub

## References

- [1] Bateman, H.: The solution of a system of differential equations occurring in the theory of radio-active decay **15**, 423–427
- [2] Bakin, R.I., Kiselev, A.A., Shvedov, A.M., Shikin, A.V.: Computational errors in the calculation of long radioactive decay chains **123**(6), 406–411 <https://doi.org/10.1007/s10512-018-0360-2> . Accessed 2026-07-19
- [3] Maren, V.: Implementation, Validation and Comparison of Different Algorithms to Solve the Bateman Equations for Very Large Systems