

A Programmer's Introduction to Mathematics:

Chapter 2 Exercise solutions

Arthur Allshire

December 10, 2018

2.6: Prove Vieta's Formulas

Exercise. Let $f(x) = a_0 + a_1x + \dots + a_nx^n$ be a degree n polynomial, and suppose it has k real roots r_1, \dots, r_n . Prove Vieta's Formulas, which are

$$\sum_{i=1}^n r_i = -\frac{a_{n-1}}{a_n}$$

$$\prod_{i=1}^n r_i = (-1)^n \frac{a_0}{a_n}.$$

Proof. Let $g(x) = \frac{f(x)}{a_n} = \frac{a_0}{a_n} + \frac{a_1}{a_n}x + \dots + x^n$. Now, given roots of $g(x)$ are r_1, \dots, r_n , the monic polynomial

$$g(x) = \prod_{i=1}^n (x - r_i) = \sum_{i=0}^n \frac{a_i}{a_n} x^i$$

Now, equating the coefficients of x^{n-1} on both sides (observe the expansion of the product on the left hand side resulting in the multiplication of r_i by x^{n-1} times), we get

$$\sum_{i=1}^n r_i = -\frac{a_{n-1}}{a_n},$$

as required. Similarly, we simply observe equate the constant term (the "coefficient of x^0 ") on both sides, observing that on the left hand side this is the product of the roots multiplied by $(-1)^n$, and divide through by $(-1)^n$ to obtain the second equation.

□