

Efficacy and Safety From the Phase 1/2 MonumenTAL-1 Study of Talquetamab, a GPRC5D×CD3 Bispecific Antibody, in Patients With Relapsed/Refractory Multiple Myeloma: Analyses at an Extended Median Follow-Up

Leo Rasche¹, Carolina Schinke², Cyrille Touzeau³, Monique C Minnema⁴, Niels WCJ van de Donk^{5,*} (n.vandedonk@amsterdamumc.nl), Paula Rodríguez-Otero⁶, María-Victoria Mateos⁷, Jing Christine Ye⁸, Chalmer Tomlinson⁹, Deeksha Vishwamitra¹⁰, Indrajeet Singh¹⁰, Xiang Qin¹⁰, Michela Campagna¹¹, Tara J Masterson¹⁰, Veronique Vreys¹², Bonnie W Lau¹⁰, Jaszianna Tolbert¹⁰, Thomas Renaud⁹, Christoph Heuck¹⁰, Ajai Chari¹³

*Presenting author.
¹University Hospital of Würzburg, Würzburg, Germany; ²Myeloma Center, University of Arkansas for Medical Sciences, Little Rock, AR, USA; ³Centre Hospitalier Universitaire de Nantes, Nantes, France; ⁴University Medical Center Utrecht, Utrecht, Netherlands; ⁵Amsterdam University Medical Center, Vrije Universiteit Amsterdam, Amsterdam, Netherlands; ⁶Clinica Universidad de Navarra, CIMA, CIBERONC, IDISNA, Pamplona, Spain; ⁷University Hospital of Salamanca/IBSAL/CIC/CIBERONC, Salamanca, Spain; ⁸MD Anderson Cancer Center, University of Texas, Houston, TX, USA; ⁹Johnson & Johnson, Raritan, NJ, USA; ¹⁰Johnson & Johnson, Spring House, PA, USA; ¹¹Johnson & Johnson, Madrid, Spain; ¹²Johnson & Johnson, Beerse, Belgium; ¹³University of California, San Francisco, CA, USA


Key Takeaway

At an extended mFU (30–38 months), patients with RRMM treated with talquetamab continue to demonstrate deep and durable responses and tolerable safety with no new discontinuations due to GPRC5D-related AEs

Conclusions

High ORRs elicited by talquetamab were durable and led to promising 36-month OS rates (45–61%)

The safety profile continued to show lower risk of high-grade infections relative to approved anti-BCMA BsAbs,^{8,9} potentially contributing to the OS benefit with talquetamab and highlighting the humoral immune preservation that enables versatile use of talquetamab including in combination



Please scan QR code

Supplementary material

<https://www.congresshub.com/Oncology/AM2025/Talquetamab/Donk-Talq>


Copies of this presentation obtained through Quick Response (QR) Codes are for personal use only and may not be reproduced without permission from ASCO® or the author of this presentation.

Acknowledgments
This study was funded by Johnson & Johnson. Medical writing support was provided by Rachael Smith, PhD, of Eloquent Scientific Solutions, and funded by Johnson & Johnson. We thank the patients who participated in the study and their families and caregivers, the physicians and nurses who cared for patients and supported this clinical trial, staff members at the study sites, and staff members involved in data collection and analyses.

Disclosures
NWCJvdD has received research funding from Amgen, BMS, Celgene, Cellectis, Johnson & Johnson, and Novartis; and reports a consulting or advisory role with AbbVie, Adaptive Biotechnologies, Amgen, Bayer, BMS, Celgene, Galapagos, Johnson & Johnson, Novartis, Pfizer, Roche, Servier, and Takeda.

Introduction

- Talquetamab is the first and only approved bispecific antibody (BsAb) targeting G protein–coupled receptor class C group 5 member D (GPRC5D) for the treatment of relapsed/refractory multiple myeloma (RRMM)¹⁻³
- In previous results from the phase 1/2 MonumenTAL-1 study (clinical cut-off: Jan 2024; median follow-up [mFU], 21–30 months), talquetamab elicited deep, durable responses with low discontinuation rates⁴



We report efficacy and ongoing safety from MonumenTAL-1 at an extended mFU of 30–38 months, the longest mFU for any anti-GPRC5D agent

Results





Table. mDOR and mPFS continued to demonstrate superior outcomes in the Q2W vs QW cohort

Outcome	QW (n=143)	Q2W (n=154)	Prior TCR QW and Q2W (n=78)
mFU, mo	38.2	31.2	30.3
mDOR, mo (95% CI) ^a	9.5 (6.7–13.4)	17.5 (12.5–25.1)	19.2 (8.1–24.7)
mPFS, mo (95% CI)	7.5 (5.7–9.4)	11.2 (7.7–14.6)	7.7 (4.1–14.5)
MRD neg (10 ⁻⁵), % (95% CI) ^b	64.3 (51.9–75.4)	65.2 (52.8–76.3)	57.1 (37.2–75.5)

^an=106 (QW), n=107 (Q2W), n=52 (prior TCR QW and Q2W). ^bAssessed in patients with evaluable samples: n=70 (QW), n=69 (Q2W), n=28 (prior TCR QW and Q2W). Only MRD assessments (10⁻⁵) within 3 months of achieving CR or sCR until death, disease progression, or subsequent therapy are considered. See Supplemental Table 1 for efficacy outcomes in the USPI population (≥4 prior LOT). mDOR, median duration of response; mPFS, median progression-free survival; MRD neg, minimal residual disease negativity; USPI, US prescribing information.





References


- Verkleij CPM, et al. *Blood Adv* 2021;5:2196-215.
- Chari A, et al. *Lancet Haematol* 2025;12:E269-E821.
- TALVEY™ (talquetamab-tgvs). Prescribing information. Horsham, PA: Janssen Biotech, Inc.; 2023.
- Rasche L, et al. Presented at EHA; June 13–16, 2024; Madrid, Spain.
- Rajkumar SV, et al. *Blood* 2011;117:4691-5.
- Kumar S, et al. *Lancet Oncol* 2016;17:328-46.
- Lee DW, et al. *Biol Blood Marrow Transplant* 2019;25:625-38.
- van de Donk NWCJ, et al. Presented at ASCO; June 2–6, 2023; Chicago, IL, USA & Virtual. #8011.
- Tomasson M, et al. *Blood* 2023;142(supplement 1):3385.

Methods

MonumenTAL-1^a phase 1/2 study design



NCT03399799/NCT04634552. ^aWith 2–3 step-up doses. ^bPatients received QW (n=70) or Q2W (n=8) dosing. ^cAssessed by IRC using International Myeloma Working Group criteria. ^dMRD was assessed using bone marrow aspirates and evaluated via next-generation sequencing. CRS and ICANS were graded by ASTCT criteria¹; all other AEs were graded by CTCAE v4.03. ^eAdverse event; ASTCT, American Society of Transplantation and Cellular Therapy; CR, complete response; CRS, cytokine release syndrome; CTCAE, Common Terminology Criteria for Adverse Events; DOR, duration of response; ECOG PS, Eastern Cooperative Oncology Group performance status; ICANS, immune effector cell–associated neurotoxicity syndrome; IMiD, immunomodulatory drug; IRC, independent review committee; LOT, line of therapy; mAb, monoclonal antibody; MRD, minimal residual disease; ORR, overall response rate; OS, overall survival; PFS, progression-free survival; PI, proteasome inhibitor; Q2W, every other week; QW, weekly; SC, subcutaneous; TCR, T-cell redirection therapy; VGPR, very good partial response.





The safety profile was consistent with previous results (Supplemental Table 2)⁴; discontinuation rates due to AEs remained low, and no new discontinuations occurred due to GPRC5D-related AEs

A new safety signal, ataxia/balance disorders, was recently identified in association with talquetamab and had low prevalence in MonumenTAL-1



Multiple Myeloma



Supplemental Table 1: Efficacy Outcomes in the USPI Population

Outcome	QW (n=100)	Q2W (n=90)	Prior TCR QW and Q2W ^a (n=58)
ORR, %	73.0	71.1	72.4
≥CR	35.0	43.3	50.0
VGPR	22.0	17.8	8.6
PR	16.0	10.0	13.8
Median time to best response of ≥CR, ^b mo (range)	2.27 (1.1–12.7)	6.24 (1.2–16.8)	2.66 (1.2–17.5)
Median time to best response of VGPR, ^c mo (range)	1.97 (1.1–6.2)	3.06 (0.3–18.9)	2.04 (1.2–2.1)
Median time to best response of PR, ^d mo (range)	1.28 (1.1–2.9)	2.07 (1.2–2.8)	1.13 (1.1–3.0)
Median DOR, mo (95% CI) ^e	10.2 (6.6–15.7)	17.9 (12.5–26.0)	19.2 (6.7–NE)
≥CR ^b	28.8 (18.9–NE)	26.1 (18.0–NE)	24.7 (19.2–NE)
VGPR ^c	6.4 (4.4–9.5)	9.3 (7.4–15.2)	4.8 (2.1–NE)
PR ^d	3.0 (1.9–5.6)	5.5 (0.9–6.5)	2.4 (1.9–4.6)
Median PFS (95% CI), mo	6.8 (5.5–10.4)	12.4 (9.6–18.2)	11.3 (4.8–21.4)
36-mo PFS, %	17.6 (10.7–26.0)	NE (NE–NE)	28.2 (16.0–41.7)
Median OS (95% CI), mo	NR (21.7–NE)	NR (33.2–NE)	30.6 (20.2–NE)
36-mo OS, %	50.5 (40.0–60.0)	NE (NE–NE)	46.4 (29.2–61.9)

Data are reported from phase 2 only.
^aPhase 2 data include the 0.4 mg/kg QW cohort only. ^bn=35 (QW), n=39 (Q2W), n=29 (prior TCR QW and Q2W). ^cn=22 (QW), n=16 (Q2W), n=5 (prior TCR QW and Q2W). ^dn=16 (QW), n=9 (Q2W), n=8 (prior TCR). ^en=73 (QW), n=64 (Q2W), n=42 (prior TCR QW and Q2W).
CR, complete response; DOR, duration of response; NE, not estimable; NR, not reached; ORR, overall response rate; OS, overall survival; PFS, progression-free survival; PR, partial response; Q2W, every other week; QW, weekly; TCR, T-cell redirection therapy; USPI, US prescribing information; VGPR, very good partial response.

Supplemental Table 2: Hematologic and Nonhematologic AEs

AE (≥30% in any cohort), n (%)	QW (n=143)		Q2W (n=154)		Prior TCR QW and Q2W (n=78)	
	Any Grade	Grade 3/4	Any Grade	Grade 3/4	Any Grade	Grade 3/4
Hematologic AE						
Anemia	65 (45.5)	46 (32.2)	67 (43.5)	39 (25.3)	38 (48.7)	22 (28.2)
Neutropenia	50 (35.0)	44 (30.8)	44 (28.6)	33 (21.4)	40 (51.3)	37 (47.4)
Thrombocytopenia	39 (27.3)	29 (20.3)	46 (29.9)	28 (18.2)	30 (38.5)	22 (28.2)
Nonhematologic AE						
CRS	113 (79.0)	3 (2.1)	116 (75.3)	1 (0.6)	57 (73.1)	1 (1.3)
Dysgeusia ^a	103 (72.0)	NA	111 (72.1)	NA	59 (75.6)	NA
Infections ^b	87 (60.8)	33 (23.1)	109 (70.8)	33 (21.4)	61 (78.2)	20 (25.6)
Skin related ^c	85 (59.4)	0	113 (73.4)	1 (0.6)	53 (67.9)	0
Nail related ^d	80 (55.9)	0	84 (54.5)	0	47 (60.3)	0
Weight decreased	59 (41.3)	3 (2.1)	64 (41.6)	9 (5.8)	29 (37.2)	1 (1.3)
Rash related ^e	57 (39.9)	2 (1.4)	48 (31.2)	8 (5.2)	25 (32.1)	2 (2.6)
Pyrexia	57 (39.9)	4 (2.8)	44 (28.6)	2 (1.3)	27 (34.6)	0
Dry mouth	38 (26.6)	0	60 (39.0)	0	34 (43.6)	0
Fatigue	36 (25.2)	5 (3.5)	44 (28.6)	1 (0.6)	25 (32.1)	1 (1.3)

^aIncluding ageusia, dysgeusia, hypogeusia, and taste disorder. Per CTCAE, the maximum possible grade of dysgeusia is 2. ^bInfections are reported as a System Organ Class. ^cIncluding skin exfoliation, dry skin, pruritus, and palmar-plantar erythrodysesthesia syndrome. ^dIncluding nail discoloration, nail disorder, onycholysis, onychomadesis, onychoclasia, nail dystrophy, nail toxicity, and nail ridging. ^eIncluding rash, maculopapular rash, erythematous rash, and erythema. AE, adverse event; CRS, cytokine release syndrome; CTCAE, Common Terminology Criteria for Adverse Events; NA, not applicable; Q2W, every other week; QW, weekly, TCR, T-cell redirection therapy.