# 01. Self-assessment questions

January 2, 2021

- Look at the scatterplot below[1]

Can you clearly see if there is a dependence between Population and Area? Are there any transformations that can help to unveil the possible dependency? If yes, would you apply such transformation on the x-axis, y-axis or both?

- Look at the two plots below [2] [3]

---

[1]Attribution Skbkekas, CC BY-SA 3.0 <http://creativecommons.org/licenses/by-sa/3.0/>, via Wikimedia Commons

[2]Attribution: Creative Commons Attribution-ShareAlike 2.5 , `https://en.wikipedia.org/wiki/File:Lightness_approximations.svg`

[3]Attribution: Krishnavedala, CC BY-SA 3.0 <https://creativecommons.org/licenses/by-sa/3.0>, via Wikimedia Commons

For each graph, tell if there is a dependence between the x and y variables. If yes, would this result in a null, positive or negative Pearson's correlation coefficient?

- Assume one of your features has 4 values, you can encode them in two ways: (i) Transform them in 0,1,2,3 or (ii) Apply One-Hot Encoding. When would you choose (i) and when (ii)?

- Suppose you have a dataset with 200 columns. One column has 0.1% of missing values. The other column has 30% of missing values. What pre-processing would you apply before applying any supervised or unsupervised machine learning methods? [See the code related to pre-processing]

- Suppose you have a column with the following values: very bad, bad, medium, good, very good, excellent, no-opinion. How would you transform it in numerical form?

- Suppose you have a column with the following values: student, medical doctor, engineer, cashier, no-info-available. How would you transform such column in numerical form?

- Write the formulas of variance (of a feature), standard deviation, Pearson's correlation coefficient.

- Explain how you construct a boxplot and what is the meaning of all the lines that compose it.

- Explain how to calculate the 35-th percentile.

.