

Contents

1	Introduction to Topology	2
2	Bases for Topologies	7
3	Closed sets, closures, and dense sets	13

1 Introduction to Topology

Problem 1.1

Fix $a < b \in \mathbb{R}$. Show explicitly that the interval (a, b) is open in $\mathbb{R}_{\text{usual}}$. Show explicitly that the interval $[a, b)$ is not open in $\mathbb{R}_{\text{usual}}$.

Solution: Let $x \in (a, b)$. Take $\epsilon = \min(x - a, b - x)$. Let $y \in B_\epsilon(x)$. If $y = x$, then $y \in (a, b)$. If $y < x < b$, we have that $x - y = |x - y| < \epsilon \implies x - y < x - a \implies y > a$. Hence $y \in (a, b)$ in this case. If $a < x < y$, we have that $y - x = |x - y| < \epsilon \implies y - x < b - x \implies y < b$. Hence $y \in (a, b)$ in this case. Thus, $B_\epsilon(x) \subseteq (a, b)$ proving that (a, b) is open in $\mathbb{R}_{\text{usual}}$. \square

To show that $[a, b)$ is not open, consider an arbitrary open ball centred at $a \in [a, b)$. Note that for all $\epsilon > 0$, $a - \frac{\epsilon}{2} \in B_\epsilon(a)$ but $a - \frac{\epsilon}{2} \notin [a, b)$. Hence every open ball centered at a is not contained in $[a, b)$ which means that $[a, b)$ is not open in $\mathbb{R}_{\text{usual}}$. \square

Problem 1.2

Let X be a set and let $\mathcal{B} = \{\{x\} : x \in X\}$. Show that the only topology on X that contains \mathcal{B} as a subset is the discrete topology.

Solution: Let \mathcal{T} be an arbitrary topology such that $\mathcal{B} \subseteq \mathcal{T}$. I claim that $\mathcal{T} = \mathcal{P}(X)$, which by definition is the discrete topology. The forward direction follows by the definition of a topology since a topology is a collection of subsets. For the converse direction, let $A \subseteq X$. If $A = \emptyset$, $A \in \mathcal{T}$ since we know that \mathcal{T} is a topology. Otherwise, we can then write $A = \bigcap_{a \in A} \{a\}$ and hence A is a union of elements of \mathcal{B} and since $\mathcal{B} \subseteq \mathcal{T}$, by the definition of a topology, $A \in \mathcal{T}$. Hence $\mathcal{T} = \mathcal{P}(X)$ as desired. \square

Problem 1.3

Fix a set X , and let $\mathcal{T}_{\text{co-finite}}$ and $\mathcal{T}_{\text{co-countable}}$ be the co-finite and co-countable topologies on X , respectively.

- Show explicitly that $\mathcal{T}_{\text{co-finite}}$ and $\mathcal{T}_{\text{co-countable}}$ are both topologies on X .
- Show that $\mathcal{T}_{\text{co-finite}} \subseteq \mathcal{T}_{\text{co-countable}}$.
- Under what circumstances does $\mathcal{T}_{\text{co-finite}} = \mathcal{T}_{\text{co-countable}}$?
- Under what circumstances does $\mathcal{T}_{\text{discrete}} = \mathcal{T}_{\text{co-countable}}$?

Solution:

- For co-finite, we first note that $\emptyset \in \mathcal{T}_{\text{co-finite}}$ by definition and $X \in \mathcal{T}_{\text{co-finite}}$ since $X \setminus X = \emptyset$ which is finite. Let $U_1, U_2 \in \mathcal{T}_{\text{co-finite}}$. If either are empty, the intersection is empty and thus remains in $\mathcal{T}_{\text{co-finite}}$. Otherwise, say $U_1 = X \setminus \{x_1, \dots, x_m\}$ and $U_2 = X \setminus \{y_1, \dots, y_n\}$. Then $U_1 \cap U_2 = X \setminus (\{x_1, \dots, x_m\} \cup \{y_1, \dots, y_n\})$ and hence is still in $\mathcal{T}_{\text{co-finite}}$. Now consider a collection $\{U_\lambda\}_{\lambda \in \Lambda}$ of open sets in $\mathcal{T}_{\text{co-finite}}$. We can assume they are all non-empty since this will not change the value of

the union. So each U_λ is of the form $X \setminus V_\lambda$ where V_λ is finite. We have that:

$$\begin{aligned}\bigcup_{\lambda \in \Lambda} U_\lambda &= \bigcup_{\lambda \in \Lambda} X \setminus V_\lambda \\ &= X \setminus \left(\bigcap_{\lambda \in \Lambda} V_\lambda \right)\end{aligned}$$

Note that the arbitrary intersection of finite sets remains finite (this is seen because the arbitrary intersection is a subset of any one of those finite sets) and hence, $\bigcup_{\lambda \in \Lambda} U_\lambda$ remains open. Hence the cofinite topology is indeed a topology. \square

For the cocountable topology, the proof is the same. Since the union of 2 countable sets is still countable, finite intersection of open sets in the cocountable topology is open. Since arbitrary intersections of countable sets is countable, arbitrary unions remain open. Hence the cocountable topology is a topology. \square

- b) This just follows from the fact that all finite sets are countable. So if $X \setminus U$ is finite (i.e $U \in \mathcal{T}_{\text{co-finite}}$), then $X \setminus U$ is countable and thus $U \in \mathcal{T}_{\text{co-countable}}$. \square
- c) This occurs if and only if X is finite. For the converse direction, we see that $\mathcal{T}_{\text{co-finite}} = \mathcal{T}_{\text{co-countable}} = \mathcal{P}(X)$ because every subset of X is finite and countable. For the forward direction, consider the contrapositive. Let X be infinite and thus there exists a countable infinite (denumerable) subset C in X . Note that $X \setminus C$ is thus an element of the cocountable topology but not the cofinite topology. \square
- d) This occurs if and only if X is countable. For the converse direction, every subset is countable, and thus every subset of X has its complement countable, thus $\mathcal{T}_{\text{co-countable}} = \mathcal{P}(X) = \mathcal{T}_{\text{discrete}}$. For the forward direction, consider the contrapositive. Suppose X is uncountable and consider any $x \in X$. Note that $X \setminus \{x\}$ remains uncountable, and so $\{x\} \in \mathcal{T}_{\text{discrete}}$ but $\{x\} \notin \mathcal{T}_{\text{co-countable}}$ and hence $\mathcal{T}_{\text{co-countable}} \neq \mathcal{T}_{\text{discrete}}$ as desired. \square

Problem 1.4

Let $(X, \mathcal{T}_{\text{co-countable}})$ be an infinite set with the co-countable topology. Show that $\mathcal{T}_{\text{co-countable}}$ is closed under countable intersections. Give an example to show that it need not be closed under arbitrary intersections.

Solution: Let $\{U_n\}_{n \in \mathbb{N}}$ be countably many sets in the co-countable topology. We can assume they are all non-empty since this will not change the value of the union. Thus they are of the form $U_n = X \setminus V_n$ where V_n is countable for all $n \in \mathbb{N}$. Then, we simply note that:

$$\begin{aligned}\bigcap_{n \in \mathbb{N}} U_n &= \bigcap_{n \in \mathbb{N}} X \setminus V_n \\ &= X \setminus \left(\bigcup_{n \in \mathbb{N}} V_n \right)\end{aligned}$$

Note that the countable union of countable sets remain countable so the countable intersection above is still in the cocountable topology. \square

To show that arbitrary intersections need not remain in the cocountable topology, consider \mathbb{R} equipped with the cocountable topology. For $x \in \mathbb{R}$ with $x \geq 0$, define $U_x = \mathbb{R} \setminus \{x\}$ which is an element of the cocountable topology. Then consider the arbitrary intersection of these sets. We see that $\bigcap_{x \geq 0} U_x = \mathbb{R}_{<0}$. Note that

$\mathbb{R}_{<0}$ is not an element of the cocountable topology since its complement is the non negative real numbers which is not countable.

Problem 1.5

Let X be a nonempty set, and fix an element $p \in X$. Recall that

$$\mathcal{T}_p = \{U \subseteq X : p \in U\} \cup \{\emptyset\}$$

is called the point topology at p on X . Show that \mathcal{T}_p is a topology on X .

Solution: We have $p \in X$ so $X \in \mathcal{T}_p$ and that $\emptyset \in \mathcal{T}_p$. For arbitrary unions, if $p \in U_\lambda$ for all $\lambda \in \Lambda$, then $p \in \bigcup U_\lambda$. Similarly, if $p \in U_1$ and $p \in U_2$, then $p \in U_1 \cap U_2$. Hence this is a topology. \square

Problem 1.6

Define the ray topology on \mathbb{R} as:

$$\mathcal{T}_{\text{ray}} = \{(a, \infty) : a \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$$

Show that \mathcal{T}_{ray} is a topology on \mathbb{R} . Be sure to think carefully about unions.

Solution: By definition, $\emptyset, \mathbb{R} \in \mathcal{T}_{\text{ray}}$. For intersections, if the empty set is contained in the intersection, the intersection is the empty set. If \mathbb{R} is contained in the intersection of 2 open sets, the other set (which is open) will be the result and thus the intersection is open. Otherwise, consider $(a, \infty) \cap (b, \infty)$. Simply note that this is equal to $(\max(a, b), \infty)$ because $x > a$ and $x > b$ if and only if $x > \max(a, b)$. For unions, if \mathbb{R} is contained in the union the result is \mathbb{R} which is open. We can assume there are no empty sets in the union as that will not change the result. Now consider an arbitrary collection of intervals (a_λ, ∞) , and let $S = \{a_\lambda : \lambda \in \Lambda\}$. We consider 2 cases, S is bounded below and S is not bounded below.

In the first case, by the completeness of \mathbb{R} , we know that $\inf S$ exists, and I claim that $\bigcup (a_\lambda, \infty) = (\inf S, \infty)$. For the forward inclusion, if $x \in (a_{\lambda_0}, \infty)$ for some $a_{\lambda_0} \in S$, we have that $x > a_{\lambda_0} \geq \inf S$ and hence $x \in (\inf S, \infty)$. For the converse inclusion, if $x > \inf S$, by the definition of the greatest lower bound, there exists an $\inf S < a_{\lambda_0} < x$ (otherwise x would be a greater lower bound). Hence $x \in \bigcup (a_\lambda, \infty)$.

Now suppose that S is not bounded below. I claim then that $\bigcup (a_\lambda, \infty) = \mathbb{R}$. The forward direction follows since each interval is a subset of \mathbb{R} . For the converse direction, given $x \in \mathbb{R}$, since S is not bounded below, there exists $a_{\lambda_0} \in S$ with $a_{\lambda_0} < x$ and thus $x \in \bigcup (a_\lambda, \infty)$ as desired.

Hence \mathcal{T}_{ray} is a topology. \square

Problem 1.7

Let (X, \mathcal{T}) be a topological space, and let $A \subseteq X$ be a set with the property that for every $x \in A$, there is an open set $U_x \in \mathcal{T}$ such that $x \in U_x \subseteq A$. Show that A is open.

Solution: This simply follows by noting that $A = \bigcup_{x \in A} U_x$. Converse direction follows since each U_x is a subset of A forward direction follows because for all $x \in X$, $x \in U_x \subseteq \bigcup_{x \in A} U_x$. Since each U_x is open, A is open as it is the union of open sets. \square

Problem 1.8

Let (X, \mathcal{T}) be a topological space, and let $f : X \rightarrow Y$ be an injective (but not necessarily surjective) function. Is $\mathcal{T}_f := \{f(U) : U \in \mathcal{T}\}$ necessarily a topology on Y ? Is it necessarily a topology on the range of f ?

Solution: This is not necessarily a topology on Y . Let $Y = \mathbb{R} \times \{0, 1\}$ and let X be $\mathbb{R}_{\text{usual}}$. Define the function $f : X \rightarrow Y$ by $x \mapsto (x, 0)$. We see that $f(U) = U \times \{0\}$ for all non empty sets $U \in \mathbb{R}$ and $f(\emptyset) = \emptyset$. In particular, $Y \notin \mathcal{T}_f$ and so this is not a topology on Y . \square

It is a topology on the range of f . We have that $f(\emptyset) = \emptyset$ and $f(X) = \text{Range } f$ (and \emptyset and X are open in X). Note that since f is injective, we have that $f(U_1 \cap U_2) = f(U_1) \cap f(U_2)$. The forward inclusion is true for all functions since if $y = f(x)$ with $x \in U_1 \cap U_2$, then $y \in f(U_1)$ and $y \in f(U_2)$. For the reverse inclusion, say $y = f(x_1)$ and $y = f(x_2)$ with $x_i \in U_i$. Since f is injective, we have that $x_1 = x_2$ and hence they are both elements of U_1 and U_2 . Hence $y \in f(U_1 \cap U_2)$. Since $f(U_1 \cap U_2) = f(U_1) \cap f(U_2)$, we note that \mathcal{T}_f is closed under finite intersections as $U_1 \cap U_2$ will be open in X whenever U_1 and U_2 are both open in X .

Now consider arbitrary unions. We first prove that for arbitrary collection of sets $\{U_\lambda\}_{\lambda \in \Lambda}$, $f(\bigcup_{\lambda \in \Lambda} U_\lambda) = \bigcup_{\lambda \in \Lambda} f(U_\lambda)$. The forward direction follows because if $y = f(x)$ where $x \in \bigcup_{\lambda \in \Lambda} U_\lambda$, then $x \in U_{\lambda_0}$ and so $y = f(x) \in f(U_{\lambda_0}) \subseteq \bigcup_{\lambda \in \Lambda} f(U_\lambda)$. For the converse direction, if $y \in \bigcup_{\lambda \in \Lambda} f(U_\lambda)$, then $y = f(x)$ where $x \in U_{\lambda_0} \subseteq \bigcup_{\lambda \in \Lambda} U_\lambda$ and hence $y = f(x) \in f(\bigcup_{\lambda \in \Lambda} U_\lambda)$. Hence \mathcal{T}_f is closed under arbitrary union since $\bigcup_{\lambda \in \Lambda} U_\lambda$ will be open in X whenever each U_λ is open in X .

Hence $\mathcal{T}_{\text{co-finite}}$ is indeed a topology on the range of f . \square

Problem 1.9

Let X be a set and \mathcal{T}_1 and \mathcal{T}_2 be two topologies on X . Is $\mathcal{T}_1 \cup \mathcal{T}_2$ a topology on X ? Is $\mathcal{T}_1 \cap \mathcal{T}_2$ a topology on X ? If yes, prove it. If not, give a counterexample.

Solution: Unions of topologies are not necessarily topologies. Consider $X = \{1, 2, 3\}$ and $\mathcal{T}_1 = \{\emptyset, \{1\}, X\}$ and $\mathcal{T}_2 = \{\emptyset, \{2\}, X\}$. Note that $\mathcal{T}_1 \cup \mathcal{T}_2 = \{\emptyset, \{1\}, \{2\}, X\}$. This is not a topology since $\{1, 2\} = \{1\} \cup \{2\} \notin \mathcal{T}_1 \cup \mathcal{T}_2$ but both of $\{1\}$ and $\{2\}$ are in $\mathcal{T}_1 \cup \mathcal{T}_2$.

Intersections of topologies are topologies. Let X be a set and \mathcal{T}_1 and \mathcal{T}_2 be arbitrary topologies. By definition, $\emptyset, X \in \mathcal{T}_1 \cap \mathcal{T}_2$. Let $U, V \in \mathcal{T}_1 \cap \mathcal{T}_2$. By the definition of a topology, $U \cap V \in \mathcal{T}_1$ and $U \cap V \in \mathcal{T}_2$. Hence, $U \cap V \in \mathcal{T}_1 \cap \mathcal{T}_2$. If $\{U_\lambda\}_{\lambda \in \Lambda}$ are all open sets in $\mathcal{T}_1 \cap \mathcal{T}_2$, then by the definition of a topology $\bigcup U_\lambda \in \mathcal{T}_1$ and $\bigcup U_\lambda \in \mathcal{T}_2$. Hence, $\bigcup U_\lambda \in \mathcal{T}_1 \cap \mathcal{T}_2$ and we have that $\mathcal{T}_1 \cap \mathcal{T}_2$ is still a topology. \square

Problem 1.10

Let X be an infinite set. Show that there are infinitely many distinct topologies on X

Solution: Note that for any $x \in X$, $\mathcal{T}_x = \{\emptyset, \{x\}, X\}$ forms a topology on X . We see that $\emptyset, X \in \mathcal{T}_x$ and that arbitrary unions and intersections stay in \mathcal{T}_x . Thus to get infinitely many distinct topologies, we just consider $\{\mathcal{T}_x : x \in X\}$. \square

Problem 1.11

Fix a set X , and let ϕ be a property that subsets A of X can have. For example, ϕ could be “ A is countable”, or “ A is finite”. ϕ could be “ A contains p ” or “ A doesn’t contain p ” for a fixed point $p \in X$. If $X = \mathbb{R}$, ϕ could be “ A is an interval” or “ A contains uncountably many irrational numbers less than π ”. Define

$$T_{\text{co-}\phi} = \{U \subseteq X : U = \emptyset, \text{ or } X \setminus U \text{ has } \phi\}$$

Under what assumptions on ϕ is $T_{\text{co-}\phi}$ a topology on X ? Which topologies we have seen so far can be described in this way, using which ϕ ?

Solution: First we note that $U \neq \emptyset \in T_{\text{co-}\phi}$ if and only if $U = X \setminus V$ where V satisfies ϕ (in particular $V = X \setminus U$). Finite intersections $(X \setminus V_1) \cap (X \setminus V_2)$ are of the form $X \setminus (V_1 \cup V_2)$, so we require that finite unions of sets satisfying ϕ still satisfy ϕ . Also $\bigcup X \setminus V_\lambda = X \setminus \bigcap V_\lambda$, so we require arbitrary intersections of sets satisfying ϕ to still satisfy ϕ . Finally since we want $X \in T_{\text{co-}\phi}$, we require that \emptyset satisfies ϕ . Cocountable, cofinite, and particular point topology all satisfy this where the properties are “is finite”, “is countable” and “does not contain p ”. We can also phrase the discrete topology in this way with the property “is a set”. We can phrase the trivial topology in this way by using the property “is the empty set”. The ray topology can be phrased as “is an interval of the form $(-\infty, a)$ or is equal to the emptyset”.

Problem 1.12

Let $\{\mathcal{T}_\alpha : \alpha \in I\}$ be a collection of topologies on a set X , where I is some indexing set. Prove that there is a unique finest topology that is refined by all the \mathcal{T}_α . That is, prove that there is a topology \mathcal{T} on X such That

- a) \mathcal{T}_α refines \mathcal{T} for every $\alpha \in I$.
- b) If \mathcal{T}' is another topology that is refined by \mathcal{T}_α for every $\alpha \in I$, then \mathcal{T} is finer than \mathcal{T}'

Solution: We simply take $\mathcal{T} = \bigcap_{\alpha \in I} \mathcal{T}_\alpha$. We see that this is a subset of each \mathcal{T}_α which means that condition a) is satisfied. Also if \mathcal{T}' is a subset of each \mathcal{T}_α it is thus a subset of their intersection which satisfies condition b). It remains to be shown that this is actually still a topology.

We see that \emptyset, X are elements of each \mathcal{T}_α and is thus an element of \mathcal{T} . If $U_1, U_2 \in \mathcal{T}$, then $U_1, U_2 \in \mathcal{T}_\alpha$ for all $\alpha \in I$, so $U_1 \cap U_2 \in \mathcal{T}_\alpha$ for all $\alpha \in I$ and hence $U_1 \cap U_2 \in \mathcal{T}$. For unions, suppose $\{U_\lambda\}_{\lambda \in \Lambda}$ is a collection of sets in \mathcal{T} . They are thus contained in each \mathcal{T}_α . Since each \mathcal{T}_α is a topology, $\bigcup_{\lambda \in \Lambda} U_\lambda \in \mathcal{T}_\alpha$ for all α and hence the union is contained in \mathcal{T} as desired. \square

Problem 1.13

This extends Problem 1.8. Show that f being injective is necessary. That is given an example of a topological space (X, \mathcal{T}) and a non-injective function $f : X \rightarrow Y$ such that \mathcal{T}_f is a topology on the range of f and an example where it is not a topology.

Solution: For an example where \mathcal{T}_f is a topology consider $f : \mathbb{R}_{\text{usual}} \rightarrow \mathbb{R}_{\text{usual}}$ where f is the constant map that maps everything to 1. \mathcal{T}_f in this case is simply $\{\emptyset, \{1\}\}$ which is just the trivial topology on the range of f which is $\{1\}$.

For an example where \mathcal{T}_f is not a topology, let $X = \{0, 1, 2, 3\}$ and $Y = \{0, 1, 2\}$. Define $f : X \rightarrow Y$ where $f(0) = f(1) = 0$, $f(2) = 1$ and $f(3) = 2$ (note that f is surjective here). Define the following topology on X : $\mathcal{T} = \{\emptyset, \{0, 2\}, \{1, 3\}, X\}$. The resulting set for \mathcal{T}_f is $\{\emptyset, \{0, 1\}, \{0, 2\}, Y\}$. Note however that this is not a topology since $\{0, 1\} \cap \{0, 2\} = \{0\}$ but $\{0\} \notin \mathcal{T}_f$.

Problem 1.14

Working in $\mathbb{R}_{\text{usual}}$:

- Show that every nonempty open set contains a rational number.
- Show that there is no uncountable collection of pairwise disjoint open subsets of \mathbb{R} .

Solution:

- Let U be a nonempty open set. Let $x \in U$. Since U is open, there exists an ϵ greater than 0 such that the interval $(x - \epsilon, x + \epsilon) \subseteq U$. By the density of the rationals, there exists a rational number q contained in the interval which is a subset of U as desired. \square
- Suppose for contradiction an uncountable collection of pairwise disjoint open subsets of \mathbb{R} exists. By part a), each of these open subsets would contain a rational number and since the subsets are all disjoint, each of these rational numbers would be distinct and we would end up with an uncountable collection of rational numbers. This is a contradiction since there are countably many rational numbers. \square

2 Bases for Topologies

Problem 2.1

Show explicitly that the collection $\mathcal{B} = \{(a, b) \subseteq \mathbb{R} : a < b\}$ is basis, and that it generates the usual topology on \mathbb{R} .

Solution: We can write \mathbb{R} as $\bigcup_{n \in \mathbb{N}} (n, n+2)$ and thus \mathcal{B} covers \mathbb{R} . Suppose (a_1, b_1) and (a_2, b_2) intersect non trivially, say x is contained in the intersection. We know that there exists an $\epsilon_1 > 0$ such that $(x - \epsilon_1, x + \epsilon_1) \subseteq (a_1, b_1)$ and $\epsilon_2 > 0$ such that $(x - \epsilon_2, x + \epsilon_2) \subseteq (a_2, b_2)$. If we take $\epsilon = \min(\epsilon_1, \epsilon_2)$, we then have that the interval $(x - \epsilon, x + \epsilon)$ is contained in both of $(x - \epsilon_1, x + \epsilon_1)$ and $(x - \epsilon_2, x + \epsilon_2)$. Hence it is contained in both of (a_1, b_1) and (a_2, b_2) . Thus \mathcal{B} is a basis.

To show that this generates the usual topology on \mathbb{R} we note by the definition of the usual topology, we have that if U open, for all $x \in U$ there exists an open interval centered at x that is contained in U (let's call this interval (a_x, b_x)). Then, we simply write $U = \bigcup_{x \in U} (a_x, b_x)$ and we have that every open set in $\mathbb{R}_{\text{usual}}$ can be written as the union of elements of \mathcal{B} and hence \mathcal{B} generates the usual topology of \mathbb{R} . \square

Problem 2.2

Show that $\mathcal{B}_{\mathbb{Q}} = \{(a, b) \subseteq \mathbb{R} : a, b \in \mathbb{Q}, a < b\}$ is a basis for the usual topology on \mathbb{R} .

Solution: Like in Problem 2.1, we write \mathbb{R} as $\bigcup_{n \in \mathbb{N}} (n, n+2)$ and thus $\mathcal{B}_{\mathbb{Q}}$ covers \mathbb{R} . Now again suppose, (a_1, b_1) and (a_2, b_2) intersect non trivially (here $a_i, b_i \in \mathbb{Q}$) at x . Problem 2.1 gives us that there exists an interval (a, b) containing x that is contained in both of these intervals. In particular $b < \min(b_1, b_2)$ and $a > \max(a_1, a_2)$. By the density of the rationals, there exists rational numbers q and r such that $b < r < \min(b_1, b_2)$ and $\max(a_1, a_2) < q < a$. Thus the interval (q, r) still contains x but is also still contained in both of (a_1, b_1) and (a_2, b_2) as desired. Hence, $\mathcal{B}_{\mathbb{Q}}$ is a basis.

To show that this generates the usual topology on \mathbb{R} , it suffices to show that open intervals can be expressed as unions of intervals in $\mathcal{B}_{\mathbb{Q}}$. Then by Problem 2.1, we have that every open set is a union of open intervals which are each a union of intervals in $\mathcal{B}_{\mathbb{Q}}$. By the density of the rationals, for all $n \in \mathbb{N}$, there exists a rational number q_n such that $a < q_n < \min(b, a + \frac{1}{n})$. Similarly, we also have that there exists r_n such that $\max(a, b - \frac{1}{n}) < r_n < b$. We then note that $(a, b) = \bigcup_{n \in \mathbb{N}} (q_n, r_n)$. The reverse inclusion follows because each $(q_n, r_n) \subseteq (a, b)$. For the forward inclusion, if $a < x < b$. We know that there exists an $n \in \mathbb{N}$ such that $\frac{1}{n} < \min(x - a, b - x)$. In particular, this means that $x > a + \frac{1}{n} > q_n$ and $x < b - \frac{1}{n} < r_n$ and we have that $x \in (q_n, r_n)$. Hence every open interval can be expressed as a union of intervals in $\mathcal{B}_{\mathbb{Q}}$ which as discussed above implies that $\mathcal{B}_{\mathbb{Q}}$ generates the usual topology of \mathbb{R} . \square

Problem 2.3

Some exercises about the Sorgenfrey line. Recall the collection $\mathcal{B} = \{[a, b) \subseteq \mathbb{R} : a < b\}$ is a basis which generates \mathcal{S} , the Lower Limit Topology. The space $(\mathbb{R}, \mathcal{S})$ is called the Sorgenfrey line.

- Show that every nonempty open set in \mathcal{S} contains a rational number.
- Show that the interval $(0, 1)$ is open in the Sorgenfrey line.
- More generally, show that for any $a < b \in \mathbb{R}$, (a, b) is open in the Sorgenfrey line.
- Is the interval $(0, 1]$ open \mathcal{S} ?
- Show that \mathcal{S} strictly refines the usual topology on \mathbb{R}
- Show that the real numbers can be written as the union of two disjoint, nonempty open sets in \mathcal{S}
- Let $\mathcal{B}_{\mathbb{Q}} = \{[a, b) : a, b \in \mathbb{Q}, a < b\}$. Show that $\mathcal{B}_{\mathbb{Q}}$ is *not* a basis for the Lower Limit Topology

Solution:

- A non empty open set U in \mathcal{S} must contain an element of the basis, say $[a, b)$. Note that $(a, b) \subseteq [a, b) \subseteq U$ and by Problem 1.14, there exists a rational number in (a, b) and thus a rational number in U . \square
- This follows from the fact that we can write $(0, 1) = \bigcup_{n \in \mathbb{N}} [\frac{1}{n}, 1)$. The reverse inclusion is true since each half open interval is contained in $(0, 1)$. The forward direction is true since for all $x \in (0, 1)$ there is a natural number n large enough so that $\frac{1}{n} < x$. \square
- We use the same construction, we write $(a, b) = \bigcup_{n \in \mathbb{N}} [a + \frac{1}{n}, b)$ where if $a + \frac{1}{n} \geq b$, we say that $[a + \frac{1}{n}, b) = \emptyset$. The argument is the exact same as above.
- Yes, we write $(0, 1] = (0, 1) \cup [\frac{1}{3}, 1]$. Then we write $[\frac{1}{3}, 1] = \bigcup_{n \geq 2} [\frac{1}{3}, 1 - \frac{1}{n})$. This shows that $[\frac{1}{3}, 1]$ and we've already shown that $(0, 1)$ is open, thus the union of the two intervals is still open. \square
- Since we showed every open interval is open in \mathcal{S} and we know from Problem 2.1 that the open intervals

form a basis for $\mathbb{R}_{\text{usual}}$, we thus have that every open set in $\mathbb{R}_{\text{usual}}$ can be represented as a union of elements in \mathcal{S} . Hence we have that \mathcal{S} refines $\mathbb{R}_{\text{usual}}$. To show that it strictly refines it, we simply remark that $[0, 1) \in \mathcal{S}$ since it is a basis element, and Problem 1.1 tells us that this is not open in $\mathbb{R}_{\text{usual}}$. \square

f) We will write $\mathbb{R} = (-\infty, 0) \cup [0, \infty)$. $(-\infty, 0)$ is open in \mathcal{S} since $(-\infty, 0) = \bigcup_{n \in \mathbb{N}} [-n, 0)$. $[0, \infty)$ is open in \mathcal{S} since $[0, \infty) = \bigcup_{n \in \mathbb{N}} [0, n)$. These 2 sets are clearly disjoint and non empty thus satisfying the conditions of the problems. \square

g) I will show that $[\sqrt{2}, 5)$ cannot be written as the union of elements in $\mathcal{B}_{\mathbb{Q}}$. Suppose for contradiction that $[\sqrt{2}, 5) = \bigcup_{\lambda \in \Lambda} [q_{\lambda}, r_{\lambda})$ for some collection of rational number intervals $[q_{\lambda}, r_{\lambda})$. We must have that each interval $[q_{\lambda}, r_{\lambda}) \subseteq [\sqrt{2}, 5)$. Thus $q_{\lambda} \geq \sqrt{2}$ for all $\lambda \in \Lambda$. Since these 2 sets are equal, we also have that $\sqrt{2} \in [q_{\lambda_0}, r_{\lambda_0})$ for some $\lambda_0 \in \Lambda$ (since $\sqrt{2} \in [\sqrt{2}, 5)$). Thus we have that $q_{\lambda_0} \leq \sqrt{2} \leq q_{\lambda_0}$ which implies that $q_{\lambda_0} = \sqrt{2}$, a contradiction. \square

Problem 2.4

Recall that the collection $\mathcal{B} = \{\{x\} : x \in X\}$ is a basis for the discrete topology on a set X . If X is a finite set with n elements, then clearly \mathcal{B} also has n elements. Is there a basis with fewer than n elements that generates the discrete topology on X ?

Solution: This is not possible. Suppose such a basis \mathcal{B}' existed. Since \mathcal{B}' has less than n elements, there exists $x \in X$ such that $\{x\} \notin \mathcal{B}'$. Suppose now that $\{x\}$ can be written as the union of elements of \mathcal{B}' . Say $\{x\} = \bigcup \mathcal{C}$ where $\mathcal{C} \subseteq \mathcal{B}'$. Then every element of \mathcal{C} must be a subset of $\{x\}$ which is not possible since $\{x\} \notin \mathcal{B}$ and we would have that each element of \mathcal{C} is the empty set. This a contradiction since the union of empty sets is empty. \square

Problem 2.5

Let $X = [0, 1]^{[0, 1]}$, the set of all functions $f : [0, 1] \rightarrow [0, 1]$. Given a subset $A \subseteq [0, 1]$, let

$$U_A = \{f \in X : f(x) = 0 \text{ for all } x \in A\}$$

Show that $\mathcal{B} = \{U_A : A \subseteq [0, 1]\}$ is a basis for a topology on X .

Solution: First let's show that \mathcal{B} covers X . We simply note that $U_{\emptyset} = X$ since the statement in the set definition is vacuously true (and hence the union of all the U_A will be equal to X). Suppose $U_A \cap U_B \neq \emptyset$. Then there exists $f \in X$ such that $f(x) = 0$ for all $x \in A$ and $f(x) = 0$ for all $x \in B$. In particular, this tells us that $f \in U_{A \cup B}$. Finally note that $U_{A \cup B} \subseteq U_A \cap U_B$ since if $g(x) = 0$ for all $x \in A \cup B$, then $g(x) = 0$ for all $x \in A$ and $g(x) = 0$ for all $x \in B$. Hence \mathcal{B} is a basis as we have checked both conditions in the definition. \square

Problem 2.6

Let \mathcal{B} be a basis on a set X and let $\mathcal{T}_{\mathcal{B}}$ be the topology that it generates. Show that:

$$\mathcal{T}_{\mathcal{B}} = \bigcap \{\mathcal{T} \subseteq \mathcal{P}(X) : \mathcal{T} \text{ is a topology on } X \text{ and } \mathcal{B} \subseteq \mathcal{T}\}$$

That is, show that $\mathcal{T}_{\mathcal{B}}$ is the intersection of all topologies that contain \mathcal{B} .

Solution: The reverse inclusion follows from the fact that $\mathcal{T}_{\mathcal{B}}$ is itself a topology that contains \mathcal{B} so if U is an element of the intersection, $U \in \mathcal{T}_{\mathcal{B}}$. For the forward inclusion, suppose $U \in \mathcal{T}_{\mathcal{B}}$. Let \mathcal{T} be an arbitrary topology of X containing \mathcal{B} . Since U is the union of elements in \mathcal{B} and each set in \mathcal{B} is in \mathcal{T} , by the definition of a topology we have that $U \in \mathcal{T}$ as desired. \square

Problem 2.7

Let $\{\mathcal{T}_{\alpha} : \alpha \in I\}$ be a collection of topologies on a set X , where I is some indexing set. Prove that there is a unique coarsest topology that refines all the \mathcal{T}_{α} . That is, prove that there is a topology \mathcal{T} on X such that

- a) \mathcal{T} refines \mathcal{T}_{α} for every $\alpha \in I$.
- b) If \mathcal{T}' is another topology that refines \mathcal{T}_{α} for every $\alpha \in I$, then \mathcal{T} is coarser than \mathcal{T}'

Solution: Define $\mathcal{S} = \bigcup_{\alpha \in I} \mathcal{T}_{\alpha}$. Similar to Problem 2.6, define a set \mathcal{T} (which we will show is a topology) that is equal to the intersection of all topologies that contain \mathcal{S} . That is:

$$\mathcal{T} = \bigcap \{ \tau \subseteq \mathcal{P}(X) : \tau \text{ is a topology on } X \text{ and } \mathcal{S} \subseteq \tau \}$$

To show that this is a topology, we remark that arbitrary intersections of topologies is still a topology (we also need to show that the intersection is not empty, but this is satisfied by the discrete topology on X). First note that $\emptyset, X \in \mathcal{T}$ since they are elements of each τ by the definition of a topology. If $\{U_{\lambda}\}_{\lambda \in \Lambda}$ are all elements of \mathcal{T} , they are elements of each τ , and thus their union will still be an element of each τ , and is thus in \mathcal{T} as well. Similarly, if $U, V \in \mathcal{T}$, $U, V \in \tau$ for all τ containing \mathcal{S} and by the definition of a topology $U \cap V \in \tau$ for all τ and thus $U \cap V \in \mathcal{T}$.

We have just shown that \mathcal{T} is a topology. Let's check the refinement conditions. We note that \mathcal{T} contains each \mathcal{T}_{α} since each $\mathcal{T}_{\alpha} \subseteq \mathcal{S} \subseteq \mathcal{T}$ (since \mathcal{T} is an intersection of sets which all contain \mathcal{S}). If \mathcal{T}' is a topology that refines each \mathcal{T}_{α} , then we have that $\mathcal{S} \subseteq \mathcal{T}'$, but by the definition of \mathcal{T} , \mathcal{T}' will be contained in the intersection and hence $\mathcal{T} \subseteq \mathcal{T}'$ i.e \mathcal{T} is coarser than \mathcal{T}' as desired. \square

Problem 2.8

Let $m, b \in \mathbb{Z}$ with $m \neq 0$. A set of the form $Z(m, b) = \{mx + b \mid x \in \mathbb{Z}\}$ is called an arithmetic progression.

- a) Show that the collection \mathcal{B} of all arithmetic progressions is a basis on \mathbb{Z} . The topology $\mathcal{T}_{\text{Furst}}$ that \mathcal{B} generates is called the Furstenberg Topology.
- b) Show that every nonempty open set in $\mathcal{T}_{\text{Furst}}$ is infinite.
- c) Let $U \in \mathcal{B}$ be a basic open set. Show that $\mathbb{Z} \setminus U$ is open.
- d) Show that $\mathcal{T}_{\text{Furst}}$ is Hausdorff (i.e for any distinct integers m, n there are disjoint open sets U and V with $m \in U$ and $n \in V$).

Solution:

- a) We first note that $Z(1, 0) = \mathbb{Z}$ and thus the set of all arithmetic progressions cover \mathbb{Z} . Now suppose $n \in Z(m_1, b_1) \cap Z(m_2, b_2)$. We have that $n = m_1x_1 + b_1 = m_2x_2 + b_2$. Consider $Z(m_1m_2, n)$. Note that $n \in Z(m_1m_2, n)$ since $n = 0m_1m_2 + n$. Next note that $Z(m_1m_2, n) \subseteq Z(m_1, b_1) \cap Z(m_2, b_2)$. This is

because if $y = m_1 m_2 x + n \in Z(m_1 m_2, n)$, we have that $y = m_1 m_2 x + m_1 x_1 + b_1 = m_1(m_2 x + x_1) + b_1 \in Z(m_1, b_1)$ and $y = m_1 m_2 x + m_2 x_2 + b_2 = m_2(m_1 x + x_2) + b_2 \in Z(m_2, b_2)$. Hence \mathcal{B} is a basis on \mathbb{Z} . \square

- b) Note that every non empty open set must contain a basis element. Since every basis element has infinitely many elements, then each non empty open set must be infinite. \square
- c) Let's write $U = Z(m, b)$ where $m \neq 0$. If $m = 1$, note that $Z(m, b) = \mathbb{Z}$ and hence, $\mathbb{Z} \setminus U = \emptyset$ which is open. Otherwise we prove the following relationship:

$$\mathbb{Z} \setminus Z(m, b) = \bigcup_{n=b+1}^{b+m-1} Z(m, n)$$

This is equivalent to showing:

$$\mathbb{Z} = \bigcup_{n=b}^{b+m-1} Z(m, n)$$

The converse inclusion is trivial since each $Z(m, n) \subseteq \mathbb{Z}$. For the forward direction, let $y \in \mathbb{Z}$. We know that $y \in Z(m, y)$. By division algorithm, we have that $y - b = mq + r$ for some $q \in \mathbb{Z}$ and $0 \leq r \leq m - 1$. So we have that $y \in Z(m, b + r + mq)$. We now quickly remark that $y \in Z(m, b + r + mq) = Z(m, b + r)$. For the forward inclusion we have that $y = mx + b + r + mq \implies y = m(x + q) + b + r \in Z(m, b + r)$. For the reverse inclusion $y = mx + b + r \implies y = mx - mq + b + r + mq = m(x - q) + b + r + mq \in Z(m, b + r + mq)$. Hence we have that $y \in Z(m, b + r)$. Since we know that $0 \leq r \leq m - 1$, this means that $Z(m, b + r)$ is an element in the union above completing the proof of the above set equality. This set equality precisely tells us that $\mathbb{Z} \setminus Z(m, b)$ is a union of arithmetic progressions and thus is open. \square

- d) Let $m \neq n \in \mathbb{Z}$. WLOG $m < n$. Take $U = Z(n - m + 1, m)$ and $V = Z(n - m + 1, n)$. It is clear that $m \in U$ and $n \in V$ (take $x = 0$ in the definition). To show that sets are disjoint, suppose that $(n - m + 1)x_1 + m = (n - m + 1)x_2 + n$ for some $x_1, x_2 \in \mathbb{Z}$ so that the intersection is non empty. This implies that $(n - m + 1)(x_1 - x_2) = n - m$. Since $0 < n - m < n - m + 1$, this implies $x_1 - x_2 = 0$ as any other value for $x_1 - x_2$ would cause $|(n - m + 1)(x_1 - x_2)| > n - m + 1 > n - m$. Thus we have that $x_1 = x_2$ which contradicts the fact that they are distinct. Hence U and V must be disjoint completing the proof that $\mathcal{T}_{\text{Furst}}$ is Hausdorff. \square

Problem 2.9

Show that the collection $\mathcal{S} = \{(-\infty, b) : b \in \mathbb{R}\} \cup \{(a, \infty) : a \in \mathbb{R}\}$ is a subbasis that generates the usual topology on \mathbb{R} .

Solution: We first show that if \mathcal{B}_1 is a base for a topology \mathcal{T} and if $\mathcal{B}_2 \subseteq \mathcal{T}$ is such that $\mathcal{B}_1 \subseteq \mathcal{B}_2$, then \mathcal{B}_2 is a base for \mathcal{T} as well. This follows quite simply because if $G \in \tau$, there is $\mathcal{C} \subseteq \mathcal{B}_1$ such that $G = \bigcup \{B : B \in \mathcal{C}\}$. But we also have that \mathcal{C} is a subset of \mathcal{B}_2 , since $\mathcal{B}_1 \subseteq \mathcal{B}_2$. Thus \mathcal{B}_2 is also a base for \mathcal{T} .

Let \mathcal{B}' be the set of all finite intersections of \mathcal{S} . Let \mathcal{B} be the base for the topology from a). We note that $\mathcal{B} \subseteq \mathcal{B}'$. This is because an interval of the form (a, b) is equal to the intersection of the intervals $(-\infty, b)$ and (a, ∞) (from Problem 2.1 we know the set of all intervals form a base for the usual topology of \mathbb{R}). From above, we get that \mathcal{B}' is a base for topology from a), and hence \mathcal{S} is a subbase.

Problem 2.10

- a) Let \mathcal{S} be a collection of subsets of a set X that covers X . Show that \mathcal{S} is a subbasis on X .
- b) Give an example of a subbasis on \mathbb{R} that does not generate the usual topology on \mathbb{R}

Solution:

- a) Let \mathcal{B} be the set of all finite intersections of \mathcal{S} . Note that $\mathcal{S} \subseteq \mathcal{B}$ since we can take intersections with just one term. Thus \mathcal{B} covers X since \mathcal{S} covers X . Let $B_1, B_2 \in \mathcal{B}$. Since B_1 and B_2 are both a finite intersection of elements in \mathcal{S} , then $B_1 \cap B_2$ is also a finite intersection of elements in \mathcal{S} , and hence $B_1 \cap B_2 \in \mathcal{B}$. In particular, for any $x \in B_1 \cap B_2$, we simply take $B_1 \cap B_2$ to be the neighbourhood of x that is contained in $B_1 \cap B_2$. Hence \mathcal{B} is a basis as desired. \square
- b) We simply take $\mathcal{S} = \mathcal{P}(\mathbb{R})$. We note that the set of finite intersections we end up with the basis $\mathcal{B} = \mathcal{P}(\mathbb{R})$. Then taking unions, we still end up with $\mathcal{T} = \mathcal{P}(\mathbb{R})$ which is the discrete topology (which is not the usual topology).

Problem 2.11

For a prime number p , let $S_p = \{n \in \mathbb{N} : n \text{ is a multiple of } p\}$

- a) Show that $\mathcal{S} = \{S_p : p \text{ is prime}\} \cup \{\{1\}\}$ is a subbasis on \mathbb{N}
- b) Describe the open sets in the topology generated by \mathcal{S}

Solution:

- a) By Problem 2.10, it suffices to show that \mathcal{S} covers \mathbb{N} . Let $n \in \mathbb{N}$. If $n = 1$, then $n \in \{1\}$ which is an element of \mathcal{S} . Otherwise, there exists a prime p that divides n , and hence $n \in S_p$ which is an element of \mathcal{S} . Since \mathcal{S} covers \mathbb{N} it is a subbasis for \mathbb{N} . \square
- b) First I'll extend the notation S_n to be the multiples of n even when n is not prime. We note that $S_n \cup S_m$ are the multiples of the least common multiple of n and m . And so by taking finite intersections, we generate the basis $\mathcal{B} = \{S_n : n \geq 2 \in \mathbb{N}\} \cup \{\{1\}\}$. The resulting topology has open sets whose elements are the union of these basis elements. So open sets in the topology are simply the union of multiples of numbers.

Problem 2.12

Fix an infinite subset A of \mathbb{Z} whose complement $\mathbb{Z} \setminus A$ is also infinite. Construct a topology on \mathbb{Z} which satisfies the following properties:

- a) A is open
- b) Singletons are never open
- c) The topology is Hausdorff

Solution: We know that since A and $\mathbb{Z} \setminus A$ are subsets of \mathbb{Z} , they are countable (and thus countably infinite since they are infinite by assumption). Let f be a bijection from A to the set of even integers, and g a bijection from $\mathbb{Z} \setminus A$ to the set of odd integers. Then construction a bijection $\tilde{h} : \mathbb{Z} \rightarrow \mathbb{Z}$ as follows:

$$\tilde{h}(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \notin A \end{cases}$$

Let $h = \tilde{h}^{-1}$. Let's equip the domain of h with the Furstenberg topology. By Problem 1.8, since h is bijective in this case, if we take $\mathcal{T}_h = \{h(U) : U \in \mathcal{T}_{\text{Furst}}\}$, we get that \mathcal{T}_h is a topology on \mathbb{Z} . We note that A is the image of the even integers under h and note that the set of even integers is open in $\mathcal{T}_{\text{Furst}}$ since they can be represented as $Z(2, 0)$. Hence the first condition is satisfied. The second condition is satisfied because Problem 2.8 tells us that non empty open sets in $\mathcal{T}_{\text{Furst}}$ are all infinite, and if we take the image of these infinite set under a bijection, they will still be infinite. So open sets in this topology are never singletons. Finally to show Hausdorff, let $a, b \in \mathbb{Z}$. By Problem 2.8, we know that there exists disjoint open sets U, V in $\mathcal{T}_{\text{Furst}}$ such that $h^{-1}(a) \in U$ and $h^{-1}(b) \in V$. Now consider $h(U)$ and $h(V)$ in \mathbb{Z} equipped with \mathcal{T}_h . We have that $a \in h(U)$ and $b \in h(V)$. We also have that they are open by the definition of \mathcal{T}_h . Finally we note that they are disjoint because if $y \in h(U) \cap h(V)$, then $y = h(u)$ for some $u \in U$ and $y = h(v)$ for some $v \in V$. Thus by injectivity of h , we get that $u = v$, which in particular gives us that $u = v \in U \cap V$ which contradicts the fact that U and V are disjoint. Thus $h(U) \cap h(V) = \emptyset$ completing the proof that \mathcal{T}_h is Hausdorff. \square

3 Closed sets, closures, and dense sets

Problem 3.1

Let (X, \mathcal{T}) be a topological space and \mathcal{B} a basis for \mathcal{T} . Let $A \subseteq X$. Show that $x \in \overline{A}$ if and only if for every basic open set U containing x , $U \cap A \neq \emptyset$

Solution: The forward direction follows by the definition because every basic open set is an open set. For the converse direction, let $x \in X$ and V be an arbitrary (not necessarily basic) open set containing x . We know from equivalent definitions of a basis that there exists a basic open set U containing x that is contained in V . By assumption, $U \cap A \neq \emptyset$ and since $U \subseteq V$, we have that $V \cap A \neq \emptyset$

Problem 3.2

Let (X, \mathcal{T}) be a topological space, $A, B \subseteq X$. Is it necessarily true that $\overline{A \cup B} = \overline{A} \cup \overline{B}$? Is it necessarily true that $\overline{A \cap B} = \overline{A} \cap \overline{B}$

Solution: It is true that $\overline{A \cup B} = \overline{A} \cup \overline{B}$. For the forward inclusion, let $x \in \overline{A \cup B}$ and suppose $x \notin \overline{A}$. Then, there exists U containing x such that $U \cap A = \emptyset$. Since $x \in \overline{A \cup B}$, we have that $(U \cap A) \cup (U \cap B) = U \cap (A \cup B) \neq \emptyset$. But since $U \cap A = \emptyset$, this gives us that $U \cap B \neq \emptyset$. Since U was arbitrary, this gives us that $x \in \overline{B}$ as desired. For the reverse inclusion, we simply use the contrapositive. Suppose $x \notin \overline{A \cup B}$, there exists an open set U containing x such that $(U \cap A) \cup (U \cap B) = U \cap (A \cup B) = \emptyset$. This gives us that $U \cap A = U \cap B = \emptyset$ and hence $x \notin \overline{A}$ and $x \notin \overline{B}$ as desired. Thus closure distributes over unions.

It is not true however that closures distribute over intersections. Consider $\mathbb{R}_{\text{usual}}$ with $A = (0, 1)$ and $B = (1, 2)$. We have that $A \cap B = \emptyset$, so $\overline{A \cap B} = \emptyset$. However, $\overline{A} = [0, 1]$ and $\overline{B} = [1, 2]$, so $\overline{A} \cap \overline{B} = \{1\} \neq \emptyset = \overline{A \cap B}$. \square

Problem 3.3

Let $A = \{\frac{1}{n} : n \in \mathbb{N}\}$. In $\mathbb{R}_{\text{usual}}$, show that $\overline{A} = A \cup \{0\}$.

Solution: For the reverse inclusion, we know that $A \subseteq \overline{A}$. We also know that for any open set U around 0, we can take an $\epsilon > 0$ such that $(-\epsilon, \epsilon) \subseteq U$. Note that for any ϵ , there exists $N \in \mathbb{N}$ large enough so that $\frac{1}{N} < \epsilon$. Thus the intersection $U \cap A$ is always non empty.

For the forward inclusion we approach via the contrapositive. Suppose $x \notin A \cup \{0\}$. If $x < 0$, take $\epsilon = |x|/2$ so that $B_\epsilon(x) \cap A = \emptyset$ because all values in $B_\epsilon(x)$ will take on strictly negative values. Now suppose $x > 0$ but $x \notin A$. Let $n \in \mathbb{N}$ be minimal so that $\frac{1}{n} < x$ (we know such an n must exist by Archimedian property). So we have that $\frac{1}{n} < x < \frac{1}{n-1}$ and we know these are strict inequalities because $x \notin A$. So $x \in (\frac{1}{n}, \frac{1}{n-1})$, and by 1.1, we know that this is an open set and there exists an $\epsilon > 0$ so that $B_\epsilon(x) \subseteq (\frac{1}{n}, \frac{1}{n-1})$. Note that this implies $B_\epsilon(x) \cap A = \emptyset$ since for all $k \leq n-1$ we have that $\frac{1}{k} \geq \frac{1}{n-1}$, and for $k \geq n$, we have that $\frac{1}{k} \leq \frac{1}{n}$. So none of the values of A are contained in this interval. In both cases, we have constructed a $B_\epsilon(x)$ so that $B_\epsilon(x) \cap A = \emptyset$. Since $B_\epsilon(x)$ is an open set in $\mathbb{R}_{\text{usual}}$ but has an empty intersection with A , we have that $x \notin \overline{A} \cup \{0\}$ as desired. \square

Problem 3.4

Show the following facts in $\mathbb{R}_{\text{usual}}$. Let $a < b < c \in \mathbb{R}$,

- $\overline{\{a\}} = \{a\}$
- $\overline{[a, b]} = [a, b]$
- $\overline{(a, b) \cup (b, c)} = [a, c]$
- $\overline{[a, b]} = [a, b]$
- If $(x_n)_{n=1}^\infty$ is a sequence converging to $x \in \mathbb{R}$, then $\overline{\{x_n : n \in \mathbb{N}\}} = \{x_n : n \in \mathbb{N}\} \cup \{x\}$

Solution:

- Converse inclusion follows because $A \subseteq \overline{A}$ for all sets A . For the forward inclusion, we use the contrapositive. If $x \notin \{a\}$, i.e $x \neq a$, we can take $\epsilon = |x - a|$, and thus $a \notin B_\epsilon(x) \implies \{a\} \cap B_\epsilon(x) = \emptyset$. Since $B_\epsilon(x)$ is an open set containing x , $x \notin \overline{\{a\}}$ as desired. \square
- For the converse inclusion if $x \in [a, b]$, we are done. So consider $x = b$. In this case, for any open set U containing B , there exists $\epsilon > 0$ such that $B_\epsilon(b) \subseteq U$. We have that $b - \min(\frac{\epsilon}{2}, \frac{b-a}{2}) \in B_\epsilon(b) \cap [a, b]$ and thus the intersection $U \cap [a, b]$ is non empty for arbitrary open sets U containing b . This completes the reverse inclusion.

For the forward inclusion use the contrapositive. If $x < a$, then take $\epsilon = a - x$ and $B_\epsilon(x) \cap [a, b] = \emptyset$. If $x > b$, take $\epsilon = x - b$ and $B_\epsilon(x) \cap [a, b] = \emptyset$. In either case, $B_\epsilon(x)$ is an open set containing x that doesn't intersect with $[a, b]$, so $x \notin \overline{[a, b]}$ as desired. \square

- For the converse inclusion if $x \in (a, b) \cup (b, c)$, we are done. So we are left to consider $x \in \{a, b, c\}$ only. For a , any open set U containing it, there exists $\epsilon > 0$ such that $B_\epsilon(a) \subseteq U$. Then $a + \min(\frac{\epsilon}{2}, \frac{b-a}{2}) \in (a, b) \cap B_\epsilon(a)$ and thus the intersection $U \cap (a, b) \cup (b, c)$ is non empty for arbitrary open sets U

containing a . For b , any open set U containing it, there exists $\epsilon > 0$ such that $B_\epsilon(b) \subseteq U$. Then $a + \min(\frac{\epsilon}{2}, \frac{c-b}{2}) \in (b, c) \cap B_\epsilon(b)$ and thus the intersection $U \cap (a, b) \cup (b, c)$ is non empty for arbitrary open sets U containing b . For c , any open set U containing it, there exists $\epsilon > 0$ such that $B_\epsilon(c) \subseteq U$. Then $c - \min(\frac{\epsilon}{2}, \frac{c-b}{2}) \in (b, c) \cap B_\epsilon(c)$ and thus the intersection $U \cap (a, b) \cup (b, c)$ is non empty for arbitrary open sets U containing c . In all cases, we have that $x \in \overline{(a, b) \cup (b, c)}$

For the forward inclusion use the contrapositive. If $x < a$, then take $\epsilon = a - x$ and $B_\epsilon(x) \cap ((a, b) \cup (b - c)) = \emptyset$. If $x > c$, take $\epsilon = x - c$ and $B_\epsilon(x) \cap ((a, b) \cup (b - c)) = \emptyset$. In either case, $B_\epsilon(x)$ is an open set containing x that doesn't intersect with $(a, b) \cup (b, c)$, so $x \notin \overline{(a, b) \cup (b, c)}$ as desired. \square

- d) Reverse inclusion follows because $A \subseteq \overline{A}$ for all sets A . For the forward inclusion use the contrapositive. If $x < a$, then take $\epsilon = a - x$ and $B_\epsilon(x) \cap [a, b] = \emptyset$. If $x > b$, take $\epsilon = x - b$ and $B_\epsilon(x) \cap [a, b] = \emptyset$. In either case, $B_\epsilon(x)$ is an open set containing x that doesn't intersect with $[a, b]$, so $x \notin \overline{[a, b]}$ as desired. \square

- e) For the reverse inclusion, we only need to show that x is in the closure, the other elements are contained because of the fact that $A \subseteq \overline{A}$ in general. Let U be an arbitrary open set containing x . There exists an $\epsilon > 0$ such that $B_\epsilon(x) \subseteq U$. By the definition of convergence, there exists $n \in \mathbb{N}$ such that $x_n \in B_\epsilon(x) \subseteq U$. Hence the intersection $U \cap \{x_n : n \in \mathbb{N}\}$ is always not empty and hence $x \in \overline{\{x_n : n \in \mathbb{N}\}}$ completing the reverse inclusion.

For the forward inclusion use the contrapositive. Suppose y is not an element of the sequence and is not equal to x . Since limits are unique in $\mathbb{R}_{\text{usual}}$, the sequence doesn't converge to y . In particular, taking $\epsilon = |y - x|/2$, there exists an $N \in \mathbb{N}$ so that $n \geq N$ implies that $x_n \in B_\epsilon(x)$. In particular, by the triangle inequality, this means that $n \geq N$ implies that $x_n \notin B_\epsilon(y)$ and that $x \notin B_\epsilon(y)$. For each $1 \leq k \leq N$, define $\epsilon_k = |y - x_k|/2 > 0$ (since $y \neq x_k$ for all $k \in \mathbb{N}$). So this construction gives us that $x_k \notin B_{\epsilon_k}(y)$. Finally let's take $\epsilon_0 = \min(\epsilon, \epsilon_1, \dots, \epsilon_N)$. We then have that $B_{\epsilon_0}(y)$ is an open set containing y that doesn't contain any elements of the sequence or x . Hence $y \notin \overline{\{x_n : n \in \mathbb{N}\}}$ as desired. \square

Problem 3.5

Let (X, \mathcal{T}) be a topological space and let $A \subseteq X$. Show that

$$\overline{A} = \bigcap \{F \subseteq X : A \subseteq F \text{ and } F \text{ is closed}\}$$

Solution: For both inclusions, we prove via the contrapositive.

(\subseteq)

Suppose x is not an element of the set on the right hand side. Thus there exists a closed set F with $A \subseteq F$ such that $x \notin F$, i.e $x \in X \setminus F$. $X \setminus F$ is an open set since F is closed and since $A \subseteq F$ we have that $A \cap (X \setminus F) = \emptyset$. Since we have found an open set containing x that doesn't intersect with A , $x \notin \overline{A}$. \square

(\supseteq)

Suppose $x \notin \overline{A}$. There exists an open set U containing x such that $U \cap A = \emptyset$. This implies that $A \subseteq X \setminus U$. Since U is open, $X \setminus U$ is closed, so this set is contained in the intersection on the right side of the equation. However, since $x \notin X \setminus U$ (since $x \in U$), we have that x cannot be an element of an intersection containing $X \setminus U$, and hence x is not an element of the right side. \square

Problem 3.6

Show that if (X, \mathcal{T}) is a topological space and $D \subseteq X$, show that D is dense if and only if for every nonempty open set $U \subseteq X$, $D \cap U \neq \emptyset$.

Solution: For the forward implication, suppose $\overline{D} = X$. Let U be an arbitrary non empty open set. Then there exists $x \in U$. By assumption, we have that $x \in \overline{D}$. So by the definition of closure, $U \cap D \neq \emptyset$ as desired.

For the reverse implication. Suppose all nonempty open sets intersect non trivially with D . We then see that $\overline{D} = X$ since for arbitrary $x \in X$ and open sets U_x containing x , we get by our assumption that $U_x \cap D \neq \emptyset$ so $x \in \overline{D}$ as desired. \square

Problem 3.7

In an arbitrary topological space, is the union of two dense sets necessarily dense? What about the intersection of two dense sets? For both questions, prove it or give a counterexample

Solution: The union is necessarily dense. Suppose D_1 and D_2 are dense. We note that $D_1 \subseteq D_1 \cup D_2$ and since D_1 is dense, $D_1 \cup D_2$ is dense. In general if A is dense with $A \subseteq B$, then B is dense.

Intersections are not necessarily dense. In $\mathbb{R}_{\text{usual}}$, we have that the set of rationals, and irrationals are both dense. However, their intersection is the empty set which is not dense. \square

Problem 3.8

Show that

$$\text{int}(A) = \bigcup \{U \subseteq X : U \subseteq A \text{ and } U \text{ is open}\}$$

Solution: For the forward inclusion, suppose $x \in \text{int}(A)$. Then there exists an open set U_x containing x with $U_x \subseteq A$ by definition. Thus, U_x is an element of the union on the right hand side. Since $x \in U_x$, then x is an element of the right side.

For the reverse inclusion, if x is an element of the union, there exists an open set U such that $x \in U$ and $U \subseteq A$. This is precisely the condition for $x \in \text{int } A$

Problem 3.9

Show for any topological space (X, \mathcal{T}) and any $A \subseteq X$, that $\text{int}(A)$ is open

Solution: This follows from problem 3.8 since the interior is a union of open sets and is thus open. \square

Problem 3.10

Show that a subset A of a topological space X is open if and only if $A = \text{int}(A)$.

Solution: The reverse implication follows since we've already showed in problem 3.9 that the interior is open. For the forward implication, suppose A is open. Using the equivalent definition for the interior in problem 3.8, we see that $\text{int}(A) \subseteq A$ since it is a union of subsets of A . Also, we see that if A is open, then A is an element of the union and so $A \subseteq \text{int}(A)$. Hence $A = \text{int}(A)$ completing the proof. \square

Problem 3.11

Compute the interior and closures of the following sets in the given spaces:

- a) $(0, 1]$ in $\mathbb{R}_{\text{usual}}$
- b) $(0, 1]$ in the Sorgenfrey line
- c) $(0, 1]$ in $(\mathbb{R}, \mathcal{T}_{\text{trivial}})$
- d) $(0, 1]$ in $(\mathbb{R}, \mathcal{T}_{\text{discrete}})$
- e) $(0, 1]$ in $(\mathbb{R}, \mathcal{T}_{\text{ray}})$
- f) $(0, 1]$ in $(\mathbb{R}, \mathcal{T}_{\text{co-finite}})$
- g) The set E of even numbers in $(\mathbb{Z}, \mathcal{T}_{\text{co-finite}})$
- h) \mathbb{Q} in $\mathbb{R}_{\text{usual}}$
- i) \mathbb{Q} in the Sorgenfrey line
- j) $\mathbb{Q} \times \mathbb{Q}$ in $\mathbb{R}_{\text{usual}}^2$
- k) $\{(x, y, z) \in \mathbb{R}^3 : x = 0\}$ in $\mathbb{R}_{\text{usual}}^3$

Solution:

- a) The closure is $[0, 1]$ via a similar proof to problem 3.4. The interior is $(0, 1)$. We note that $(0, 1)$ is open by problem 1.1 and that it is a subset of $[0, 1]$. Thus the interior must contain $(0, 1)$. However, $1 \notin \text{int}(0, 1]$ (so the interior cannot be $(0, 1]$) since every open set containing 1 will contain an element larger than 1 which is not in $(0, 1]$. Thus the interior must be $(0, 1)$. \square
- b) We showed that $(0, 1]$ is open in the Sorgenfrey line in Problem 2.3. Thus by problem 3.10, we have that $(0, 1]$ is its own interior. We also have that $\mathbb{R} \setminus (0, 1] = (-\infty, -1) \cup [-1, 0] \cup (1, \infty)$. By Problem 2.3, we know that the topology of the Sorgenfrey line refines the usual topology of \mathbb{R} . Since $(-\infty, -1)$ and $(1, \infty)$ are open in the usual topology on \mathbb{R} , they are open in the Sorgenfrey line. $[-1, 0]$ is open because we can write it as $[-1, -\frac{1}{2}) \cup (-\frac{1}{2}, 0]$ which are both open by 2.3. Hence, $\mathbb{R} \setminus (0, 1]$ is open, implying that $(0, 1]$ is closed and equal to its own closure. \square
- c) The only open/closed sets in $\mathcal{T}_{\text{trivial}}$ are \emptyset and \mathbb{R} . By using 3.8, the only open set contained by $(0, 1]$ is \emptyset so the interior is \emptyset . The only closed set containing $(0, 1]$ is \mathbb{R} , so the closure is \mathbb{R} by problem 3.5. \square
- d) In the discrete topology every set is both open and closed. So the interior and closure of $(0, 1]$ is itself. \square

- e) In \mathcal{T}_{ray} , open sets are of the form (a, ∞) (or the empty set and \mathbb{R}), and so closed sets are of the form $(-\infty, b]$. Note that there does not exist a nonempty open set that is contained in $(0, 1]$ and hence the interior is equal to \emptyset . Note that $(0, 1] \subseteq (-\infty, 1]$ and $(-\infty, 1]$ is closed, so the closure must be a closed subset of $(-\infty, 1]$. However, every proper closed subset of $(-\infty, 1]$ is of the form $(-\infty, b]$ for $b < 1$ so $1 \notin (-\infty, b]$. This implies that the closure must be equal to $(-\infty, 1]$ since $(0, 1]$ itself must be a subset of its closure. \square
- f) The closed sets in $\mathcal{T}_{\text{co-finite}}$ are the finite sets and \mathbb{R} itself. Since $(0, 1]$ is infinite, there are no finite sets that contain it, so the only closed set containing it is \mathbb{R} . Thus, by problem 3.5, the closure must be \mathbb{R} . Since the complement of $(0, 1]$ is infinite, every subset of $(0, 1]$ will contain infinitely many values and so every non empty subset will not be open in $\mathcal{T}_{\text{co-finite}}$. The only open subset of $(0, 1]$ is \emptyset . Thus by problem 3.8, the interior will be equal to \emptyset . \square
- g) We can generalize the above argument to say that if X is a set equipped with the co-finite topology and $A \subseteq X$ is such that A is infinite and $X \setminus A$ is infinite, then the closure of A is X and the interior of A is \emptyset . If A is infinite, there are no finite (i.e closed) sets that contain it except for X . So by problem 3.5, the closure of A must be X . If $X \setminus A$ is infinite, every subset of A will contain infinitely many elements in its complement (since it must contain $X \setminus A$). Thus the only open set contained in A is the empty set, which by problem 3.8 tells us that the interior is \emptyset . So in this case, the set of even numbers is infinite and so is the complement (odd integers), and thus the interior is \emptyset and closure is X . \square
- h) We've shown in problem 1.14 that every open set in $\mathbb{R}_{\text{usual}}$ intersects non trivially with \mathbb{Q} . Furthermore, problem 3.6 tells us that this implies that \mathbb{Q} is dense, and hence $\overline{\mathbb{Q}} = \mathbb{R}$. Next, we show that there are no nonempty open subsets of \mathbb{Q} . Suppose such an open subset U existed. Let $q \in U$. Let V be an arbitrary open set containing q . Since U is open there must be open interval $(q - \epsilon, q + \epsilon) \subseteq U$. However this open interval intersects non trivially with the irrationals by problem 1.14. This contradicts the fact that U is contained in the rationals. Hence the only non empty open subset contained in \mathbb{Q} is \emptyset which by problem 3.8 we have that the interior is \emptyset . \square
- i) \mathbb{Q} is still dense in the Sorgenfrey Line because every open set U contains a basis element of the form $[a, b)$ which will always contain a rational number. Hence $\overline{\mathbb{Q}} = \mathbb{R}$. We can always follow the same proof as for $\mathbb{R}_{\text{usual}}$ to show that interior is empty because any open set must contain an interval $[a, b)$ (which is a basis element). Since $(a, b) \subseteq [a, b)$, we have that this interval must contain an irrational number. The proof continues in the same way and we get that the interior is \emptyset . \square
- j) $\mathbb{Q} \times \mathbb{Q}$ is still dense in $\mathbb{R}_{\text{usual}}^2$ since given a non empty open set U , say it contains a point (x, y) . There exists an open ball such that $B_\epsilon(x, y) \subseteq U$ ($\epsilon > 0$). Consider the interval $(x - \frac{\epsilon}{\sqrt{2}}, x + \frac{\epsilon}{\sqrt{2}}) \in \mathbb{R}$, by density of rationals (problem 1.14), there is a rational number q_1 in this interval. Similarly, there exists a rational number q_2 in the interval $(y - \frac{\epsilon}{\sqrt{2}}, y + \frac{\epsilon}{\sqrt{2}})$. Then, $d((q_1, q_2), (x, y)) = \sqrt{|q_1 - x|^2 + |q_2 - y|^2} < \sqrt{\left(\frac{\epsilon}{\sqrt{2}}\right)^2 + \left(\frac{\epsilon}{\sqrt{2}}\right)^2} = \epsilon$. So $(q_1, q_2) \in U \cap \mathbb{Q} \times \mathbb{Q}$ and hence $\overline{\mathbb{Q} \times \mathbb{Q}} = \mathbb{R}^2$.
- A similar justification will allow us to show that the interior is empty. Suppose a non empty open subset U of $\mathbb{Q} \times \mathbb{Q}$ existed. Say $(x, y) \in U$. There exists an open ball such that $B_\epsilon(x, y) \subseteq U$ ($\epsilon > 0$). Consider the interval $(x - \epsilon, x + \epsilon)$, there exists an irrational number z in this interval. Then $d((z, y), (x, y)) = |z - x| < \epsilon$, so $(z, y) \in B_\epsilon(x, y) \subseteq U$. However, $(z, y) \notin \mathbb{Q} \times \mathbb{Q}$. So there cannot be a non empty open subset of $\mathbb{Q} \times \mathbb{Q}$ implying that the interior is the empty set. \square

Problem 3.12

Let A be a subset of a topological space X . Show that $\partial(A) = \overline{A} \cap \overline{X \setminus A} = \overline{A} \setminus \text{int}(A)$.

Solution: The first equality $\partial(A) = \overline{A} \cap \overline{X \setminus A}$ follows exactly from the definition. $x \in \partial(A)$ if and only if every open set intersects non trivially with A and $X \setminus A$ which is equivalent to x being an element of \overline{A} and $\overline{X \setminus A}$.

To complete the proof we show $\partial(A) = \overline{A} \setminus \text{int}(A)$.

$$\begin{aligned}
 x \in \partial(A) &\iff \text{for all open sets } U_x \text{ containing } x, U_x \cap A \neq \emptyset \text{ and } U_x \cap (X \setminus A) \neq \emptyset \\
 &\iff \text{for all open sets } U_x \text{ containing } x, U_x \cap A \neq \emptyset \text{ and } U_x \not\subseteq A \\
 &\iff x \in \overline{A} \text{ and } x \notin \text{int}(A) \\
 &\iff x \in \overline{A} \setminus \text{int}(A). \quad \square
 \end{aligned}$$

Problem 3.13

Let A be a subset of a topological space X . Show that $\overline{A} = A \cup \partial(A)$ and $\text{int}(A) = A \setminus \partial(A)$.

Solution:

a) $\overline{A} = A \cup \partial(A)$

We know that $A \subseteq \overline{A}$ and that $\partial(A) = \overline{A} \cap \overline{X \setminus A}$ by problem 3.12, so we also have that $\partial(A) \subseteq \overline{A}$. Thus the reverse inclusion has been shown. For the forward inclusion, again from problem 3.12, we have that $\partial(A) = \overline{A} \setminus \text{int}(A) \implies \overline{A} \subseteq \partial(A) \cup \text{int}(A) \subseteq \partial(A) \cup A$ as desired ($\text{int}(A)$ is contained in A since it is a union of subsets of A). \square

b) $\text{int}(A) = A \setminus \partial(A)$

We know that $\text{int}(A) \subseteq A$ as mentioned above. We also have from problem 3.12 that $\partial(A) = \overline{A} \setminus \text{int}(A)$ so $x \in \text{int}(A) \implies x \notin \partial(A)$. This completes the forward inclusion. For the reverse inclusion, if $x \in A$ and $x \notin \partial(A)$, there exists a neighbourhood of U_x of x such that $U_x \cap A = \emptyset$ or $U_x \cap (X \setminus A) = \emptyset$. Since $x \in A$, we know that $U_x \cap A \neq \emptyset$, so we must have that $U_x \cap (X \setminus A) = \emptyset$. This is equivalent to $U_x \subseteq A$ and hence $x \in \text{int}(A)$.

Problem 3.14

Let A be a subset of a topological space X . Show that $X = \text{int}(A) \sqcup \partial(A) \sqcup \text{int}(X \setminus A)$

Solution: First for showing equality of the union, suppose $x \in X$ with $x \notin \text{int}(A)$ and $x \notin \text{int}(X \setminus A)$. Then for all open sets U_x containing x , $U_x \not\subseteq A$ which is equivalent to $U_x \cap (X \setminus A) \neq \emptyset$ and $U_x \not\subseteq (X \setminus A)$ which is equivalent to $U_x \cap A \neq \emptyset$. These 2 combined precisely are the definition of $x \in \partial(A)$.

For disjointness, $\text{int}(A) \subseteq A$ and $\text{int}(X \setminus A) \subseteq X \setminus A$ so $\text{int}(A)$ and $\text{int}(X \setminus A)$ are disjoint since A and $X \setminus A$ are disjoint. $\partial(A)$ and $\text{int}(A)$ are disjoint since by problem 3.12, $\partial(A) = \overline{A} \setminus \text{int}(A)$ so if $x \in \text{int}(A)$, $x \notin \partial(A)$ so there cannot be an element in their intersection. Finally, we note that $\partial(A) = \partial(X \setminus A)$ because the definition is symmetric when we replace A with $X \setminus A$. We then get that $\text{int}(X \setminus A)$ is disjoint with $\partial(X \setminus A) = \partial(A)$. We have shown all 3 sets are pairwise disjoint and thus the union is indeed a disjoint union. \square

Problem 3.15

A subset A of a topological space X is called regular open if $\text{int}(\overline{A}) = A$. Regular open sets play an important role in set theoretic topology.

- Show that in $\mathbb{R}_{\text{usual}}$, any open interval (a, b) is regular open
- Let A be a subset of a topological space X . Is it true that $\text{int}(\overline{A}) = \overline{\text{int}(A)}$? If not, is there containment one way or the other?
- Show that the intersection of two regular open sets is again regular open (in any topological space)
- Is the union of two regular open sets again regular open? Prove it or give a counterexample.
- Given a subset A of a topological space X , let $A^\perp = X \setminus \overline{A}$. Show that a set A is regular open if and only if $(A^\perp)^\perp = A$.

Solution:

- We know that $\overline{(a, b)} = [a, b]$ by 3.4. The interior of $[a, b]$ is (a, b) . It suffices to show that a and b both cannot be in the interior, and since we know that the interior must contain all open sets, it follows that the interior is (a, b) . Note that every open set U containing a , contains an open interval $B_\epsilon(a) \subseteq U$. We see that this is not contained in (a, b) since $a - \frac{\epsilon}{2}$ is not in (a, b) . Similarly for b , any open set V containing b contains an open interval $B_\epsilon(b) \subseteq V$ which is not contained in (a, b) since $b + \frac{\epsilon}{2}$ is not in (a, b) . \square
- We note that this equality doesn't necessarily hold because $\text{int}(\overline{A})$ is a open set while $\overline{\text{int}(A)}$ is a closed set and in a lot of topological spaces, there are very few clopen sets. A concrete counter example will be (a, b) in $\mathbb{R}_{\text{usual}}$. In part a), we showed that $\text{int}(\overline{A}) = (a, b)$. We can also see that $\overline{\text{int}(A)} = [a, b]$ since $A = \text{int}(A)$ as (a, b) is open. This example also shows us that the reverse inclusion is false.

The forward implication is not true either. Recall from problem 3.11 if we take \mathbb{R} equipped with the co-finite topology, and take $A = (0, 1]$ (or any infinite subset whose complement is infinite), $\text{int}(A) = \emptyset$ and $\overline{A} = \mathbb{R}$. Thus, $\text{int}(\overline{A}) = \emptyset$ and $\overline{\text{int}(A)} = \mathbb{R}$. So we don't necessarily have that, $\text{int}(\overline{A}) \subseteq \overline{\text{int}(A)}$. \square

- Suppose A and B are regular open sets. We have that $A \cap B = \text{int}(\overline{A \cap B})$ by definition of regular open sets. We seek to show that $A \cap B = \text{int}(\overline{A \cap B})$. For the forward inclusion we note that $A \cap B \subseteq \overline{A \cap B}$. Thus, taking the interior of both sides and we get that $\text{int}(A \cap B) \subseteq \text{int}(\overline{A \cap B})$ (problem 3.8 gives us the property that interior maintains the inclusion). Since A and B are both open, we have that $A \cap B$ is open and thus $\text{int}(A \cap B) = A \cap B$ (problem 3.10). Hence $A \cap B = \text{int}(A \cap B) \subseteq \text{int}(\overline{A \cap B})$.

For the converse inclusion we first remark that in general, the interior distributes over intersection, i.e if $\text{int}(C \cap D) = \text{int}(C) \cap \text{int}(D)$. The forward direction follows from the fact that $C \cap D$ is a subset of both C and D so $\text{int}(C \cap D) \subseteq \text{int}(C)$ and $\text{int}(C \cap D) \subseteq \text{int}(D)$. For the reverse inclusion we note that if $x \in \text{int}(C)$ and $x \in \text{int}(D)$, there exists open sets U_1 and U_2 containing x such that $U_1 \subseteq C$ and $U_2 \subseteq D$. Then if we take $U_1 \cap U_2$, it is still open and we have that $U_1 \cap U_2 \subseteq C \cap D$ and so $x \in \text{int}(C \cap D)$. Also note that $\overline{A \cap B} \subseteq \overline{A} \cap \overline{B}$ because $A \cap B \subseteq A$ and $A \cap B \subseteq B$, and so $\overline{A \cap B}$ is a subset of both \overline{A} and \overline{B} (problem 3.5). Again, we can take the interior of both sides which tells us that $\text{int}(\overline{A \cap B}) \subseteq \text{int}(\overline{A} \cap \overline{B}) = \text{int}(\overline{A}) \cap \text{int}(\overline{B}) = A \cap B$. The first equality follows from the fact that interior splits over intersections, and the second because A and B are regular open sets. This completes the proof of the converse direction and the set equality. \square

- The union of regular open sets is not necessarily open. In $\mathbb{R}_{\text{usual}}$, take the intervals $(0, 1)$ and $(1, 2)$

which are regular open by a). The union is $(0, 1) \cup (1, 2)$. This however is not regular open as its closure is $[0, 2]$ (by problem 3.4) whose interior is $(0, 2) \neq (0, 1) \cup (1, 2)$.

- e) It suffices to show that $(A^\perp)^\perp = \text{int}(\overline{A})$ for all subsets A . Unravelling the definition, $(A^\perp)^\perp = X \setminus \overline{(X \setminus \overline{A})}$. Note that $\text{int}(\overline{A}) = X \setminus \overline{(X \setminus \overline{A})} \iff X \setminus \text{int}(\overline{A}) = \overline{(X \setminus \overline{A})}$. Now note the following chain of equivalences to see this equality:

$$\begin{aligned} x \notin \overline{A} &\iff \text{For all neighbourhoods } U_x \text{ of } x, U_x \not\subseteq \overline{A} \\ &\iff \text{For all neighbourhoods } U_x \text{ of } x, U_x \cap (X \setminus \overline{A}) \neq \emptyset \\ &\iff x \in X \setminus \overline{A} \end{aligned}$$

As discussed, this set equality implies that A is regular open if and only if $(A^\perp)^\perp = A$. \square

Problem 3.16

Let A be a subset of \mathbb{R}^n with its usual topology. Show that $x \in \overline{A}$ if and only if there exists a sequence of elements of A that converges to x .

Solution: For the forward direction, let $x \in \overline{A}$. For $n \in \mathbb{N}$, since $x \in \overline{A}$, $B_{\frac{1}{n}}(x) \cap A \neq \emptyset$. So for each $n \in \mathbb{N}$, choose an $x_n \in B_{\frac{1}{n}}(x) \cap A$ forming a sequence of elements in A (invoking the axiom of choice). Note that $(x_n)_{n=1}^\infty$ converges to x because for all $\epsilon > 0$, choose N such that $\frac{1}{N} < \epsilon$, and for $n \geq N$, $x_n \in B_{\frac{1}{n}}(x) \subseteq B_{\frac{1}{N}}(x) \subseteq B_\epsilon(x)$.

For the converse direction, suppose $(x_n)_{n=1}^\infty$ is a sequence in A that converges to $x \in X$. Let U be an arbitrary neighbourhood of x . There exists an $\epsilon > 0$ such that $B_\epsilon(x) \subseteq U$. Since the sequence converges to x , there exists a $x_{n_0} \in B_\epsilon(x) \subseteq U$. Since the sequence x_n is contained in A , the intersection $U \cap A$ is non-empty (it contains x_{n_0}) and hence $x \in \overline{A}$. \square

Problem 3.17

We have already learned that \mathbb{Q} is dense in \mathbb{R} with its usual topology. Is $\mathbb{Q} \setminus \{0\}$ dense? How about if you remove finitely many points from \mathbb{Q} ? Is there an infinite set of points you can remove from \mathbb{Q} that leaves the resulting set dense?

Solution: I show that if we remove finitely many points from \mathbb{Q} , the resulting set is still dense. Suppose $A = \{a_1, \dots, a_n\} \subseteq \mathbb{Q}$. Consider $\mathbb{Q} \setminus A$. Let U be an arbitrary non empty open set in $\mathbb{R}_{\text{usual}}$. Let $x \in U$, and thus there exists an interval $I = (x - \epsilon, x + \epsilon) \subseteq U$ (where $\epsilon > 0$). If $A \cap I = \emptyset$, then we get that $I \cap (\mathbb{Q} \setminus A) \neq \emptyset$ since we know the interval must contain a rational number and that rational number cannot be in A . Now suppose $A \cap I \neq \emptyset$. Define $m = \max A \cap I$ (we can take this maximum because A is finite). Consider the interval $(m, x + \epsilon) \subseteq I$. This interval must contain a rational number q , and note that $q \notin A$ because that would contradict the maximality of m . Thus we have that $I \cap (\mathbb{Q} \setminus A) \neq \emptyset$ since it contains this rational number q . In both cases, we have that $I \cap (\mathbb{Q} \setminus A) \neq \emptyset$ and since $I \subseteq U$, this means that $U \cap (\mathbb{Q} \setminus A) \neq \emptyset$. Since U was arbitrary, $\mathbb{Q} \setminus A$ is dense. This also tells us that $\mathbb{Q} \setminus \{0\}$ is dense.

We can indeed remove infinitely many sets and still have the resulting set be dense. Consider $\mathbb{Q} \setminus \mathbb{N}$. Let U be an arbitrary non empty open set in $\mathbb{R}_{\text{usual}}$. Let $x \in U$, and thus there exists an interval $I = (x - \epsilon, x + \epsilon) \subseteq U$ (where $\epsilon > 0$). If $\mathbb{N} \cap I = \emptyset$, then we get that $I \cap (\mathbb{Q} \setminus \mathbb{N}) \neq \emptyset$ since we know the interval must contain a rational number and that rational number cannot be a natural number. Now suppose $\mathbb{N} \cap I \neq \emptyset$. Define $m = \min \mathbb{N} \cap I$ (we can take this minimum because \mathbb{N} is well ordered under the usual ordering, and so this

subset of \mathbb{N} must contain a minimal element). Consider the interval $(x - \epsilon, m) \subseteq I$. This interval must contain a rational number q , and note that $q \notin \mathbb{N}$ because that would contradict the minimality of m . Thus we have that $I \cap (\mathbb{Q} \setminus \mathbb{N}) \neq \emptyset$ since it contains this rational number q . In both cases, we have that $I \cap (\mathbb{Q} \setminus \mathbb{N}) \neq \emptyset$ and since $I \subseteq U$, this means that $U \cap (\mathbb{Q} \setminus \mathbb{N}) \neq \emptyset$. Since U was arbitrary, $\mathbb{Q} \setminus \mathbb{N}$ is dense. \square

Problem 3.18

Let (X, \mathcal{T}) be a topological space, and let D_1 and D_2 be dense open subsets of X . Prove that $D_1 \cap D_2$ is dense and open. Give an example in $\mathbb{R}_{\text{usual}}$ to show that this does not extend even to countably infinite intersections. That is, give an example of a collection $\{D_n : n \in \mathbb{N}\}$ of dense open subsets of $\mathbb{R}_{\text{usual}}$ such that $\bigcap_{n=1}^{\infty} D_n$ is not open (as you will soon see, such an intersection must be dense).

Solution: Let D_1, D_2 be dense and open. Their intersection is open by the definition of a topology. To show that they are dense, let U be an arbitrary non empty subset of X . Since D_1 is dense, $U \cap D_1$ is non empty. Since U and D_1 are both open, $U \cap D_1$ is open. Since D_2 is dense, $U \cap D_1 \cap D_2$ is not empty and hence $D_1 \cap D_2$ is dense.

Let $D_n = \mathbb{R} \setminus \{\frac{1}{n}\}$. Problem 3.4, gives us that singletons are equal to their closures and are thus closed, so D_n is open for all $n \in \mathbb{N}$. It is dense because every non empty open set in \mathbb{R} must contain at least 2 distinct elements (well they contain infinitely many but this is enough) since they aren't singletons, and we see that this implies that it must intersect with \mathbb{R} with only 1 element removed. Note however that $S = \bigcup_{n=1}^{\infty} D_n$ is not open because $0 \in S$, but for every open set U containing 0, there is an interval $B_{\epsilon}(0) \subseteq U$, but we know that there exists $\frac{1}{n} < \epsilon$ and so every open set will contain an element not in S and thus S is not open. \square

Problem 3.19

Recall the Furstenberg topology $\mathcal{T}_{\text{Furst}}$ on \mathbb{Z} (problem 2.8), introduced in the exercises from the previous section. To remind you, this is the topology on \mathbb{Z} generated by the basis consisting of all infinite arithmetic progressions in \mathbb{Z} . Earlier, you proved that every nonempty open subset in $\mathcal{T}_{\text{Furst}}$ is infinite. You also proved that for every basic open set U in $\mathcal{T}_{\text{Furst}}$, $\mathbb{Z} \setminus U$ is open. We now know this is the same as saying every basic open subset is closed. You are going to use this topology to give a slick, elegant proof that there are infinitely many prime numbers.

- a) Using the notation from when this topology was first introduced, show that:

$$\mathbb{Z} \setminus \{-1, 1\} = \bigcup_{p \text{ is prime}} Z(p, 0)$$

- b) Assume for the sake of contradiction that there are only finitely many primes. Deduce from this assumption that $\mathbb{Z} \setminus \{-1, 1\}$ is closed.
- c) Find a contradiction resulting from the previous part, and conclude that there must be infinitely many primes.

Solution:

- a) For the reverse inclusion, note that $Z(p, 0) = \{np : n \in \mathbb{Z}\}$. Note that no scalar multiple of a prime is equal to 1 or -1 which completes the reverse inclusion. For the forward inclusion, if $n \in \mathbb{Z} \setminus \{-1, 1\}$ we consider 2 cases, $n = 0$ or otherwise. If $n = 0$, note that $0 = 7(0)$, so $0 \in Z(7, 0)$ and since 7 is prime n is an element of the right hand side. Now suppose n is non zero, by prime factorization, there exists a prime p such that $p \mid n$, i.e $n = pk$ for some $k \in \mathbb{Z}$. Thus $n \in Z(p, 0)$ and thus is an element of

the right hand side. In both cases we see that the forward inclusion stands. \square

- b) As mentioned in the problem statement, we've shown that every basic open set is closed. Thus, if there are finitely many primes, we have expressed $\mathbb{Z} \setminus \{-1, 1\}$ as the union of finitely many closed sets, and by the definition of a topology, this means that it must be closed. \square
- c) As mentioned in the problem statement, we've shown that all non empty open sets are infinite. If $\mathbb{Z} \setminus \{-1, 1\}$ is closed, then $\{-1, 1\}$ must be open. This is a contradiction as this set is not infinite, hence there must be infinitely many primes. \square