Supplemental Table S1. Postpartum incidence of negative health events and the proportion of primiparous and multiparous cows included in each model for evaluating the association between different biomarkers and negative health events for a prospective cohort study involving 72 farms across the northeastern US. Cows diagnosed with the negative health event of interest prior to the postpartum blood sample collection were removed from the postpartum biomarker models. Herds that did not have accurate records for the negative health event of interest were removed from the analysis

	Prepartum NEFA[1]		Postpartum NEFA and BHB		Haptoglobin	
Outcome[2]	n cows (% primiparous; % multiparous)	Disorder Incidence	n cows (%primiparous; % multiparous)	Disorder Incidence	n cows (%primiparous; % multiparous)	Disorder Incidence
DA	1468 (32%; 68%)	2.3%	1461 (32%; 68%)	1.4%	981 (30%; 70%)	1.9%
CK	1444 (32%; 68%)	7.1%	1398 (33%; 67%)	3.7%	944 (31%; 69%)	4.7%
MET[3]	1369 (32%; 68%)	12.4%	1290 (31%; 69%)	1.9%	865 (29%; 71%)	2.4%
CULL30	1468 (32%; 68%)	2.5%	1473 (32%; 68%)	2.5%	988 (30%; 70%)	3.2%
DA, CK, MET[3], or any combination	1345 (32%; 68%)	18.1%	1219 (32%; 68%)	5.7%	824 (30%; 70%)	7.0%

[1]NEFA = nonesterified fatty acids

[2]DA = displaced abomasum, CK = clinical ketosis, MET = metritis, and CULL30 = culling within 30 DIM.

[3]In the prepartum NEFA model, retained placenta was assessed concurrently with MET.