Lecture 1: Diversification and Mean-Variance

Mark Hendricks

Autumn 2024

FINM 36700: Portfolio Management

The Big Picture

The Big Picture

Diversification

Mean-Variance

Excess Returns

Appendi



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 2/69

The Big Picture Diversification Mean-Variance Excess Returns Appendix

Key results

- ▶ Portfolio risk is a nonlinear function of security risk.
- ▶ If we assume frictionless markets, then we can analytically solve for "optimal" return-risk allocations.
- ► The optimal formula penalizes securities for marginal risk (covariance), not total risk (volatility.)
- ► The result is analytical, efficiently implemented, and maximizes portfolio Sharpe Ratio.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 3/69

Key questions

- ► What do we mean by "optimal"?
- ▶ What is the right measure of risk?
- ► How do we forecast returns?
- ► How well does this work in practice?



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 4/69

Portfolio Management Procedure

- 1. Define the security universe
- 2. Model security risk and performance
- Forecast returns
- 4. Define the portfolio's objective
- 5. Define portfolio's constraints
- 6. Simulate the candidate portfolios
- 7. Optimize among the portfolios
- 8. Assess the constructed portfolio's performance



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 5/69

Getting Started

To begin,

- we need to understand diversification and non-linearity of risk.
- we will examine the full portfolio optimization process.

During the course, we will dig deeper into each of the steps of the process.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 6/69

Outline

Diversification



Hendricks, Autumn 2024 FINM 36700: Lecture 1 7/69

Return notation: one-period

notation	description	formula	example
r^{i}	return rate of asset i		
$r^{\scriptscriptstyle f}$	risk-free return rate		
$ ilde{\pmb{r}}^i$	excess return rate of asset i	$r^i - r^f$	



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 8/69

Two investments: bonds and stocks

Consider the following portfolio example

Table: Portfolio example

	return	allocation weight
bonds	r^b	W
stocks	rs	1 - w

Table: Return statistics notation

mean	variance	correlation
	0	
μ	σ^2	ho



Hendricks, Autumn 2024 FINM 36700: Lecture 1 9/69

Portfolio return stats

Investment portfolio return r^p has mean and variance of

$$\mu^p = w\mu^b + (1-w)\mu^s$$

$$\sigma_p^2 = w^2 \sigma_b^2 + (1 - w)^2 \sigma_s^2 + 2w(1 - w)\rho \sigma_s \sigma_b$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 10/69

Perfect correlation

Suppose that $\rho = 1$.

► Then the volatility (standard deviation) of the portfolio is proportional to the asset allocation weights:

$$\sigma_p = w\sigma_b + (1 - w)\sigma_s$$

▶ Thus, both mean and volatility are linear in the allocations.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 11/69

Imperfect correlation

Suppose that $\rho < 1$.

► The volatility function is convex,

$$\sigma_p < w\sigma_b + (1-w)\sigma_s$$

▶ Yet the mean return is still linear in the portfolio allocation:

$$\mu^p = w\mu^b + (1-w)\mu^s$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 12/69

Diversification

Portfolio diversification refers to this case where

- mean returns are linear in allocations
- while volatility of returns is less than linear in allocation.

This only required $\rho < 1$.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 13/69

A perfect hedge

For ho=-1 ,

- ► The portfolio variance can be as small as desired, by choosing the appropriate allocation, w.
- ▶ In fact, $\sigma_p = 0$ if

$$w = \frac{\sigma_s}{\sigma_b + \sigma_s}$$

► Thus, a riskless portfolio can be formed from the two risky assets.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 14/69





Figure: Diversification of investment portfolio between two risky assets.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 15/69



The Big Picture

Figure: Diversification over investment in U.S. market index and 10-year T-note. Source: CRSP and N.Y. Fed. July 1971 to June 2012.

 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 16/69

Allocation among *n* assets

Consider the following portfolio allocation problem:

- n risky securities,
- return volatility (std.dev.) denoted σ_i
- return covariance between security i and j denoted by $\sigma_{i,j}$.
- w^i denotes the fraction of the portfolio allocated to asset i, with $\sum_{i=1}^{n} w^i = 1$.

Then

$$\sigma_p^2 = \sum_{i=1}^n \sum_{j=1}^n w^i w^j \sigma_{i,j}$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 17/69

Variance of the equally weighted portfolio

Consider an equally-weighted portfolio, with $w^i=1/n$ for each asset. Then

$$\sigma_p^2 = \frac{1}{n^2} \sum_{i=1}^n \sigma_i^2 + \frac{1}{n^2} \sum_{i \neq i} \sum_{i=1}^n \sigma_{i,j}$$

In the earlier example with bonds and stocks, n = 2,

$$\sigma_p^2 = \frac{1}{4}\sigma_b^2 + \frac{1}{4}\sigma_s^2 + \frac{1}{2}\underbrace{\sigma_{b,s}}_{\rho\sigma_b\sigma_s}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 18/69

Portfolio variance as average covariances

Use the following notation for averaging the variances and covariances across the n assets:

$$\operatorname{avg}\left[\sigma_{i}^{2}\right] \equiv \frac{1}{n} \sum_{i=1}^{n} \sigma_{i}^{2}$$

$$\operatorname{avg}\left[\sigma_{i,j}\right] \equiv \frac{1}{n(n-1)} \sum_{i \neq i} \sum_{i=1}^{n} \sigma_{i,j}$$

So the portfolio variance can be written as

$$\sigma_p^2 = \frac{1}{n} \operatorname{avg} \left[\sigma_i^2 \right] + \frac{n-1}{n} \operatorname{avg} \left[\sigma_{i,j} \right]$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 19/69

Portfolio irrelevance of individual security variance

As number of securities in portfolio, n, gets large,

$$\lim_{n\to\infty}\sigma_p^2=\operatorname{avg}\left[\sigma_{i,j}\right]$$

- ▶ Individual security variance is unimportant!
- Overall portfolio variance is average of individual security covariance.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 20/69

Diversified portfolio

Obtained this result using equally-weighted portfolio, $w^i = 1/n$.

▶ Don't need equal weighting, just that

$$\lim_{n\to\infty} w^i = 0$$

- ► That is, as *n* gets large the portfolio must have trivial exposure to security *i*.
- ➤ This is the sense in which portfolio must be diversified for individual variances to become unimportant.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 21/69

Portfolio variance decomposition

Above we saw the equally-weighted portfolio variance:

$$\sigma_p^2 = \frac{1}{n} \operatorname{avg}\left[\sigma_i^2\right] + \frac{n-1}{n} \operatorname{avg}\left[\sigma_{i,j}\right]$$

Variance has a term which can be diversified to zero, and another term that remains.

Suppose that asset returns have

- ightharpoonup identical volatilities, $\sigma_i = \sigma$
- ightharpoonup identical correlations, $\rho_{i,j} = \rho$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 22/69

Systematic risk

$$\sigma_p^2 = \frac{1}{n}\sigma^2 + \frac{n-1}{n}\rho\sigma^2$$

$$\lim_{n\to\infty}\sigma_p^2\to\underbrace{\rho\sigma^2}_{\text{systematic}}$$

- \blacktriangleright A fraction, ρ , of the variance is systematic.
- ▶ No amount of diversification¹ can get portfolio variance lower:

$$\sigma_p^2 \ge \rho \sigma^2$$



¹Inequality holds for any n and any set of allocations $\{w^i\}$.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 23/69

Idiosyncratic risk

$$\sigma_p^2 = \frac{1}{n}\sigma^2 + \frac{n-1}{n}\rho\sigma^2$$

- ▶ Idiosyncratic risk refers to the diversifiable part of σ_p^2 .
- An equally-weighted portfolio ² has idiosyncratic risk equal to $\frac{1}{n}\sigma^2$.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 24/69

²For general weights, w^i , remaining idiosyncratic risk is bounded by $\max_i w^i \sigma^2$.

Correlation and diversified portfolios

$$\sigma_p^2 = \frac{1}{n}\sigma^2 + \frac{n-1}{n}\rho\sigma^2$$

For $\rho = 1$, there is no possible diversification, regardless of n.

$$\sigma_p^2 = \sigma^2$$

For $\rho=0$, there is no systematic risk, only variance is remaining idiosyncratic:

$$\sigma_p^2 = \frac{1}{n}\sigma^2$$

And as n gets large the portfolio is riskless,

$$\lim_{n\to\infty}\sigma_p^2=0$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 25/69

The Big Picture Diversification Mean-Variance Excess Returns Appendix

Riskless portfolios

- Above, we found that a riskless portfolio could be created if $\rho = -1$.
- ► Here, we found that a riskless portfolio can be created if $\rho = 0$.

Question:

How did the assumptions behind these conclusions differ?



Hendricks, Autumn 2024 FINM 36700: Lecture 1 26/69

Answer:

- ▶ In the case of just two underlying assets, complete diversification is achieved with $\rho = -1$.
- ▶ In the case of many assets, complete diversification is achieved when all assets are uncorrelated, and the number of assets in the portfolio goes to infinity.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 27/69

Outline

The Big Picture

Diversification

Mean-Variance

Excess Returns

Appendia



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 28/69

Mean-variance comparisons


We want to compare risk and return...

- Use mean return to score the portfolio's benefits.
- Use variance (or volatility) of return to score the portfolio's risk.

Consider the case of two assets:



Hendricks, Autumn 2024 FINM 36700: Lecture 1 29/69



The Big Picture

Figure: Example in mean-volatility space of diversification between two assets.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 30/69

Diversification across *n* assets


With n securities, there is further potential for diversification.

- ► The set of all possible portfolios formed from this basis of assets forms a convex set in mean-variance space.
- ► The boundary of this set is known as the mean-variance frontier, and it forms a parabola.
- ► The boundary of the set in mean-volatility space forms a hyperbola.

We use **MV** frontier to refer to both the mean-variance and mean-volatility frontiers.




 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 31/69



The Big Picture

Figure: Mean-variance frontier formed by 25 U.S. equity portfolios, sorted by size and and book/market.

 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 32/69



The Big Picture

Figure: Mean-volatility frontier formed by 25 U.S. equity portfolios, sorted by size and and book/market.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 33/69

Efficient portfolios

The top segment of the MV frontier is the set of efficient MV portfolios.

- ► These portfolios maximize mean return given the return variance.
- ► Contrast this with the lower segment of the MV frontier, the inefficient MV portfolios.
- ► The inefficient MV portfolios minimize mean return given the return variance.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 34/69

Importance of MV analysis

- ► MV analysis is the most widely used tool in portfolio allocation.
- ▶ The model gives a tractable way to balance risk and return.
- ► Later in the course, we will a connection between MV analysis and beta-factor models.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 35/69

The Big Picture Diversification Mean-Variance Excess Returns Appendix

Notation

Suppose there are n risky assets.

- **r** is an $n \times 1$ random vector. Each element is the return on one of the n assets.
- Let μ denote the $n \times 1$ vector of mean returns. Let **Σ** denote the $n \times n$ covariance matrix of returns.

$$oldsymbol{\mu} = \mathbb{E}\left[oldsymbol{r}
ight] \ oldsymbol{\Sigma} = \mathbb{E}\left[\left(oldsymbol{r} - oldsymbol{\mu}
ight)\left(oldsymbol{r} - oldsymbol{\mu}
ight)'
ight]$$

- For now, we suppose no risk-free rate is available.
- ▶ Assume ∑ is positive definite—no asset is a linear function of the others.

CHICAGO

Hendricks, Autumn 2024 FINM 36700: Lecture 1 36/69

Portfolios

- ▶ An investor chooses a **portfolio**, defined as a $n \times 1$ vector of allocation weights, ω .
- ► These allocation weights must sum to unity:

$$\omega' \mathbf{1} = 1$$

where **1** denotes a $n \times 1$ vector of ones.

ightharpoonup No shorting restriction here: elements of ω can be negative.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 37/69

Return moments

The portfolio return on some portfolio, ω^p , is

$$r^{p}=\left(\omega^{p}
ight) ^{\prime}oldsymbol{r}.$$

The portfolio return moments are

$$\mu^p =: \mathbb{E}\left[r^p\right] = \left(\omega^p\right)' \mu$$
 $\sigma_p^2 =: \operatorname{var}(r^p) = \left(\omega^p\right)' \mathbf{\Sigma} \omega^p$
 $\operatorname{cov}(r^p, \mathbf{r}) = \mathbf{\Sigma} \omega^p$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 38/69

MV Portfolio

A Mean-Variance (MV) portfolio is a vector, ω^* , which solves the following optimization for some number μ^p :

min
$$\omega' \mathbf{\Sigma} \omega$$

s.t. $\omega' \mu = \mu^p$
 $\omega' \mathbf{1} = 1$

- Note that the objective function is convex in w, given that Σ is positive definite.
- ► The constraint set is also convex.
- ▶ Thus, the solution, ω^* is characterized by the first-order conditions.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 39/69

MV solution

Thus, a portfolio ω^* is MV iff exists $\delta \in (-\infty, \infty)$ such that

$$egin{aligned} oldsymbol{\omega}^* = & \delta oldsymbol{\omega}^{ t t} + (1 - \delta) oldsymbol{\omega}^{ t v} \ & \omega^{ t t} \equiv \underbrace{\left(rac{1}{\mathbf{1}' oldsymbol{\Sigma}^{-1} \mu}
ight)}_{ ext{scaling}} oldsymbol{\Sigma}^{-1} oldsymbol{\mu}, \qquad oldsymbol{\omega}^{ t v} \equiv \underbrace{\left(rac{1}{\mathbf{1}' oldsymbol{\Sigma}^{-1} \mathbf{1}}
ight)}_{ ext{scaling}} oldsymbol{\Sigma}^{-1} oldsymbol{1} \end{aligned}$$

 $oldsymbol{\omega}^{ t t}$ and $oldsymbol{\omega}^{ t v}$ are themselves MV portfolios $(\delta=0,1)$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 40/69

GMV and zero-tangency portfolios

 ω^{v} is the Global Minimum Variance (GMV) portfolio. It solves,


$$\min_{\omega} \;\; \omega' \mathbf{\Sigma} \omega$$
 s.t. $\omega' \mathbf{1} = 1$

► This is the same as the MV problem, but dropping the first constraint, ($\omega' \mu = \mu^p$.)

 ω^{t} is the portfolio tangent to the mean-volatility frontier and going through the origin. (See next slide.)



Hendricks, Autumn 2024 FINM 36700: Lecture 1 41/69



The Big Picture

Figure: Illustration of two useful MV portfolios. The Global-Minimum-Variance portfolio as well as the zero-tangency portfolio.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 42/69

MV investors

Consider MV investors, the investors for whom mean and variance of returns are sufficient statistics of the investment.

- ightharpoonup Such investors will hold an MV portfolio, ω^* .
- ▶ Thus, these investors are holding linear combination of just two risky portfolios, ω^{t} and ω^{v} .
- ➤ So if in real markets all investors were MV investors, everyone would simply invest in two funds.
- Those wanting higher mean returns would hold more in the high-return MV, ω^t , while those wanting safer returns would hold more in the low-return MV, ω^v .



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 43/69

Outline

The Big Picture

Diversification

Mean-Variance

Excess Returns

Appendi



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 44/69

With a riskless asset

Now consider the existence a risk-free asset with return, r^{f} .

- ightharpoonup Suppose there are still n risky assets available, still notating the risky returns as r
- ▶ Let w denote a n × 1 vector of portfolio allocations to the n risky assets.
- Since the total portfolio allocations must add to one, we have allocation to the risk-free rate =1-w'1



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 45/69

Mean excess returns

 μ denotes the vector of mean returns of risky assets, $\mathbb{E}\left[r \right]$.

Let μ^p denote the mean return on a portfolio.

$$\mu^p = \left(1 - \mathbf{w}'\mathbf{1}\right)r^{\scriptscriptstyle f} + \mathbf{w}'\boldsymbol{\mu}$$

Use the following notation for excess returns:

$$ilde{m{\mu}} = m{\mu} - \mathbf{1} r^{\scriptscriptstyle f}$$

Thus the mean return and mean excess return of the portfolio are

$$\mu^{p} = r^{f} + \mathbf{w}' \tilde{\boldsymbol{\mu}}$$
 $\tilde{\mu}^{p} = \mathbf{w}' \tilde{\boldsymbol{\mu}}$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 46/69

Variance of returns

- ► The risk-free rate has zero variance and zero correlation with any security.
- ▶ Let Σ continue to denote the $n \times n$ covariance matrix of *risky* assets, (and is positive semi-definite.)
- ▶ The return variance of the portfolio, \mathbf{w}^p is

$$\sigma_p^2 = \mathbf{w}' \mathbf{\Sigma} \mathbf{w}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 47/69

The MV problem with a riskless asst

A Mean-Variance portfolio with risk-free asset ($\mathring{\text{MV}}$) is a vector, \boldsymbol{w}^* , which solves the following optimization for some mean excess return number $\tilde{\mu}^p$:

min
$$oldsymbol{w}'oldsymbol{\Sigma}oldsymbol{w}$$
 s.t. $oldsymbol{w}'oldsymbol{ ilde{\mu}}= ilde{\mu}^p$

- ▶ In contrast to the MV problem, there is only one constraint.
- ► The allocation weight vector, **w** need not sum to one, as the remainder is invested in the risk-free rate.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 48/69

Solving the MV problem

Solving the problem is straitforward:

- 1. Set up the Lagrangian with just one constraint.
- 2. The FOC is sufficient given the convexity of the problem.
- 3. Finally, substitute the Lagrange multiplier using the constraint.

Refer to the solution as an MV portfolio.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 49/69

MV solution

$$\mathbf{w}^* = ilde{\delta} \; \mathbf{w}^{ exttt{t}}$$

for the portfolio

$$oldsymbol{w}^{ ext{t}} = \underbrace{\left(rac{1}{\mathbf{1}'oldsymbol{\Sigma}^{-1} ilde{oldsymbol{\mu}}}oldsymbol{\Sigma}^{-1} ilde{oldsymbol{\mu}}$$

and allocation

$$ilde{\delta} = \left(rac{\mathbf{1}'\mathbf{\Sigma}^{-1} ilde{oldsymbol{\mu}}}{(ilde{oldsymbol{\mu}})'\mathbf{\Sigma}^{-1} ilde{oldsymbol{\mu}}}
ight) ilde{\mu}^{oldsymbol{p}}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 50/69

MV portfolio variance formula

The return variance of an MV portfolio is given by

$$\frac{(\tilde{\mu}^p)^2}{(\tilde{\mu})' \mathbf{\Sigma}^{-1} \tilde{\mu}}$$

This implies that the return volatility (standard-deviation) is linear in the absolute value of the mean excess return:

$$rac{| ilde{\mu}^{
ho}|}{\sqrt{(ilde{oldsymbol{\mu}})'\,oldsymbol{\Sigma}^{-1} ilde{oldsymbol{\mu}}}}$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 51/69

Tangency portfolio

The result is that any \widetilde{MV} portfolio is a combination of the tangency portfolio, \boldsymbol{w}^{t} , and a position in the riskless asset.

- The tangency portfolio, \mathbf{w}^{t} invests 100% in risky assets, $\mathbf{1}'\mathbf{w}^{t} = 1$.
- ▶ w^t is the unique portfolio which is on the risky MV frontier as well as the MV frontier expanded by the risk-free asset.
- \mathbf{w}^{t} is the point on the risky MV frontier at which the tangency line goes through the risk-free rate. (See the figure below.)



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 52/69




Figure: Illustration of the MV frontier when a riskless asset is available. In this case, the MV portfolio frontier consists of two straight lines. The curved frontier is the MV frontier when a riskless asset is unavailable.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 53/69

Tangency portfolio and the Sharpe ratio

For an arbitrary portfolio, \mathbf{w}^p ,

$$\mathsf{SR}(\boldsymbol{w}^p) = \frac{\mu^p - r^t}{\sigma^p} = \frac{\tilde{\mu}^p}{\sigma^p}$$

The tangency portfolio, \mathbf{w}^{t} , is the portfolio on the risky MV frontier with maximum Sharpe ratio.

$$\mathsf{SR}\left(oldsymbol{w}^*
ight) = \pm \sqrt{\left(ilde{oldsymbol{\mu}}
ight)' oldsymbol{\Sigma}^{-1} ilde{oldsymbol{\mu}}}$$

The SR magnitude is constant across all MV portfolios. (Sign depends on whether part of the efficient or inefficient frontier.)



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 54/69

Capital Market Line

The Capital Market Line (CML) is the efficient portion of the MV frontier.

- The CML shows the risk-return tradeoff available to MV investors.
- ► The slope of the CML is the maximum Sharpe ratio which can be achieved by any portfolio.
- ► The inefficient portion of the MV frontier acheives the minimum (negative) Sharpe ratio by shorting the tangency portfolio.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 55/69




Figure: Illustration of the $\tilde{\text{MV}}$ frontier when a riskless asset is available. In this case, the $\tilde{\text{MV}}$ portfolio frontier consists of two straight lines. The curved frontier is the $\tilde{\text{MV}}$ frontier when a riskless asset is unavailable.

Hendricks, Autumn 2024 FINM 36700: Lecture 1 56/69

Two-fund separation

Two-fund separation. Every MV portfolio is the combination of the risky portfolio with maximal Sharpe Ratio and the risk-free rate.

Thus, for an MV investor the asset allocation decision can be broken into two parts:

- 1. Find the tangency portfolio of risky assets, $\boldsymbol{w}^{\text{t}}$.
- 2. Choose an allocation between the risk-free rate and the tangency portfolio.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 57/69

Intuition of asset allocation

The two-fund separation says that

- ► Any investment in risky assets should be in the tangency portfolio since it offers the maximum Sharpe Ratio.
- One must decide the desired level of risk in the investment, which determines the split between the riskless asset and the tangency portfolio.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 58/69

Conclusion

- Non-additivity of portfolio risk requires us to consider mathematics of diversification.
- Mean-variance optimization is the dominant approach in industry.
- ▶ But implementation will raise a number of challenges, related to computation and statistics.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 59/69

References

▶ Back, Kerry. Asset Pricing and Portfolio Choice Theory. 2010. Chapter 5.

Develops the mathematical formulas for optimization among n assets.

▶ Bodie, Kane, and Marcus. *Investments*. 2011. Chapter 7. Develops the intuition of mean-variance space and optimal portfolios.



Hendricks, Autumn 2024 FINM 36700: Lecture 1 60/69

Outline

The Big Picture

The Big Picture

Diversification

Mean-Variance

Excess Returns

Appendix



Appendix

 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 61/69

Solving the MV problem: FOC

Solving with Lagrangian multipliers, (γ_1 and γ_2 ,) gives the unconstrained optimization:

$$\mathcal{L} = \frac{1}{2} \boldsymbol{\omega}' \boldsymbol{\Sigma} \boldsymbol{\omega} - \gamma_1 \left(\boldsymbol{\omega}' \boldsymbol{\mu} - \mu^p \right) - \gamma_2 \left(\boldsymbol{\omega}' \mathbf{1} - 1 \right)$$

The first derivative equations are (in matrix notation,)

$$rac{\partial \mathcal{L}}{\partial oldsymbol{\omega}'} = \mathbf{\Sigma} oldsymbol{\omega} - \gamma_1 oldsymbol{\mu} - \gamma_2 \mathbf{1}$$

Get the first-order conditions of optimization by setting equal to zero and solve for ω^* :

$$oldsymbol{\omega}^* = \! oldsymbol{\Sigma}^{-1} egin{bmatrix} oldsymbol{\mu} & oldsymbol{1} \end{bmatrix} egin{bmatrix} \gamma_1 \ \gamma_2 \end{bmatrix}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 62/69

Solving the MV problem: portfolios $\omega^{ t t}$ and $\omega^{ t v}$

Rewrite this as

$$\boldsymbol{\omega}^* = \gamma_1 \mathbf{\Sigma}^{-1} \boldsymbol{\mu} + \gamma_2 \mathbf{\Sigma}^{-1} \mathbf{1}$$

which can be rewritten as the sum of two portfolios:

$$oldsymbol{\omega}^* = \!\! \gamma_1 \left(\mathbf{1}' \mathbf{\Sigma}^{-1} oldsymbol{\mu}
ight) oldsymbol{\omega}^{ exttt{t}} + \gamma_2 \left(\mathbf{1}' \mathbf{\Sigma}^{-1} \mathbf{1}
ight) oldsymbol{\omega}^{ exttt{v}}$$

where

$$\omega^\mathtt{t} \equiv rac{1}{\mathbf{1}'\mathbf{\Sigma}^{-1}\mu}\mathbf{\Sigma}^{-1}\mu, \qquad \omega^\mathtt{v} \equiv rac{1}{\mathbf{1}'\mathbf{\Sigma}^{-1}\mathbf{1}}\mathbf{\Sigma}^{-1}\mathbf{1}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 63/69

Solving the MV problem: eliminate γ_2

Note that ω^{t} and ω^{v} are proper portfolios:

$$\left(oldsymbol{\omega}^{\mathtt{t}}
ight)'\mathbf{1}=1, \qquad \left(oldsymbol{\omega}^{\mathtt{v}}
ight)'\mathbf{1}=1$$

Given that $\mathbf{1}'\omega^*=\mathbf{1}'\omega^{\mathsf{t}}=\mathbf{1}'\omega^{\mathsf{v}}=1$, the equation above implies

$$1 = \gamma_1 \left(\mathbf{1}' \mathbf{\Sigma}^{-1} \boldsymbol{\mu} \right) + \gamma_2 \left(\mathbf{1}' \mathbf{\Sigma}^{-1} \mathbf{1} \right)$$

Use this to rewrite the MV vector as

$$\omega^* = \delta \omega^{\mathtt{t}} + (1 - \delta) \omega^{\mathtt{v}}$$

where

$$\delta \equiv \gamma_1 \left(\mathbf{1}' \mathbf{\Sigma}^{-1} \boldsymbol{\mu} \right)$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 64/69

MV formulas

For any MV portfolio ω^* , consider the mean, μ^p and variance σ_p^2 ,

Sub out γ_1 to get δ in terms of μ^p ,

$$\delta = \frac{\mu^{p} - \mu' \omega^{v}}{\mu' \omega^{t} - \mu' \omega^{v}}$$

The return variance, σ_p^2 , is a quadratic function of μ^p ,

$$\sigma_{p}^{2} = \frac{1}{\phi_{0}\phi_{2} - \phi_{1}^{2}} \left[\phi_{0} - 2\phi_{1} \left(\mu^{p} \right) + \phi_{2} \left(\mu^{p} \right)^{2} \right]$$

where the coefficients, ϕ are characterized by

$$\phi_0 = \mu' \mathbf{\Sigma}^{-1} \mu, \qquad \phi_1 = \mu' \mathbf{\Sigma}^{-1} \mathbf{1}, \qquad \phi_2 = \mathbf{1}' \mathbf{\Sigma}^{-1} \mathbf{1}$$

Hendricks, Autumn 2024 FINM 36700: Lecture 1 65/69

Two-fund separation

Consider any three MV portfolios, ω_a , ω_b , ω^p , which must satisfy the following for some δ_a , δ_b , δ_p ,

$$\omega_a = \delta_a \omega^{t} + (1 - \delta_a) \omega^{v}$$

$$\omega_b = \delta_b \omega^{t} + (1 - \delta_b) \omega^{v}$$

$$\omega^{p} = \delta_p \omega^{t} + (1 - \delta_p) \omega^{v}$$

- $ightharpoonup \omega^{t}$ and ω^{v} are not unique in being able to decompose the MV portfolio, ω^{p} .
- lacktriangle Any MV portfolio can be written as a combo of ω_a and ω_b .

$$m{\omega}^{m{p}} = artheta m{\omega}_{m{a}} + (1 - artheta) m{\omega}_{m{b}}, \qquad artheta \equiv rac{\delta_{m{p}} - \delta_{m{b}}}{\delta_{m{a}} - \delta_{m{b}}}$$



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 66/69

Uncorrelated MV portfolios

Using 2-fund separation, convenient to decompose MV portfolios into two orthogonal portfolios.

- For any MV portfolio, $\omega^p \neq \omega^v$, there exists another MV portfolio, ω_o such that ω_o orthogonal to ω^p .
- If ω^p has mean return μ^p , then the orthogonal MV portfolio ω_o has mean return, μ_o , where

$$\mu_o = \frac{\phi_1 \mu^p - \phi_0}{\phi_2 \mu^p - \phi_1}$$

$$\phi_0 = \mu' \mathbf{\Sigma}^{-1} \mu, \qquad \phi_1 = \mu' \mathbf{\Sigma}^{-1} \mathbf{1}, \qquad \phi_2 = \mathbf{1}' \mathbf{\Sigma}^{-1} \mathbf{1}$$



Hendricks, Autumn 2024 FINM 36700: Lecture 1 67/69

Geometry of uncorrelated portfolios


In mean-volatility space, the orthogonal MV portfolio has a simple geometry.

- ▶ Draw the tangent line at the point of some MV portfolio.
- ► Find the value on this tangent line for volatility of zero, (where it hits the vertical axis.)
- The mean return at this point is, μ_o , the mean return of the orthogonal MV portfolio.

See the following figure.



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 68/69



 Hendricks,
 Autumn 2024
 FINM 36700: Lecture 1
 69/69