


# Probabilistic Graphical Models: Problem Set 3 (Solutions)

Svante Linusson, Liam Solus  
KTH Royal Institute of Technology

27 September 2021

1. Consider the DAG  $\mathcal{G} = ([8], E)$  depicted below:




- (a) What is the essential graph of  $\mathcal{G}$ ?  
 (b) How many DAGs are in the Markov equivalence class of  $\mathcal{G}$ ?

---

**Solution:**

**a)** We note that the v-structures are  $1 \rightarrow 2 \leftarrow 3$  and  $3 \rightarrow 6 \leftarrow 4$  so those four edges cannot change direction. Then we notice that  $6 \rightarrow 8$  and  $6 \rightarrow 7$  cannot be reversed since that would cause a new v-structure with the edge  $3 \rightarrow 6$ . The other edges can be given different directions however. The answer is thus



**b)** The edges  $1 - 4 - 5$  can be directed in three ways without creating a v-structure and the edge  $7 - 8$  can be directed either way without creating a v-structure. And there is no risk of creating a cycle in either case. They are independent so the total number is  $2 \cdot 3 = 6$ .

---

2. Explain why the edge  $a \rightarrow b$  in an essential graph  $\mathcal{D}$  cannot be reversed if it is strongly protected.

---

**Solution:**

As explained at the end of lecture 3, we look at the four cases in the definition of strongly protected. Note that they are **induced** subgraphs. For the first two a v-structure is formed or broken if  $a \rightarrow b$  is reversed, which would change the MEC. For the third case it would create a cycle. The fourth case is slightly more complicated. If  $a \rightarrow b$  is reversed then we must direct  $c_1 \rightarrow a$  to avoid a cycle with  $b, a, c_1$  and similarly we must have  $c_2 \rightarrow a$ . But this now creates a v-structure  $c_1 \rightarrow a \leftarrow c_2$ .

Problem Set 3 (Solutions)

3. For a DAG  $\mathcal{G} = (V, E)$ ,  $u, v \in V$ . Prove that if  $u, v$  are not adjacent then for either  $C = \text{pa}_{\mathcal{G}}(u)$  or  $C = \text{pa}_{\mathcal{G}}(v)$ , there is no d-connecting path between  $u$  and  $v$  given  $C$  in  $\mathcal{G}$ . \_\_\_\_\_

**Solution:**

Let first  $C = \text{pa}_{\mathcal{G}}(u)$  and assume that there is a d-connecting path between  $u$  and  $v$ . That path must start from  $u$  with an edge directed away  $u$  and to be d-connecting it must continue to be a directed path until it meets an ancestor of some node in  $C$  but that would create a cycle, so it cannot happen. Thus the path must be a directed path from  $u$  to  $v$ .

If we instead let  $C = \text{pa}_{\mathcal{G}}(v)$ , then we can similarly conclude that the only possible d-connecting path would be a directed path from  $v$  to  $u$ . Since it would create a cycle to have both a directed path from  $u$  to  $v$  and another from  $v$  to  $u$  we can conclude that for at least one of  $C = \text{pa}_{\mathcal{G}}(u)$  or  $C = \text{pa}_{\mathcal{G}}(v)$ , there is no d-connecting path between  $u$  and  $v$  given  $C$  in  $\mathcal{G}$ .

4. Let  $\mathcal{G}$  and  $\mathcal{H}$  be two DAGs on node set  $V$  such that  $\mathcal{G} \leq \mathcal{H}$  (i.e.  $\text{CI}(\mathcal{H}) \subseteq \text{CI}(\mathcal{G})$ ). Prove that if  $\mathcal{G}$  contains the v-structure  $x \rightarrow z \leftarrow y$  then either  $\mathcal{H}$  contains the same v-structure or  $x$  and  $y$  are adjacent in  $\mathcal{H}$ . \_\_\_\_\_

**Solution:**

Suppose this is not the case and  $\mathcal{H}$  does not contain the v-structure and  $X$  and  $Y$  are not adjacent in  $\mathcal{H}$ . From Exercise 3 we can conclude that if there is an edge between two vertices in  $\mathcal{G}$  then they are adjacent also in  $\mathcal{H}$ , thus  $Z$  must be adjacent to both  $X$  and  $Y$  in  $\mathcal{H}$ . By our assumption,  $Z$  is a parent of either  $X$  or  $Y$  in  $\mathcal{H}$ . This implies by Exercise 3, that there exists a conditioning set  $C$  that includes node  $Z$  (and does not include either node  $X$  or node  $Y$ ) such that there is no d-connecting path between  $X$  and  $Y$  given  $C$  in  $\mathcal{H}$ . But the path  $X \rightarrow Z \leftarrow Y$  in  $\mathcal{G}$  is d-connecting given any set that includes  $Z$  (and excludes  $X$  and  $Y$ ), including the set  $C$ , which contradicts the fact that  $\mathcal{G} \leq \mathcal{H}$ .

5. An edge  $i \rightarrow j$  in a DAG  $\mathcal{G} = ([m], E)$  is called **covered** if  $\text{pa}_{\mathcal{G}}(j) = \text{pa}_{\mathcal{G}}(i) \cup \{i\}$ . In this problem we will show that two DAGs  $\mathcal{G} = ([m], E)$  and  $\mathcal{G}' = ([m], E')$  are Markov equivalent if and only if there exists a sequence of DAGs  $\mathcal{G}_1 := \mathcal{G}, \dots, \mathcal{G}_M := \mathcal{G}'$  such that the only difference between  $\mathcal{G}_i$  and  $\mathcal{G}_{i+1}$  for all  $i \in [M - 1]$  is the reversal of a single covered edge.

- (a) Let  $\mathcal{G} = ([m], E)$  be a DAG containing the edge  $i \rightarrow j$  and let  $\mathcal{G}' = ([m], E')$  be the directed graph produced by reversing the edge  $i \rightarrow j$  in  $\mathcal{G}$ . Show that  $\mathcal{G}'$  is a DAG that is Markov equivalent to  $\mathcal{G}$  if and only if  $i \rightarrow j$  is a covered edge in  $\mathcal{G}$ .
- (b) Consider two Markov equivalent DAGs  $\mathcal{G} = ([m], E)$  and  $\mathcal{G}' = ([m], E')$ . Fix a linear extension  $\pi = \pi_1 \dots \pi_m$  of  $\mathcal{G}$  and for  $i \in [m]$  define

$$P_i = \{j \in [m] : j \rightarrow i \in \Delta(\mathcal{G}, \mathcal{G}')\},$$

where

$$\Delta(\mathcal{G}, \mathcal{G}') = \{j \rightarrow i \in E : j \leftarrow i \in E'\}.$$

Let  $k$  be the smallest number such that  $P_{\pi_k} \neq \emptyset$  and let  $s$  be the largest number such that  $\pi_s \in P_{\pi_k}$ . Prove that  $\pi_s \rightarrow \pi_k$  is a covered edge in  $\mathcal{G}$ .

- (c) Prove the theorem stated at the start of the problem.
- (d) Implement an algorithm that takes in a DAG  $\mathcal{G}$  and computes all elements of its Markov equivalence class.

**Solution:**

a) We use the theorem by Verma and Pearl that two DAGs are Markov equivalent (MEQ) iff they have the same skeleton and the same v-structures. Assume first that the graphs are MEQ. If  $\exists x \in \text{pa}_{\mathcal{G}}(j) \setminus (\text{pa}_{\mathcal{G}}(i) \cup \{i\})$  then also  $x \notin \text{ch}_{\mathcal{G}}(i)$  since that would mean that  $i \rightarrow x \rightarrow j \rightarrow i$  is a cycle in  $\mathcal{G}'$ . Since  $\mathcal{G}$  and  $\mathcal{G}'$  have the same skeleton, there is no edge between  $i$  and  $x$  in  $\mathcal{G}$  either so  $i \rightarrow j \leftarrow x$  is a v-structure in  $\mathcal{G}$  which would not be in  $\mathcal{G}'$ . A contradiction and thus no such  $x$  exists. Similarly, if  $\exists x \in \text{pa}_{\mathcal{G}}(i) \setminus \text{pa}_{\mathcal{G}}(j)$  we use the same reasoning with  $\mathcal{G}$  and  $\mathcal{G}'$  interchanged.

The other direction is easier. If  $i \rightarrow j$  is a covered edge in  $\mathcal{G}$  then it cannot be part of a v-structure so neither skeleton nor v-structures will change if we reverse it. Remains to show that  $\mathcal{G}'$  is a DAG. But if we created a cycle including the edges  $x \rightarrow j \rightarrow i$  for some  $x \in \text{pa}_{\mathcal{G}}(j)$ , then it would already have been a cycle in  $\mathcal{G}$  by the edge  $x \rightarrow i$ , which we know it is not.

**b)** Recall that we have defined the edges to point from the smaller number to the larger number in a linear extension. Since  $\mathcal{G}$  and  $\mathcal{G}'$  are MEQ they have the same skeleton. First, if there is a node  $x$ ,  $\pi_x < \pi_s$  with  $\pi_x \rightarrow \pi_s$  in  $\mathcal{G}$  then by choice of  $k$ ,  $\pi_x \rightarrow \pi_s$  also in  $\mathcal{G}'$ . That means there has to be an edge  $\pi_x \rightarrow \pi_k$  in  $\mathcal{G}'$  to avoid making  $\pi_x \rightarrow \pi_s \leftarrow \pi_k$  a v-structure. By choice of  $s$   $\pi_x \rightarrow \pi_k$  also in  $\mathcal{G}$ . Second, if there is a node  $x$ ,  $\pi_x < \pi_s$  with  $\pi_x \rightarrow \pi_k$  in  $\mathcal{G}$ , we would with a similar reasoning obtain that  $\pi_x \rightarrow \pi_s$  in both graphs. Finally, if there is a node  $x$ ,  $\pi_s \pi_x < \pi_k$  with  $\pi_x \rightarrow \pi_k$  in  $\mathcal{G}$ , then by choice of  $s$ ,  $\pi_x \rightarrow \pi_k$  also in  $\mathcal{G}'$ . To not have different v-structures both graphs must have an edge  $\pi_s \rightarrow \pi_x$ , but this would create a cycle in  $\mathcal{G}'$ , a contradiction. Thus  $\pi_s \rightarrow \pi_k$  is covered.

**c)** Given two MEQ graphs  $\mathcal{G}$ ,  $\mathcal{G}'$  having  $n$  edges with different orientation. Part b) shows how to find a covered edge and a) shows we can reverse it and obtain new DAG still MEQ and having different orientation for  $n - 1$  edges compared to  $\mathcal{G}'$ . Induction over  $n$  proves that direction. The other direction follows from a).