

Список литературы

- [1] G. Arratia, K. A. González-Rodríguez, and C. Hernández-Guerrero, *Fossil Record*, **21**(1), 93–107 (2018). A new Pachyrhizodontid Fish (Actinopterygii, Teleostei) from the Muhi Quarry (Albian-Cenomanian), Hidalgo, Mexico.
- [2] L. Agassiz, *Recherches sur les Poissons Fossiles. Tome II*, 1833.
- [3] R. Doiuchi and T. Nakabo, *Ichthyological Research*, **52**, 132–151 (2005). The *Sphyraena obtusata* group (Perciformes: Sphyraenidae) with a description of a new species from southern Japan.
- [4] A. J. Mainwaring, page 162 (1978). Anatomical and Systematic review of the Pachycormidae, a family of Mesozoic fossil fishes.
- [5] M. R. Sánchez-Villagra, R. J. Asher, A. D. Rincón, A. A. Carlini, P. A. Meylan, and R. W. Purdy, *Special Papers in Palaeontology*, **71**, 105–112 (2004). New faunal reports for the cerro La Cruz locality (Lower Miocene), north-eastern Venezuela.
- [6] R. Betancur-R, E. O. Wiley, G. Arratia, A. Acero, N. Bailly, M. Miya, G. Lecointre, and G. Ortí, *BMC Evolutionary Biology*, **17**(162), 1–40 (2017). Phylogenetic classification of bony fishes.
- [7] H. W. Fowler, *Proceedings of the Academy of Natural Sciences of Philadelphia*, **55**(1903), 743–752 (1903). New and little known Mugilidae and Sphyraenidae.
- [8] R. Brzobohatý and D. Nolf, *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre*, **81**, 279–295 (2011). Fish otoliths from the middle Eocene (Bartonian) of Yebra de Basa, province of Huesca, Spain.
- [9] E. D. Cope, *Proceedings of the American Philosophical Society*, **14**(94), 361–364 (1875). Synopsis of the Vertebrata of the Miocene of Cumberland County, New Jersey.
- [10] J. D. Bryant, *Journal of Vertebrate Paleontology*, **11**(4), 472–489 (1991). New early Barstovian (Middle Miocene) vertebrates from the upper Torreya Formation, Eastern Florida Panhandle.
- [11] K. Okonechnikov, O. Golosova, M. Fursov, A. Varlamov, Y. Vaskin, I. Efremov, O. G. German Grehov, D. Kandrov, K. Rasputin, M. Syabro, and T. Tleukenov, *Bioinformatics*, **28**, 1166–1167 (2012). Unipro UGENE: A unified bioinformatics toolkit.
- [12] C. Arambourg, *Notes et Mémoires sur le Moyen-Orient*, **3**, 1–210 (1966). Les poissons oligocènes de l'Iran.
- [13] J. Amalfitano, L. Giusberti, E. Fornaciari, and G. Carnevale, *Rivista Italiana di Paleontologia e Stratigrafia*, **123**(3), 475–485 (2017). A reappraisal of the Italian record of the cretaceous Pachycormid Fish *Protosphyraena* Leidy, 1857.
- [14] L. Dollo and R. Storms, *Zoologischer Anzeiger*, **11**, 265–267 (1888). Sur les Téléostéens du Rupélien.
- [15] F. Ronquist, M. Teslenko, P. Van Der Mark, D. L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M. A. Suchard, and J. P. Huelsenbeck, *Systematic Biology*, **61**(3), 539–542 (2012). MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space.
- [16] R Core Development Team, R: A language and environment for statistical computing, 2017.
- [17] A. F. Bannikov, *Iskopaemye pozvonochnye Rossii i sopredel'nykh stran. Iskopaemye kolyucheperye ryby (Teleostei, Acanthopterygii)*, GEOS, Moscow, 2010.
- [18] J. Leidy, *Proceedings of the Academy of Natural Sciences of Philadelphia*, **8**, 220–221 (1856). Notice of remains of extinct vertebrated animals of New-Jersey, collected by Prof. Cook of the State Geological Survey under the direction of Dr. W. Kitchell.
- [19] W. v. W. Weiler, *Geologica Hungarica. Series Palaeontologica*, **15**(1-30) (1938). Neue Untersuchungen an Mitteloligozänen Fischen Ungarns.

- [20] D. A. Bone, J. A. Todd, and S. Tracey, *Tertiary Research*, **12**(3-4), 131–137 (1991). Fossils from the Bracklesham Group exposed in the M27 Motorway excavations, Southampton, Hampshire.
- [21] FAO-FIGIS in *A World Overview of Species of Interest to Fisheries*; FAO, Rome, 2005; pages 1–3.
- [22] R. S. Rana, *Journal of the Palaeontological Society of India*, **35**, 105–120 (1990). Palaeontology and palaeoecology of the intertrappean (Cretaceous-Tertiary transition) beds of the Peninsular India.
- [23] J. Leidy, *Proceedings of the Academy of Natural Sciences of Philadelphia*, **7**, 395–397 (1855). Indications of twelve species of fossil Fishes.
- [24] J. R. Grubich, A. N. Rice, and M. W. Westneat, *Zoology*, jan (2008). Functional morphology of bite mechanics in the Great Barracuda (*Sphyraena barracuda*).
- [25] D. Bardack, *University of Kansas Paleontological Contributions*, **10**, 1–86 (1969). Anatomy and evolution of Chirocentrid fishes.
- [26] E. Casier, *Mémoires du Musée Royal d’Histoire Naturelle de Belgique*, **104**, 1–267 (1946). La faune ichthyologique de l’Yprésien de la Belgique.
- [27] O. Mateus, P. M. Callapez, M. J. Polcyn, A. S. Schulp, A. O. Gonçalves, and L. L. Jacobs, *Biodiversity of Angola*, pages 53–76 (2019). The Fossil Record of Biodiversity in Angola Through Time: A Paleontological Perspective.
- [28] M. Friedman and G. Carnevale, *Journal of the Geological Society*, **175**(4), 569–579 (2018). The Bolca Lagerstätten: shallow marine life in the Eocene.
- [29] K. A. Monsch *The Phylogeny of the Scombroid Fishes* PhD thesis, (2000).
- [30] A. F. Bannikov, *Studie Ricerche sui Giacimenti Terziari di Bolca*, **9**, 65–76 (2008). Revision of the Atheriniform fish genera *Rhamphognathus* Agassiz and *Mesogaster* Agassiz (Teleostei) From the Eocene of Bolca, northern Italy.
- [31] J. S. Nelson, T. Grande, and M. V. H. Wilson, *Fishes of the World*, John Wiley & Sons, New Jersey, 5 ed., 2016.
- [32] M. A. Pastore, *Journal of the Marine Biological Association of the United Kingdom*, **89**(6), 1299–1303 (2009). *Sphyraena intermedia* sp. nov. (Pisces: Sphyraenidae): a potential new species of barracuda identified from the central Mediterranean Sea.
- [33] P. L. Forey, *Bulletin of the British Museum (Natural History)*, **28**(2), 123–204 (1977). The osteology of *Notelops* Woodward, *Rhacolepis* Agassiz and *Pachyrhizodus* Dixon (Pisces: Teleostei).
- [34] G. Mas, *Bolleti de la Societat d’Historia Natural de les Balears*, **43**, 39–61 (2000). Ictiofauna del Pliocè mitjà-superior de la conca sedimentària de Palma (Illes Balears, Mediterrània Occidental). Implicacions paleoambientals.
- [35] D. Nolf, *Handbook of Paleoichthyology. Otolithi Piscium*, Gustav Fischer Verlag, 1985.
- [36] I. Hays, *Transactions of the American Philosophical Society*, **3**, 471–477 (1830). Description of a fragment of the head of a new fossil animal, discovered in a Marl Pit, near Moorestown, New Jersey.
- [37] J. D. Carrillo-Briceño, A. E. Reyes-Cespedes, R. Salas-Gismondi, and R. Sánchez, *Swiss Journal of Palaeontology*, **0123456789** (2018). A new vertebrate continental assemblage from the Tortonian of Venezuela.
- [38] R. Fricke, M. Kulbick, and L. Wantiez, *Stuttgarter Beiträge zur Naturkunde A, New Series*, **4**, 341–463 (2011). Checklist of the fishes of New Caledonia, and their distribution in the Southwest Pacific Ocean (Pisces).

- [39] A. S. Woodward, *Catalogue of the Fossil Fishes in the British Museum (Natural History). Part III*, Taylor & Francis, London, 1895.
- [40] P. Artedi, *Petri Artedi Angermannia-Sueci synonymia nominum piscium fere omnium;... Ichthyologiae pars IV. Editio II. Grypeswaldiae*, 1793.
- [41] R. Fricke, W. N. Eschmeyer, and R. van der Laan, Eschmeyer’s catalog of Fishes: Genera, species, references, 2019.
- [42] F. Fanti, D. Minelli, G. L. Conte, and T. Miyashita, *Zoological Letters*, **2**, 2–18 (2016). An exceptionally preserved Eocene Shark and the rise of modern predator–prey interactions in the coral reef food web.
- [43] D. P. de Sylva, *Studies in Tropical Oceanography*, **1**, 1–179 (1963). Systematics and life history of the great barracuda *Sphyaena barracuda* (Walbaum).
- [44] F. T. W. Dames, *Sitzungsber . d . kgl . pr . Akad . d . Wiss. zu Berlin*, **6**, 129–135 (1883). Über eine tertiäre Wirbelthier-fauna von der westlichen Insel Birket-el-Qurun im Fajum (Aegypten).
- [45] M. E. Páramo-Fonseca, *Bolteín Geológico Ingeominas*, **39**, 47–83 (2001). Los peces de la familia Pachyrhizodontidae (Teleostei) del Turoniano del Valle Superior del Magdalena.
- [46] G. P. Whitley, *Australian Zoologist*, **11**(2), 129–150 (1947). New sharks and fishes from Western Australia. Part 3.
- [47] T. A. Deméré, M. A. Roeder, R. M. Chandler, and J. A. Minch in *Miocene and Cretaceous Depositional Environments, Northwestern Baja California, Mexico*, J. A. Minch and J. R. Ashby, Eds.; Pacific Section AAPG, Baja California, 1984; pages 47–56.
- [48] J. R. Bourque in *Morphology and Evolution of Turtles*; Springer Sciences+Business Media, Dordrecht, 2013; pages 459–475.
- [49] F. Santini, G. Carnevale, and L. Sorenson, *Italian Journal of Zoology*, **82**(1), 133–142 (2015). First timetree of Sphyraenidae (Percomorpha) reveals a Middle Eocene crown age and an Oligo–Miocene radiation of Barracudas.
- [50] D. Darriba, G. L. Taboada, R. Doallo, and D. Posada, *Nature Methods*, aug (2012). jModelTest 2: more models, new heuristics and parallel computing.
- [51] W. F. Rapp, *Journal of Paleontology*, **20**(5), 510–513 (1946). Check list of the fossil fishes of New Jersey.
- [52] S. Marsili, G. Carnevale, E. Danese, G. Bianucci, and W. Landini, *Annales de Paléontologie*, jan (2007). Early Miocene vertebrates from Montagna della Maiella, Italy.
- [53] E. Casier, *Bulletin du Musée Royal d’Histoire Naturelle de Belgique*, **20**(23), 1–8 (1944). Contributions a l’etude des poissons fossiles de la Belgique. VII. Morphologie du dentaire de *Sphyraenodus lerichei* Casier.
- [54] B. Huyghebaert and D. Nolf, *Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie*, **16**(4), 139–170 (1979). on fish-otoliths, published since 1968.
- [55] W. E. Bemis, A. Giuliano, and B. McGuire, *Zoology*, **108**(4), 317–327 (2005). Structure, attachment, replacement and growth of teeth in bluefish, *Pomatomus saltatrix* (Linnaeus, 1766), a teleost with deeply socketed teeth.
- [56] I. Nakamura, *FAO Species Catalogue. Volume 5. Billfishes of the World*, Vol. 5, FAO, Rome, 1985.
- [57] E. T. Newton, *Quarterly Journal of the Geological Society*, **33**(1-4), 505–523 (1877). On the remains of *Hypsodon*, *Portheus*, and *Ichthyodectes* from British Cretaceous strata, with descriptions of new species.

- [58] F. Bassani, *Atti della Reale Accademia delle Scienze Fisiche e Matematiche*, **3**(2), 1–100 (1889). Ricerche sui pesci fossili di Chiavon (Strati di Sotzka - Miocene Inferiore).
- [59] A. S. Woodward, *Catalogue of the Fossil Fishes in the British Museum (Natural History). Part IV*, Taylor & Francis, London, 1901.
- [60] J. Leidy, *Journal of the Academy of Natural Sciences*, **8**(3), 209–261 (1877). Description of vertebrate remains chiefly from the phosphate beds of South Carolina.
- [61] E. I. White, *Bulletin of the Geological Survey of Nigeria*, **10**, 1–82 (1926). Eocene Fishes from Nigeria.
- [62] A. J. W. Hendy, D. S. Jones, F. Moreno, V. Zapata, and C. A. Jaramillo, *Swiss Journal of Palaeontology*, pages 1–31 (2015). Neogene molluscs, shallow marine paleoenvironments, and chronostratigraphy of the Guajira Peninsula, Colombia.
- [63] K. Katoh and D. M. Standley, *Molecular Biology and Evolution*, **30**(4), 772–780 (2013). MAFFT multiple sequence alignment software version 7: Improvements in performance and usability.
- [64] D. Nolf, *Bulletin de la Societe Belge de Geologie, Paleontologie et Hydrologie*, **81**(1-2), 95–109 (1972). Deuxième note sur les Téléostéens des sabens de Lede (Éocène Belge).
- [65] H. Senou in *The Living Marine Resources of the Western Central Pacific. Volume 6*, K. E. Carpenter and V. H. Niem, Eds.; FAO, Rome, 2001; pages 3685–3697.
- [66] D. D. Gillette, *Journal of Vertebrate Paleontology*, **4**(2), 172–186 (1984). A marine ichthyofauna from the Miocene of Panama, and the Tertiary Caribbean Faunal Province.
- [67] A. S. Woodward, *Catalogue of the Fossil Fishes in the British Museum (Natural History). Part I*, Taylor & Francis, London, 1889.
- [68] L. Agassiz, *Nomina Systematica Generum Piscium, tam Viventum Quam Fossilum*, 1846.
- [69] L. Agassiz, *Recherches sur les Poisson Fossiles. Tome III*, 1833.
- [70] C. E. Ray, A. Wetmore, D. H. Dunkle, and P. Drez, *Smithsonian Miscellaneous Collections*, **153**(3), 1–25 (1968). Fossil vertebrates from the marine Pleistocene of southeastern Virginia.
- [71] F. Quillévéré, E. Koskeridou, J.-J. Cornée, P. Moissette, A. Girone, and K. Agiadi, *Quaternary Science Reviews*, **196**, 80–99 (2018). Pleistocene marine fish invasions and paleoenvironmental reconstructions in the eastern Mediterranean.
- [72] O. Schultz, R. Brzobohatý, and O. Kroupa, *Annalen des Naturhistorischen Museums in Wien, Serie A*, **112**, 489–506 (2010). Fish teeth from the Middle Miocene of Kienberg at Mikulov, Czech Republic, Vienna Basin.
- [73] S. Díaz-Franco and R. Rojas-Consuegra, *Solenodon*, **8**, 124–129 (2009). Dientes fósiles de *Sphyaena* (Perciformes: Sphyaenidae) en el Terciario de Cuba Occidental.
- [74] J. Böhm in *Die Diamantenwüste Sudwestafrikasfrikas. Vol II*, E. Kaiser, Ed.; 1926; pages 55–87.
- [75] J. W. Westgate, *Journal of Vertebrate Paleontology*, **4**(4), 536–546 (1984). Lower vertebrates from the late Eocene Crow Creek local fauna, St. Francis County, Arkansas.
- [76] M. D. Gottfried, K. E. Samonds, S. A. Ostrowski, T. H. Andrianavalona, and T. N. Ramihangihajason, *PLoS ONE*, **12**(5), 1–9 (2017). New evidence indicates the presence of Barracuda (Sphyaenidae) and supports a tropical marine environment in the Miocene of Madagascar.
- [77] ICZN, *International Code of Zoological Nomenclature*, The International Trust for Zoological Nomenclature, London, 4th ed., 1999.
- [78] R. Harlan, *Journal fo the Academy of Natural Sciences of Philadelphia*, **3**, 331–338 (1824). On a new fossil genus, of the order Enalio Sauri (of Conybeare): and on a new species of Ichthyosaurus.

- [79] E. Casier, *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, **20**(11), 11–15 (1944). Contributions à l'étude des Poissons fossiles de la Belgique. VI. Sur le *Sphyaenodus* de l'Eocene e sur la présence d'un Sphyaenidé dans le Bruxellien (Lutétien inférieur).
- [80] A. S. Woodward, *Catalogue of the Fossil Fishes in the British Museum (Natural History). Part II*, Taylor & Francis, London, 1891.
- [81] STRINGER, GARY L., Department of Ge, *AAPG Bulletin*, **85** (2003). Paleoenvironmental Interpretations Based on Vertebrate Fossil Assemblages: An Example of their Utilization in the Gulf Coast.
- [82] M. E. Páramo-Fonseca, *Comptes Rendus de l'Academie de Sciences - Serie IIa: Sciences de la Terre et des Planetes*, **325**(2), 147–150 (1997). *Bachea huilensis* nov. gen., nov. sp., premier Tselfatioidei (Teleostei) de Colombie.
- [83] F. Chapman, *Transactions and Proceedings of the Royal Society of New Zealand*, **64**, 117–121 (1935). Descriptions of fossil fish from New Zealand.
- [84] D. P. de Sylva and F. Williams in *Smiths' Sea Fishes*, M. M. Smith and P. C. Heemstra, Eds.; Macmillan South Africa, Johannesburg, 1986; pages 721–726.
- [85] C. E. Ray and D. J. Bohaska, *Geology and Paleontology of the Lee Creek Mine, North Carolina, III*, number 90, smithsonia ed., 2001.
- [86] C. Patterson, *Bulletin of Marine Science*, **52**(1), 29–59 (1993). An overview of the early fossil record of Acanthomorphs.
- [87] D. Bardack and G. Sprinkle, *Fieldiana Geology*, **16**, 297–340 (1969). Morphology and relationships of saurocephalid fishes.
- [88] K. A. Monsch, *Transactions of the Royal Society of Edinburgh: Earth Sciences*, **95**(November 2016), 445–489 (2005). Revision of the scombroid fishes from the Cenozoic of England.
- [89] L. Agassiz, *Recherches sur les Poissons Fossiles. Tome IV*, 1833.
- [90] F. Moreno, A. J. W. Hendy, L. Quiroz, N. Hoyos, D. S. Jones, V. Zapata, S. Zapata, G. A. Ballen, E. Cadena, A. L. Cárdenas, J. D. Carrillo-Briceño, J. D. Carrillo, D. Delgado-Sierra, J. Escobar, J. I. Martínez, C. Martínez, C. Montes, J. Moreno, N. Pérez, R. Sánchez, C. Suárez, M. C. Vallejo-Pareja, and C. A. Jaramillo, *Swiss Journal of Palaeontology*, **134**, 5–43 (2015). Revised stratigraphy of Neogene strata in the Cocinetas Basin, La Guajira, Colombia.
- [91] L. Agassiz, *Recherches sur les Poissons Fossiles. Tome I*, 1833.
- [92] R. Van der Laan, *European Journal of Taxonomy*, (466), 1–167 (2018). Family-group names of fossil fishes.
- [93] S. E. Meek and R. G. Newland, *Proceedings of the Academy of Natural Sciences of Philadelphia*, **36**, 67–75 (1884). A review of the American species of the genus *Sphyaena*.
- [94] G. L. Stringer, S. Q. Breard, and M. Kontrovitz, *Gulf Coast Association of Geological Societies Transactions*, **LI**, 321–328 (2001). Biostratigraphy and Paleoecology of Diagnostic Invertebrates and Vertebrates from the Type Locality of the Oligocene Rosefiled Marl Beds, Louisiana.
- [95] E. D. Cope, *Transactions of the American Philosophical Society*, **14**, 1–252 (1869). Synopsis of the Batrachia and Reptilia of North America. Part I.
- [96] A. A. Switchenska in *Ocherki po Filogenii i Sistematike Iskopaemykh Ryb I Beschelyustnykh*; 1968; pages 157–161.
- [97] L. Agassiz, *Recherches sur les Poisson Fossiles. Tome V*, 1843.
- [98] L. W. Viñola-López, R. Rojas-Consuegra, and O. Jiménez-Vásquez, *Novitates Caribaea*, **11**, 89–94 (2017). Nuevos registros de *Sphyaena* (Perciformes: Sphyaenidae) para el Neógeno de Cuba y La Española.

- [99] NCBI, *Entrez Programming Utilities Help*, number Md, NCBI, Bethesda, 2018.
- [100] L. Taverne and B. Chanet, *Geodiversitas*, **22**(1), 23–34 (2000). *Faugichthys loryi* n. gen., n. sp. (Teleostei, Ichthyodectiformes) de l’Albien terminal (Crétacé inférieur marin) du vallon de la Fauge (Isère, France) et considérations sur la phylogénie des Ichthyodectidae.
- [101] J. L. B. Smith, *Ichthyological Bulletin of the Department of Ichthyology of Rhodes University*, **3**, 37–46 (1956). The fishes of the family Sphyaenidae in the western Indian Ocean.
- [102] V. d. A. Távora, A. A. R. dos Santos, and R. N. Araújo, *Boletim do Museu Paraense Emilio Goeldi Ciencias Naturais*, **5**(2), 207–224 (2010). Localidades fossilíferas da Formação Pirabas (Mioceno Inferior).