## Список литературы

- [1] Arratia, G., González-Rodríguez, K. A., and Hernández-Guerrero, C. (2018) A new Pachyrhizodontid Fish (Actinopterygii, Teleostei) from the Muhi Quarry (Albian-Cenomanian), Hidalgo, Mexico. Fossil Record, 21, 93–107.
- [2] Agassiz, L. (1833) Recherches sur les Poissons Fossiles. Tome II.
- [3] Doiuchi, R. and Nakabo, T. (2005) The *Sphyraena obtusata* group (Perciformes: Sphyraenidae) with a description of a new species from southern Japan. *Ichthyological Research*, **52**, 132–151.
- [4] Mainwaring, A. J. (1978) Anatomical and Systematic review of the Pachycormidae, a family of Mesozoic fossil fishes. p. 162.
- [5] Sánchez-Villagra, M. R., Asher, R. J., Rincón, A. D., Carlini, A. A., Meylan, P. A., and Purdy, R. W. (2004) New faunal reports for the cerro La Cruz locality (Lower Miocene), north-eastern Venezuela. Special Papers in Palaeontology, 71, 105–112.
- [6] Betancur-R, R., Wiley, E. O., Arratia, G., Acero, A., Bailly, N., Miya, M., Lecointre, G., and Ortí, G. (2017) Phylogenetic classification of bony fishes. BMC Evolutionary Biology, 17, 1–40.
- [7] Fowler, H. W. (1903) New and little known Mugilidae and Sphyraenidae. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **55**, 743–752.
- [8] Brzobohatý, R. and Nolf, D. (2011) Fish otoliths from the middle Eocene (Bartonian) of Yebra de Basa, province of Huesca, Spain. Bulletin de l'Institut Royal des Sciences Naturelles de Belqique, Sciences de la Terre, 81, 279–295.
- [9] Cope, E. D. (1875) Synopsis of the Vertebrata of the Miocene of Cumberland County, New Jersey. *Proceedings of the American Philosophical Society*, **14**, 361–364.
- [10] Bryant, J. D. (1991) New early Barstovian (Middle Miocene) vertebrates from the upper Torreya Formation, Eastern Florida Panhandle. *Journal of Vertebrate Paleontology*, 11, 472–489.
- [11] Okonechnikov, K., et al. (2012) Unipro UGENE: A unified bioinformatics toolkit. *Bioinformatics*, **28**, 1166–1167.
- [12] Arambourg, C. (1966) Les poissons oligocènes de l'Iran. *Notes et Mémoires sur le Moyen-Orient*, **3**, 1–210.
- [13] Amalfitano, J., Giusberti, L., Fornaciari, E., and Carnevale, G. (2017) A reappraisal of the Italian record of the cretaceous Pachycormid Fish Protosphyraena Leidy, 1857. Rivista Italiana di Paleontologia e Stratigrafia, 123, 475–485.
- [14] Dollo, L. and Storms, R. (1888) Sur les Téléostéens du Rupélien. Zoologischer Anzeiger, 11, 265–267.
- [15] Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D. L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M. A., and Huelsenbeck, J. P. (2012) MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space. Systematic Biology, 61, 539–542.
- [16] R Core Development Team (2017), R: A language and environment for statistical computing.
- [17] Bannikov, A. F. (2010) Iskopaemye pozvonochnye Rossii i sopredel'nykh stran. Iskopaemye kolyucheperye ryby (Teleostei, Acanthopterigii). GEOS.
- [18] Leidy, J. (1856) Notice of remains of extinct vertebrated animals of New-Jersey, collected by Prof. Cook of the State Geological Survey under the direction of Dr. W. Kitchell. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 8, 220–221.
- [19] Weiler, W. v. W. (1938) Neue Untersuchungen an Mitteloligozänen Fischen Ungarns. Geologica Hungarica. Series Palaeontologica, 15.

- [20] Bone, D. A., Todd, J. A., and Tracey, S. (1991) Fossils from the Bracklesham Group exposed in the M27 Motorway excavations, Southampton, Hampshire. *Tertiary Research*, 12, 131–137.
- [21] FAO-FIGIS (2005) Trachurus trachurus. A World Overview of Species of Interest to Fisheries, pp. 1–3, FAO.
- [22] Rana, R. S. (1990) Palaeontology and palaeoecology of the intertrappean (Cretaceous-Tertiary transition) beds of the Peninsular India. *Journal of the Palaeontological Society of India*, **35**, 105–120.
- [23] Leidy, J. (1855) Indications of twelve species of fossil Fishes. Proceedings of the Academy of Natural Sciencies of Philadelphia, 7, 395–397.
- [24] Grubich, J. R., Rice, A. N., and Westneat, M. W. (2008) Functional morphology of bite mechanics in the Great Barracuda (*Sphyraena barracuda*). *Zoology*, **111**, 16–29.
- [25] Bardack, D. (1969) Anatomy and evolution of Chirocentrid fishes. *University of Kansas Paleontological Contributions*, **10**, 1–86.
- [26] Casier, E. (1946) La faune ichthyologique de l'Yprésien de la Belgique. Mémoires du Musée Royal d'Histoire Naturelle de Belgique, 104, 1–267.
- [27] Mateus, O., Callapez, P. M., Polcyn, M. J., Schulp, A. S., Gonçalves, A. O., and Jacobs, L. L. (2019) The Fossil Record of Biodiversity in Angola Through Time: A Paleontological Perspective. *Biodiversity of Angola*, pp. 53–76.
- [28] Friedman, M. and Carnevale, G. (2018) The Bolca Lagerstätten: shallow marine life in the Eocene. Journal of the Geological Society, 175, 569–579.
- [29] Monsch, K. A. (2000) The Phylogeny of the Scombroid Fishes. Ph.D. thesis.
- [30] Bannikov, A. F. (2008) Revision of the Atheriniform fish genera Rhamphognathus Agassiz and Mesogaster Agassiz (Teleostei) From the Eocene of Bolca, northern Italy. Studie Ricerche sui Giacimenti Terziari di Bolca, 9, 65–76.
- [31] Nelson, J. S., Grande, T., and Wilson, M. V. H. (2016) Fishes of the World. John Wiley & Sons, 5 edn.
- [32] Pastore, M. A. (2009) Sphyraena intermedia sp. nov. (Pisces: Sphyraenidae): a potential new species of barracuda identified from the central Mediterranean Sea. Journal of the Marine Biological Association of the United Kingdom, 89, 1299–1303.
- [33] Forey, P. L. (1977) The osteology of *Notelops* Woodward, *Rhacolepis* Agassiz and *Pachyrhizodus* Dixon (Pisces: Teleostei). *Bulletin of the British Museum (Natural History)*, **28**, 123–204.
- [34] Mas, G. (2000) Ictiofauna del Pliocè mitjà-superior de la conca sedimentària de Palma (Illes Balears, Mediterrània Occidental). Implicacions paleoambientals. *Bolleti de la Societat d'Historia Natural de les Balears*, **43**, 39–61.
- [35] Nolf, D. (1985) Handbook of Paleoichthyology. Otolithi Piscium. Gustav Fischer Verlag.
- [36] Hays, I. (1830) Description of a fragment of the head of a new fossil animal, discovered in a Marl Pit, near Moorestown, New Jersey. Transactions of the American Philosophical Society, 3, 471–477.
- [37] Carrillo-Briceño, J. D., Reyes-Cespedes, A. E., Salas-Gismondi, R., and Sánchez, R. (2018) A new vertebrate continental assemblage from the Tortonian of Venezuela. Swiss Journal of Palaeontology, 0123456789.
- [38] Fricke, R., Kulbick, M., and Wantiez, L. (2011) Checklist of the fishes of New Caledonia, and their distribution in the Southwest Pacific Ocean (Pisces). Stuttgarter Beiträge zur Naturkunde A, New Series, 4, 341–463.

- [39] Woodward, A. S. (1895) Catalogue of the Fossil Fishes in the British Museum (Natural History).

  Part III. Taylor & Francis.
- [40] Artedi, P. (1793) Petri Artedi Angermannia-Sueci synonymia nominum piscium fere omnium;... Ichthyologiae pars IV. Editio II. Grypeswaldiae.
- [41] Fricke, R., Eschmeyer, W. N., and van der Laan, R. (2019), Eschmeyer's catalog of Fishes: Genera, species, references.
- [42] Fanti, F., Minelli, D., Conte, G. L., and Miyashita, T. (2016) An exceptionally preserved Eocene Shark and the rise of modern predator–prey interactions in the coral reef food web. *Zoological Letters*, 2, 2–18.
- [43] de Sylva, D. P. (1963) Systematics and life history of the great barracuda Sphyraena barracuda (Walbaum). Studies in Tropical Oceanography, 1, 1–179.
- [44] Dames, F. T. W. (1883) Über eine tertiäre Wirbelthier-fauna von der westlichen Insel Birket-el-Qurun im Fajum (Aegypten). Sitzungsber . d . kgl . pr . Akad . d . Wiss. zu Berlin, 6, 129–135.
- [45] Páramo-Fonseca, M. E. (2001) Los peces de la familia Pachyrhizodontidae (Teleostei) del Turoniano del Valle Superior del Magdalena. *Bolteín Geológico Ingeominas*, **39**, 47–83.
- [46] Whitley, G. P. (1947) New sharks and fishes from Western Australia. Part 3. Australian Zoologist, 11, 129–150.
- [47] Deméré, T. A., Roeder, M. A., Chandler, R. M., and Minch, J. A. (1984) Paleontology of the middle Miocene Los Indios Member of the Rosarito Formation, northwestern Baja California, Mexico. Minch, J. A. and Ashby, J. R. (eds.), Miocene and Cretaceous Depositional Environments, Northwestern Baja California, Mexico, pp. 47–56, Pacific Section AAPG.
- [48] Bourque, J. R. (2013) Fossil Kinosternidae from the Oligocene and Miocene of Florida, USA. *Morphology and Evolution of Turtles*, pp. 459–475, Springer Sciences+Business Media.
- [49] Santini, F., Carnevale, G., and Sorenson, L. (2015) First timetree of Sphyraenidae (Percomorpha) reveals a Middle Eocene crown age and an Oligo-Miocene radiation of Barracudas. *Italian Journal of Zoology*, 82, 133–142.
- [50] Darriba, D., Taboada, G. L., Doallo, R., and Posada, D. (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, **9**, 772–772.
- [51] Rapp, W. F. (1946) Check list of the fossil fishes of New Jersey. Journal of Paleontology, 20, 510–513.
- [52] Marsili, S., Carnevale, G., Danese, E., Bianucci, G., and Landini, W. (2007) Early Miocene vertebrates from Montagna della Maiella, Italy. *Annales de Paléontologie*, 93, 27–66.
- [53] Casier, E. (1944) Contributions a l'étude des poissons fossiles de la Belgique. VII. Morphologie du dentaire de Sphyraenodus lerichei Casier. Bulletin du Museé Royal d'Histoire Naturelle de Belgique, 20, 1–8.
- [54] Huyghebaert, B. and Nolf, D. (1979) on fish-otoliths, published since 1968. Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie, 16, 139–170.
- [55] Bemis, W. E., Giuliano, A., and McGuire, B. (2005) Structure, attachment, replacement and growth of teeth in bluefish, *Pomatomus saltatrix* (Linnaeus, 1766), a teleost with deeply socketed teeth. *Zoology*, **108**, 317–327.
- [56] Nakamura, I. (1985) FAO Species Catalogue. Volume 5. Billifishes of the World, vol. 5. FAO.
- [57] Newton, E. T. (1877) On the remains of Hypsodon, Portheus, and Ichthyodectes from British Cretaceous strata, with descriptions of new species. Quarterly Journal of the Geological Society, 33, 505–523.

- [58] Bassani, F. (1889) Ricerche sui pesci fossili di Chiavon (Strati di Sotzka Miocene Inferiore). Atti della Reale Accademia delle Scienze Fisiche e Matematiche, 3, 1–100.
- [59] Woodward, A. S. (1901) Catalogue of the Fossil Fishes in the British Museum (Natural History). Part IV. Taylor & Francis.
- [60] Leidy, J. (1877) Description of vertebrate remains chiefly from the phosphate beds of South Carolina. Journal of the Academy of Natural Sciences, 8, 209–261.
- [61] White, E. I. (1926) Eocene Fishes from Nigeria. Bulletin of the Geological Survey of Nigeria, 10, 1–82.
- [62] Hendy, A. J. W., Jones, D. S., Moreno, F., Zapata, V., and Jaramillo, C. A. (2015) Neogene molluscs, shallow marine paleoenvironments, and chronostratigraphy of the Guajira Peninsula, Colombia. Swiss Journal of Palaeontology, pp. 1–31.
- [63] Katoh, K. and Standley, D. M. (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution, 30, 772–780.
- [64] Nolf, D. (1972) Deuxième note sur les Téléostéens des sabens de Lede (Éocène Belge). Bulletin de la Societe Belge de Geologie, Paleontologie et Hydrologie, 81, 95–109.
- [65] Senou, H. (2001) Sphyraenidae. Carpenter, K. E. and Niem, V. H. (eds.), The Living Marine Resources of the Western Central Pacific. Volume 6, pp. 3685–3697, FAO.
- [66] Gillette, D. D. (1984) A marine ichthyofauna from the Miocene of Panama, and the Tertiary Caribbean Faunal Province. *Journal of Vertebrate Paleontology*, 4, 172–186.
- [67] Woodward, A. S. (1889) Catalogue of the Fossil Fishes in the British Museum (Natural History).

  Part I. Taylor & Francis.
- [68] Agassiz, L. (1846) Nomina Systematica Generum Piscium, tam Viventum Quam Fossilum.
- [69] Agassiz, L. (1833) Recherches sur les Poisson Fossiles. Tome III.
- [70] Ray, C. E., Wetmore, A., Dunkle, D. H., and Drez, P. (1968) Fossil vertebrates from the marine Pleistocene of southeastern Virginia. *Smithsonian Miscellaneous Collections*, **153**, 1–25.
- [71] Quillévéré, F., Koskeridou, E., Cornée, J.-J., Moissette, P., Girone, A., and Agiadi, K. (2018) Pleistocene marine fish invasions and paleoenvironmental reconstructions in the eastern Mediterranean. Quaternary Science Reviews, 196, 80–99.
- [72] Schultz, O., Brzobohatý, R., and Kroupa, O. (2010) Fish teeth from the Middle Miocene of Kienberg at Mikulov, Czech Republic, Vienna Basin. Annalen des Naturhistorischen Museums in Wien, Serie A, 112, 489–506.
- [73] Diaz-Franco, S. and Rojas-Consuegra, R. (2009) Dientes fósiles de *Sphyraena* (Perciformes: Sphyraenidae) en el Terciario de Cuba Occidental. *Solenodon*, **8**, 124–129.
- [74] Böhm, J. (1926) Ueher tertiâre Versteinerungen von den hogenfelser Diamantfeldern. Kaiser, E. (ed.), Die Diamantenwüste Sudwestafrikasfrikas. Vol II, pp. 55–87.
- [75] Westgate, J. W. (1984) Lower vertebrates from the late Eocene Crow Creek local fauna, St. Francis County, Arkansas. *Journal of Vertebrate Paleontology*, 4, 536–546.
- [76] Gottfried, M. D., Samonds, K. E., Ostrowski, S. A., Andrianavalona, T. H., and Ramihangihajason, T. N. (2017) New evidence indicates the presence of Barracuda (Sphyraenidae) and supports a tropical marine environment in the Miocene of Madagascar. PLoS ONE, 12, 1–9.
- [77] ICZN (1999) International Code of Zoological Nomenclature. The International Trust for Zoological Nomenclature, 4th edn.

- [78] Harlan, R. (1824) On a new fossil genus, of the order Enalio Sauri (of Conybeare): and on a new species of Ichthyosaurus. *Journal fo the Academy of Natural Sciences of Philadelphia*, **3**, 331–338.
- [79] Casier, E. (1944) Contributions à l'étude des Poissons fossiles de la Belgique. VI. Sur le Sphyraenodus de l'Eocene e sur la présence d'un Sphyraenidé dans le Bruxellien (Lutétien inférieur). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, 20, 11–15.
- [80] Woodward, A. S. (1891) Catalogue of the Fossil Fishes in the British Museum (Natural History).

  Part II. Taylor & Francis.
- [81] STRINGER, GARY L., Department of Ge (2003) Paleoenvironmental Interpretations Based on Vertebrate Fossil Assemblages: An Example of their Utilization in the Gulf Coast. AAPG Bulletin, 85.
- [82] Páramo-Fonseca, M. E. (1997) Bachea huilensis nov. gen., nov. sp., premier Tselfatioidei (Teleostei) de Colombie. Comptes Rendus de l'Academie de Sciences Serie IIa: Sciences de la Terre et des Planetes, 325, 147–150.
- [83] Chapman, F. (1935) Descriptions of fossil fish from New Zealand. Transactions and Proceedings of the Royal Society of New Zealand, 64, 117–121.
- [84] de Sylva, D. P. and Williams, F. (1986) Sphyraenidae. Smith, M. M. and Heemstra, P. C. (eds.), Smiths' Sea Fishes, pp. 721–726, Macmillan South Africa.
- [85] Ray, C. E. and Bohaska, D. J. (2001) Geology and Paleontology of the Lee Creek Mine, North Carolina, III. No. 90, smithsonia edn.
- [86] Patterson, C. (1993) An overview of the early fossil record of Acanthomorphs. *Bulletin of Marine Science*, **52**, 29–59.
- [87] Bardack, D. and Sprinkle, G. (1969) Morphology and relationships of saurocephalid fishes. *Fieldiana Geology*, **16**, 297–340.
- [88] Monsch, K. A. (2005) Revision of the scombroid fishes from the Cenozoic of England. Transactions of the Royal Society of Edinburgh: Earth Sciences, 95, 445–489.
- [89] Agassiz, L. (1833) Recherches sur les Poissons Fossiles. Tome IV.
- [90] Moreno, F., et al. (2015) Revised stratigraphy of Neogene strata in the Cocinetas Basin, La Guajira, Colombia. Swiss Journal of Palaeontology, 134, 5–43.
- [91] Agassiz, L. (1833), Recherches sur les Poissons Fossiles. Tome I.
- [92] Van der Laan, R. (2018) Family-group names of fossil fishes. European Journal of Taxonomy, pp. 1–167.
- [93] Meek, S. E. and Newland, R. G. (1884) A review of the American species of the genus *Sphyraena*. Proceedings of the Academy of Natural Sciences of Philadelphia, **36**, 67–75.
- [94] Stringer, G. L., Breard, S. Q., and Kontrovitz, M. (2001) Biostratigraphy and Paleoecology of Diagnostic Invertebrates and Vertebrates from the Type Locality of the Oligocene Rosefiled Marl Beds, Louisiana. *Gulf Coast Association of Geological Societies Transactions*, LI, 321–328.
- [95] Cope, E. D. (1869) Synopsis of the Batrachia and Reptilia of North America. Part I. Transactions of the American Philosophical Society, 14, 1–252.
- [96] Switchenska, A. A. (1968) A new genus from the family Sphyraeniadae from the middle Miocene of Transcaucasia. Ocherki po Filogenii i Sistematike Iskopaemykh Ryb I Beschelyustnykh, pp. 157–161.
- [97] Agassiz, L. (1843) Recherches sur les Poisson Fossiles. Tome V.

- [98] Viñola-López, L. W., Rojas-Consuegra, R., and Jiménez-Vásquez, O. (2017) Nuevos registros de Sphyraena (Perciformes: Sphyraenidae) para el Neógeno de Cuba y La Española. Novitates Caribaea, 11, 89–94.
- [99] NCBI (2018) Entrez Programming Utilities Help. No. Md, NCBI.
- [100] Taverne, L. and Chanet, B. (2000) Faugichthys loryi n. gen., n. sp. (Teleostei, Ichthyodectiformes) de l'Albien terminal (Crétacé inférieur marin) du vallon de la Fauge (Isère, France) et considérations sur la phylogénie des Ichthyodectidae. Geodiversitas, 22, 23–34.
- [101] Smith, J. L. B. (1956) The fishes of the family Sphyraenidae in the western Indian Ocean. *Ichthyological Bulletin of the Department of Ichthyology of Rhodes University*, **3**, 37–46.
- [102] Távora, V. d. A., dos Santos, A. A. R., and Araújo, R. N. (2010) Localidades fossilíferas da Formação Pirabas (Mioceno Inferior). *Boletim do Museu Paraense Emilio Goeldi Ciencias Naturais*, 5, 207–224.