

Unit 4: Inference for numerical data

1. Inference using the t -distribution

Sta 101 - Fall 2015

Duke University, Department of Statistical Science

1. Housekeeping

2. Main ideas

1. T corrects for uncertainty introduced by plugging in s for σ
2. When comparing means of two groups, details depend on paired or independent
3. All other details of the inferential framework is the same...

- ▶ Exams returned at the end of lab tomorrow
- ▶ MT grades will be posted for everyone on ACES
- ▶ MT course eval – anonymous, appreciate feedback
- ▶ RA 4 when you get back from fall break

1. Housekeeping

2. Main ideas

1. T corrects for uncertainty introduced by plugging in s for σ
2. When comparing means of two groups, details depend on paired or independent
3. All other details of the inferential framework is the same...

1. Housekeeping

2. Main ideas


1. T corrects for uncertainty introduced by plugging in s for σ
2. When comparing means of two groups, details depend on paired or independent
3. All other details of the inferential framework is the same...

2. T corrects for uncertainty introduced by plugging in s for σ

- ▶ CLT says $\bar{x} \sim N\left(\text{mean} = \mu, SE = \frac{\sigma}{\sqrt{n}}\right)$, but, in practice, we use s instead of σ .
 - Plugging in an estimate introduces additional uncertainty.
 - We make up for this by using a more “conservative” distribution than the normal distribution.

2. T corrects for uncertainty introduced by plugging in s for σ

- ▶ CLT says $\bar{x} \sim N\left(\text{mean} = \mu, SE = \frac{\sigma}{\sqrt{n}}\right)$, but, in practice, we use s instead of σ .
 - Plugging in an estimate introduces additional uncertainty.
 - We make up for this by using a more “conservative” distribution than the normal distribution.
- ▶ t -distribution also has a bell shape, but its tails are *thicker* than the normal model's
 - Observations are more likely to fall beyond two SDs from the mean than under the normal distribution.
 - Extra thick tails help mitigate the effect of a less reliable estimate for the standard error of the sampling distribution.




- ▶ Always centered at zero, like the standard normal (z) distribution

- ▶ Always centered at zero, like the standard normal (z) distribution
- ▶ Has a single parameter, *degrees of freedom* (df), that is tied to sample size.
 - one sample: $df = n - 1$
 - two (independent) samples: $df = \min(n_1 - 1, n_2 - 1)$


- ▶ Always centered at zero, like the standard normal (z) distribution
- ▶ Has a single parameter, *degrees of freedom* (df), that is tied to sample size.
 - one sample: $df = n - 1$
 - two (independent) samples: $df = \min(n_1 - 1, n_2 - 1)$

What happens to shape of the t -distribution as df increases?



- ▶ Always centered at zero, like the standard normal (z) distribution
- ▶ Has a single parameter, *degrees of freedom* (df), that is tied to sample size.
 - one sample: $df = n - 1$
 - two (independent) samples: $df = \min(n_1 - 1, n_2 - 1)$


What happens to shape of the t -distribution as df increases?



*Approaches
normal.*

- ▶ Always centered at zero, like the standard normal (z) distribution
- ▶ Has a single parameter, *degrees of freedom* (df), that is tied to sample size.
 - one sample: $df = n - 1$
 - two (independent) samples: $df = \min(n_1 - 1, n_2 - 1)$

What happens to shape of the t -distribution as df increases?



Approaches normal.

Why?

1. Housekeeping

2. Main ideas

1. T corrects for uncertainty introduced by plugging in s for σ
2. When comparing means of two groups, details depend on paired or independent
3. All other details of the inferential framework is the same...

Trace metals in drinking water affect the flavor and an unusually high concentration can pose a health hazard. Ten pairs of data were taken measuring zinc concentration in bottom water and surface water at 10 randomly sampled locations.

Location	bottom	surface
1	0.43	0.415
2	0.266	0.238
3	0.567	0.39
4	0.531	0.41
5	0.707	0.605
6	0.716	0.609
7	0.651	0.632
8	0.589	0.523
9	0.469	0.411
10	0.723	0.612


Trace metals in drinking water affect the flavor and an unusually high concentration can pose a health hazard. Ten pairs of data were taken measuring zinc concentration in bottom water and surface water at 10 randomly sampled locations.

Location	bottom	surface
1	0.43	0.415
2	0.266	0.238
3	0.567	0.39
4	0.531	0.41
5	0.707	0.605
6	0.716	0.609
7	0.651	0.632
8	0.589	0.523
9	0.469	0.411
10	0.723	0.612

Water samples collected at the same location, on the surface and in the bottom, cannot be assumed to be independent of each other, hence we need to use a *paired* analysis.

Example 2: Gender gap in salaries

Since 2005, the American Community Survey¹ polls ~3.5 million households yearly. The following summarizes distribution of salaries of males and females from a random sample of individuals who responded to the 2012 ACS:



¹Aside: Surge of media attention in spring 2012 when the House of Representatives voted to eliminate the survey. Daniel Webster, Republican congressman from Florida: “in the end this is not a scientific survey. It’s a random survey.”

How are the two examples different from each other? How are they similar to each other?

Suppose we want to compare the average zinc concentration levels in the bottom and surface:

Suppose we want to compare the average zinc concentration levels in the bottom and surface:

- ▶ Two sets of observations with a special correspondence (not independent): *paired*


Suppose we want to compare the average zinc concentration levels in the bottom and surface:

- ▶ Two sets of observations with a special correspondence (not independent): *paired*
- ▶ Synthesize down to differences in outcomes of each pair of observations, subtract using a consistent order

Suppose we want to compare the average zinc concentration levels in the bottom and surface:

- ▶ Two sets of observations with a special correspondence (not independent): *paired*
- ▶ Synthesize down to differences in outcomes of each pair of observations, subtract using a consistent order

Location	bottom	surface	difference
1	0.43	0.415	0.015
2	0.266	0.238	0.028
3	0.567	0.39	0.177
4	0.531	0.41	0.121
5	0.707	0.605	0.102
6	0.716	0.609	0.107
7	0.651	0.632	0.019
8	0.589	0.523	0.066
9	0.469	0.411	0.058
10	0.723	0.612	0.111



For comparing average zinc concentration levels in the bottom and surface when the data are paired:

For comparing average zinc concentration levels in the bottom and surface when the data are paired:

- ▶ *Parameter of interest:* Average difference between the bottom and surface zinc measurements of *all* drinking water.

$$\mu_{diff}$$

For comparing average zinc concentration levels in the bottom and surface when the data are paired:

- ▶ *Parameter of interest:* Average difference between the bottom and surface zinc measurements of *all* drinking water.

$$\mu_{diff}$$

- ▶ *Point estimate:* Average difference between the bottom and surface zinc measurements of drinking water from the *sampled* locations.

$$\bar{x}_{diff}$$

For comparing average salaries in two independent groups

For comparing average salaries in two independent groups

- *Parameter of interest:* Average difference between the average salaries of *all* males and females in the US.

$$\mu_m - \mu_f$$

For comparing average salaries in two independent groups

- ▶ *Parameter of interest:* Average difference between the average salaries of *all* males and females in the US.

$$\mu_m - \mu_f$$

- ▶ *Point estimate:* Average difference between the average salaries of *sampled* males and females in the US.

$$\bar{x}_m - \bar{x}_f$$

- ▶ Dependent (paired) groups (e.g. pre/post weights of subjects in a weight loss study, twin studies, etc.)

$$SE_{\bar{x}_{diff}} = \frac{s_{diff}}{\sqrt{n_{diff}}}$$

- ▶ Independent groups (e.g. grades of students across two sections)

$$SE_{\bar{x}_1 - \bar{x}_2} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

- ▶ For the same data, $SE_{paired} < SE_{independent}$, so be careful about calling data paired

1. Housekeeping

2. Main ideas

1. T corrects for uncertainty introduced by plugging in s for σ
2. When comparing means of two groups, details depend on paired or independent
3. All other details of the inferential framework is the same...

3. All other details of the inferential framework is the same...

$$HT : test\ statistic = \frac{point\ estimate - null}{SE}$$

3. All other details of the inferential framework is the same...

$$HT : \text{test statistic} = \frac{\text{point estimate} - \text{null}}{SE}$$

$$CI : \text{point estimate} \pm \text{critical value} \times SE$$

3. All other details of the inferential framework is the same...

$$HT : \text{test statistic} = \frac{\text{point estimate} - \text{null}}{SE}$$

$$CI : \text{point estimate} \pm \text{critical value} \times SE$$

One mean:

$$df = n - 1$$

HT:

$$H_0 : \mu = \mu_0$$

$$T_{df} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

CI:

$$\bar{x} \pm t_{df}^* \frac{s}{\sqrt{n}}$$

3. All other details of the inferential framework is the same...

$$HT : \text{test statistic} = \frac{\text{point estimate} - \text{null}}{SE}$$

$$CI : \text{point estimate} \pm \text{critical value} \times SE$$

One mean:

$$df = n - 1$$

HT:

$$H_0 : \mu = \mu_0$$

$$T_{df} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

CI:

$$\bar{x} \pm t_{df}^* \frac{s}{\sqrt{n}}$$

Paired means:

$$df = n_{diff} - 1$$

HT:

$$H_0 : \mu_{diff} = 0$$

$$T_{df} = \frac{\bar{x}_{diff} - 0}{\frac{s_{diff}}{\sqrt{n_{diff}}}}$$

CI:

$$\bar{x}_{diff} \pm t_{df}^* \frac{s_{diff}}{\sqrt{n_{diff}}}$$

3. All other details of the inferential framework is the same...

$$HT: \text{test statistic} = \frac{\text{point estimate} - \text{null}}{SE}$$

$$CI: \text{point estimate} \pm \text{critical value} \times SE$$

One mean:

$$df = n - 1$$

HT:

$$H_0 : \mu = \mu_0$$

$$T_{df} = \frac{\bar{x} - \mu}{\frac{s}{\sqrt{n}}}$$

CI:

$$\bar{x} \pm t_{df}^* \frac{s}{\sqrt{n}}$$

Paired means:

$$df = n_{diff} - 1$$

HT:

$$H_0 : \mu_{diff} = 0$$

$$T_{df} = \frac{\bar{x}_{diff} - 0}{\frac{s_{diff}}{\sqrt{n_{diff}}}}$$

CI:

$$\bar{x}_{diff} \pm t_{df}^* \frac{s_{diff}}{\sqrt{n_{diff}}}$$

Independent means:

$$df = \min(n_1 - 1, n_2 - 1)$$

HT:

$$H_0 : \mu_1 - \mu_2 = 0$$

$$T_{df} = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

CI:

$$\bar{x}_1 - \bar{x}_2 \pm t_{df}^* \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$