

Unit 2: Probability and distributions

1. Probability and conditional probability

Sta 101 - Fall 2015

Duke University, Department of Statistical Science

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

4. Summary

- ▶ Piazza – make sure you're enrolled for our class: Sta 101.002
- ▶ Team names – if we're missing yours you must email David asap (after class), or you won't be eligible for today's RA scores

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing

2. Application of the addition rule depends on disjointness of events

3. Bayes' theorem works for all types of events

4. Summary

- ▶ 15 minutes individual – turn your clicker over when you're done
- ▶ 10 minutes team – put your team name on the front of the scratch off sheet + Lab Time + put **only** the names of the members who are present today on the back

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

4. Summary

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing

2. Application of the addition rule depends on disjointness of events

3. Bayes' theorem works for all types of events

4. Summary

1. Disjoint and independent do not mean the same thing

- ▶ *Disjoint (mutually exclusive) events* cannot happen at the same time
 - A voter cannot register as a Democrat and a Republican at the same time
 - But s/he might be a Republican and a Moderate at the same time – *non-disjoint events*
 - For disjoint A and B: $P(A \text{ and } B) = 0$

1. Disjoint and independent do not mean the same thing

- ▶ *Disjoint (mutually exclusive) events* cannot happen at the same time
 - A voter cannot register as a Democrat and a Republican at the same time
 - But s/he might be a Republican and a Moderate at the same time – *non-disjoint events*
 - For disjoint A and B: $P(A \text{ and } B) = 0$
- ▶ If A and B are *independent events*, having information on A does not tell us anything about B (and vice versa)
 - If A and B are independent:
 - $P(A | B) = P(A)$
 - $P(A \text{ and } B) = P(A) \times P(B)$

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing

2. Application of the addition rule depends on disjointness of events

3. Bayes' theorem works for all types of events

4. Summary

2. Application of the addition rule depends on disjointness of events


- ▶ *General addition rule:* $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶ $A \text{ or } B = \text{either } A \text{ or } B \text{ or both}$

2. Application of the addition rule depends on disjointness of events

- ▶ *General addition rule:* $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶ $A \text{ or } B = \text{either } A \text{ or } B \text{ or both}$

disjoint events:

$$\begin{aligned} &P(A \text{ or } B) \\ &= P(A) + P(B) - P(A \text{ and } B) \\ &= 0.4 + 0.3 - 0 = 0.7 \end{aligned}$$




2. Application of the addition rule depends on disjointness of events

- ▶ *General addition rule:* $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
- ▶ $A \text{ or } B = \text{either } A \text{ or } B \text{ or both}$


disjoint events:

$$\begin{aligned}P(A \text{ or } B) \\&= P(A) + P(B) - P(A \text{ and } B) \\&= 0.4 + 0.3 - 0 = 0.7\end{aligned}$$



non-disjoint events:

$$\begin{aligned}P(A \text{ or } B) \\&= P(A) + P(B) - P(A \text{ and } B) \\&= 0.4 + 0.3 - 0.02 = 0.68\end{aligned}$$



1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing

2. Application of the addition rule depends on disjointness of events

3. Bayes' theorem works for all types of events

4. Summary

3. Bayes' theorem works for all types of events

► *Bayes' theorem:* $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A \mid B) \times P(B)$

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A \mid B) \times P(B)$

disjoint events:

- ▶ We know $P(A \mid B) = 0$,
since if B happened A
could not have happened

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A \mid B) \times P(B)$

disjoint events:

- ▶ We know $P(A \mid B) = 0$,
since if B happened A
could not have happened
- ▶ $P(A \text{ and } B)$
 $= P(A \mid B) \times P(B)$

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A \mid B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A \mid B) \times P(B)$

disjoint events:

- ▶ We know $P(A \mid B) = 0$,
since if B happened A
could not have happened
- ▶ $P(A \text{ and } B)$
 $= P(A \mid B) \times P(B)$
 $= 0 \times P(B) = 0$

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A | B) \times P(B)$

disjoint events:

- ▶ We know $P(A | B) = 0$,
since if B happened A
could not have happened
- ▶ $P(A \text{ and } B)$
 $= P(A | B) \times P(B)$
 $= 0 \times P(B) = 0$

independent events:

- ▶ We know $P(A | B) = P(A)$,
since knowing B doesn't
tell us anything about A

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A | B) \times P(B)$

disjoint events:

- ▶ We know $P(A | B) = 0$,
since if B happened A
could not have happened
- ▶ $P(A \text{ and } B)$
 $= P(A | B) \times P(B)$
 $= 0 \times P(B) = 0$

independent events:

- ▶ We know $P(A | B) = P(A)$,
since knowing B doesn't
tell us anything about A
- ▶ $P(A \text{ and } B)$
 $= P(A | B) \times P(B)$

3. Bayes' theorem works for all types of events

- ▶ *Bayes' theorem:* $P(A | B) = \frac{P(A \text{ and } B)}{P(B)}$
- ▶ ... can be rewritten as: $P(A \text{ and } B) = P(A | B) \times P(B)$

disjoint events:

- ▶ We know $P(A | B) = 0$,
since if B happened A
could not have happened
- ▶ $P(A \text{ and } B)$
 $= P(A | B) \times P(B)$
 $= 0 \times P(B) = 0$

independent events:

- ▶ We know $P(A | B) = P(A)$,
since knowing B doesn't
tell us anything about A
- ▶ $P(A \text{ and } B)$
 $= P(A | B) \times P(B)$
 $= P(A) \times P(B)$

Application exercise: 2.1 Probability and conditional probability

See the course website for instructions.

1. Housekeeping

2. Readiness assessment

3. Main ideas

1. Disjoint and independent do not mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events

4. Summary

1. Disjoint and independent do not mean the same thing
2. Application of the addition rule depends on disjointness of events
3. Bayes' theorem works for all types of events