





1. Linear, Gaussian
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
1
0
-1
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


2. Curved exponential
n = 100
n = 250
n = 500
n = 1000
1.0
0.5
0.0
$f_0(x)$
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth



3. Cubic, flat at the center
n = 100
n = 250
n = 500
n = 1000
1.0
0.5
0.0
-0.5
$f_0(x)$
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


4. Heteroscedastic, beta design
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
3
2
1
0
-1
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


5. Student t with 3 degrees of freedom





6. Skewed Pareto errors





7. Constant function





8. Plateau followed by a rise
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
1.5
1.0
0.5
0.0
-0.5
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


9. Unequal slopes at the center
n = 100
n = 250
n = 500
n = 1000
1.5
1.0
0.5
0.0
-0.5
$f_0(x)$
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


10. Increasing function with a jump
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
2
1
0
-1
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


11. Sparse design near the center
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
1.5
1.0
0.5
0.0
-0.5
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth


12. Student t with 1.5 degrees of freedom
n = 100
n = 250
n = 500
n = 1000
$f_0(x)$
2
1
0
-1
-2
0.25
0.50
0.75
Covariate
cosi-m+
HulC (Bonferroni)
Isotonic fit
Truth