

Modern Quantum Chemistry, Szabo & Ostlund

HW

王石嵘

September 7, 2019

Contents

2	Many-electron Wave Functions and Operators	2
2.1	The Electronic Problem	2
2.1.1	Atomic Units	2
2.1.2	The B-O Approximation	2
2.1.3	The Antisymmetry or Pauli Exclusion Principle	2
2.2	Orbitals, Slater Determinants, and Basis Functions	2
2.2.1	Spin Orbitals and Spatial Orbitals	2
Ex 2.1		2
2.2.2	Hartree Products	2
Ex 2.2		2
2.2.3	Slater Determinants	2
Ex 2.3		2
Ex 2.4		2
Ex 2.5		3
2.2.4	The Hartree-Fock Approximation	3
2.2.5	The Minimal Basis H ₂ Model	3
Ex 2.6		3
2.2.6	Excited Determinants	3
2.2.7	Form of the Exact Wfn and CI	3
Ex 2.7		3
2.3	Operators and Matrix Elements	3
2.3.1	Minimal Basis H ₂ Matrix Elements	3
Ex 2.8		3
Ex 2.9		4
2.3.2	Notations for 1- and 2-Electron Integrals	4
2.3.3	General Rules for Matrix Elements	4
Ex 2.10		4
Ex 2.11		4
Ex 2.12		4
Ex 2.13		5
Ex 2.14		5
2.3.4	Derivation of the Rules for Matrix Elements	5
Ex 2.15		5
Ex 2.16		5
2.3.5	Transition from Spin Orbitals to Spatial Orbitals	5
Ex 2.17		5
Ex 2.18		6
2.3.6	Coulomb and Exchange Integrals	6
Ex 2.19		6
Ex 2.20		7
Ex 2.21		7
Ex 2.22		7

2 Many-electron Wave Functions and Operators

2.1 The Electronic Problem

2.1.1 Atomic Units

2.1.2 The B-O Approximation

2.1.3 The Antisymmetry or Pauli Exclusion Principle

2.2 Orbitals, Slater Determinants, and Basis Functions

2.2.1 Spin Orbitals and Spatial Orbitals

Ex 2.1 Consider $\langle \chi_k | \chi_m \rangle$. If $k = m$,

$$\langle \chi_{2i-1} | \chi_{2i-1} \rangle = \langle \psi_i^\alpha | \psi_i^\alpha \rangle \langle \alpha | \alpha \rangle = 1 \quad (2.2.1)$$

$$\langle \chi_{2i} | \chi_{2i} \rangle = \langle \psi_i^\beta | \psi_i^\beta \rangle \langle \alpha | \alpha \rangle = 1 \quad (2.2.2)$$

thus

$$\langle \chi_k | \chi_k \rangle = 1 \quad (2.2.3)$$

If $k \neq m$, three cases may occur as below

$$\langle \chi_{2i-1} | \chi_{2j-1} \rangle = \langle \psi_i^\alpha | \psi_j^\alpha \rangle \langle \alpha | \alpha \rangle = 0 \cdot 1 = 0 \quad (i \neq j) \quad (2.2.4)$$

$$\langle \chi_{2i-1} | \chi_{2j} \rangle = \langle \psi_i^\alpha | \psi_j^\beta \rangle \langle \alpha | \beta \rangle = S_{ij} \cdot 0 = 0 \quad (2.2.5)$$

$$\langle \chi_{2i} | \chi_{2j} \rangle = \langle \psi_i^\beta | \psi_j^\beta \rangle \langle \beta | \beta \rangle = 0 \cdot 1 = 0 \quad (i \neq j) \quad (2.2.6)$$

thus

$$\langle \chi_k | \chi_m \rangle = 0 \quad (k \neq m) \quad (2.2.7)$$

Overall,

$$\langle \chi_k | \chi_m \rangle = \delta_{km} \quad (2.2.8)$$

2.2.2 Hartree Products

Ex 2.2

$$\begin{aligned} \mathcal{H}\Psi^{HP} &= \sum_{i=1}^N h(i) \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) \cdots \chi_k(\mathbf{x}_N) \\ &= \varepsilon_i \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) \cdots \chi_k(\mathbf{x}_N) + \chi_i(\mathbf{x}_1) [\varepsilon_j \chi_j(\mathbf{x}_2)] \cdots \chi_k(\mathbf{x}_N) + \cdots + \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) \cdots [\varepsilon_k \chi_k(\mathbf{x}_N)] \\ &= (\varepsilon_i + \varepsilon_j + \cdots + \varepsilon_k) \Psi^{HP} \end{aligned} \quad (2.2.9)$$

2.2.3 Slater Determinants

Ex 2.3

$$\begin{aligned} \langle \Psi | \Psi \rangle &= \frac{1}{2} (\langle \chi_i | \chi_i \rangle \langle \chi_j | \chi_j \rangle - \langle \chi_i | \chi_j \rangle \langle \chi_j | \chi_i \rangle - \langle \chi_j | \chi_i \rangle \langle \chi_i | \chi_j \rangle + \langle \chi_j | \chi_j \rangle \langle \chi_i | \chi_i \rangle) \\ &= \frac{1}{2} (1 + 0 + 0 + 1) = 1 \end{aligned} \quad (2.2.10)$$

Ex 2.4 According to Ex. 2.2, we know that $\chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2)$ are an eigenfunction of \mathcal{H} and has the eigenvalue $\varepsilon_i \varepsilon_j$. Similarly, we have the same conclusion for $\chi_i(\mathbf{x}_2) \chi_j(\mathbf{x}_1)$.

For the antisymmetrized wave function,

$$\begin{aligned} \langle \Psi | \mathcal{H} | \Psi \rangle &= \frac{1}{2} (\langle \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) | \mathcal{H} | \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) \rangle - \langle \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) | \mathcal{H} | \chi_j(\mathbf{x}_1) \chi_i(\mathbf{x}_2) \rangle \\ &\quad - \langle \chi_j(\mathbf{x}_1) \chi_i(\mathbf{x}_2) | \mathcal{H} | \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) \rangle + \langle \chi_j(\mathbf{x}_1) \chi_i(\mathbf{x}_2) | \mathcal{H} | \chi_j(\mathbf{x}_1) \chi_i(\mathbf{x}_2) \rangle) \\ &= \frac{1}{2} (\varepsilon_i + \varepsilon_j - 0 - 0 + \varepsilon_i + \varepsilon_j) \\ &= \varepsilon_i + \varepsilon_j \end{aligned} \quad (2.2.11)$$

Ex 2.5

$$\begin{aligned}
\langle K | L \rangle &= \frac{1}{2} \langle \chi_i(\mathbf{x}_1) \chi_j(\mathbf{x}_2) - \chi_j(\mathbf{x}_1) \chi_i(\mathbf{x}_2) | \chi_k(\mathbf{x}_1) \chi_l(\mathbf{x}_2) - \chi_l(\mathbf{x}_1) \chi_k(\mathbf{x}_2) \rangle \\
&= \frac{1}{2} (\langle \chi_i | \chi_k \rangle \langle \chi_j | \chi_l \rangle - \langle \chi_i | \chi_l \rangle \langle \chi_j | \chi_k \rangle - \langle \chi_j | \chi_k \rangle \langle \chi_i | \chi_l \rangle + \langle \chi_j | \chi_l \rangle \langle \chi_i | \chi_k \rangle) \\
&= \frac{1}{2} (\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk} - \delta_{jk} \delta_{il} + \delta_{jl} \delta_{ik}) \\
&= \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}
\end{aligned} \tag{2.2.12}$$

2.2.4 The Hartree-Fock Approximation

2.2.5 The Minimal Basis H₂ Model

Ex 2.6

$$\langle \psi_1 | \psi_1 \rangle = \frac{1}{2(1 + S_{12})} (\langle \phi_1 | \phi_1 \rangle + 2 \langle \phi_1 | \phi_2 \rangle + \langle \phi_2 | \phi_2 \rangle) = \frac{2 + 2S_{12}}{2(1 + S_{12})} = 1 \tag{2.2.13}$$

$$\langle \psi_2 | \psi_2 \rangle = \frac{1}{2(1 - S_{12})} (\langle \phi_1 | \phi_1 \rangle - 2 \langle \phi_1 | \phi_2 \rangle + \langle \phi_2 | \phi_2 \rangle) = \frac{2 - 2S_{12}}{2(1 - S_{12})} = 1 \tag{2.2.14}$$

$$\langle \psi_1 | \psi_2 \rangle = \frac{1}{2\sqrt{1 + S_{12}}\sqrt{1 - S_{12}}} (\langle \phi_1 | \phi_1 \rangle - \langle \phi_2 | \phi_2 \rangle) = 0 \tag{2.2.15}$$

2.2.6 Excited Determinants

2.2.7 Form of the Exact Wfn and CI

Ex 2.7 Size of full CI matrix

$$C_{72}^{42} = 164307576757973059488 \approx 1.64 \times 10^{20} \tag{2.2.16}$$

The number of singly excited determinants

$$42 \times 30 = 1260 \tag{2.2.17}$$

The number of doubly excited determinants

$$C_{42}^2 C_{30}^2 = 374535 \tag{2.2.18}$$

2.3 Operators and Matrix Elements

2.3.1 Minimal Basis H₂ Matrix Elements

Ex 2.8

$$\begin{aligned}
\langle \Psi_{12}^{34} | h(1) | \Psi_{12}^{34} \rangle &= \frac{1}{2} \langle \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) | h(1) | \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \rangle \\
&= \frac{1}{2} (\langle \chi_3 | h(1) | \chi_3 \rangle - 0 - 0 + \langle \chi_4 | h(1) | \chi_4 \rangle) \\
&= \frac{1}{2} (\langle \chi_3 | h(1) | \chi_3 \rangle + \langle \chi_4 | h(1) | \chi_4 \rangle)
\end{aligned} \tag{2.3.1}$$

thus

$$\langle \Psi_{12}^{34} | \mathcal{O}_1 | \Psi_{12}^{34} \rangle = \langle 3 | h | 3 \rangle + \langle 4 | h | 4 \rangle \tag{2.3.2}$$

$$\begin{aligned}
\langle \Psi_0 | h(1) | \Psi_{12}^{34} \rangle &= \frac{1}{2} \langle \chi_1(\mathbf{x}_1) \chi_2(\mathbf{x}_2) - \chi_2(\mathbf{x}_2) \chi_1(\mathbf{x}_1) | h(1) | \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \rangle \\
&= \frac{1}{2} (0 - 0 - 0 + 0) \\
&= 0
\end{aligned} \tag{2.3.3}$$

thus

$$\langle \Psi_0 | \mathcal{O}_1 | \Psi_{12}^{34} \rangle = 0 \tag{2.3.4}$$

Similarly, we get

$$\langle \Psi_{12}^{34} | \mathcal{O}_1 | \Psi_0 \rangle = 0 \tag{2.3.5}$$

Ex 2.9 From Eq. (2.92) in textbook, we get

$$\langle \Psi_0 | \mathcal{H} | \Psi_0 \rangle = \langle 1 | h | 1 \rangle + \langle 2 | h | 2 \rangle + \langle 12 | 12 \rangle - \langle 12 | 21 \rangle \quad (2.3.6)$$

From Ex 2.8, we get

$$\langle \Psi_0 | \mathcal{O}_1 | \Psi_{12}^{34} \rangle = \langle \Psi_{12}^{34} | \mathcal{O}_1 | \Psi_0 \rangle = 0 \quad (2.3.7)$$

thus

$$\begin{aligned} \langle \Psi_0 | \mathcal{H} | \Psi_{12}^{34} \rangle &= \langle \Psi_0 | \mathcal{O}_2 | \Psi_{12}^{34} \rangle \\ &= \frac{1}{2} \left\langle \chi_1(\mathbf{x}_1) \chi_2(\mathbf{x}_2) - \chi_1(\mathbf{x}_2) \chi_2(\mathbf{x}_1) \left| \frac{1}{r_{12}} \right| \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \right\rangle \\ &= \langle 12 | 34 \rangle - \langle 12 | 43 \rangle \end{aligned} \quad (2.3.8)$$

$$\begin{aligned} \langle \Psi_{12}^{34} | \mathcal{H} | \Psi_0 \rangle &= \langle \Psi_{12}^{34} | \mathcal{O}_2 | \Psi_0 \rangle \\ &= \frac{1}{2} \left\langle \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \left| \frac{1}{r_{12}} \right| \chi_1(\mathbf{x}_1) \chi_2(\mathbf{x}_2) - \chi_2(\mathbf{x}_2) \chi_1(\mathbf{x}_1) \right\rangle \\ &= \langle 34 | 12 \rangle - \langle 34 | 21 \rangle \end{aligned} \quad (2.3.9)$$

$$\begin{aligned} \langle \Psi_{12}^{34} | \mathcal{H} | \Psi_{12}^{34} \rangle &= \left\langle \Psi_{12}^{34} \left| h(1) + h(2) + \frac{1}{r_{12}} \right| \Psi_{12}^{34} \right\rangle \\ &= 2 \times \frac{1}{2} \langle \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) | h(1) | \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \rangle \\ &\quad + \frac{1}{2} \left\langle \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \left| \frac{1}{r_{12}} \right| \chi_3(\mathbf{x}_1) \chi_4(\mathbf{x}_2) - \chi_3(\mathbf{x}_2) \chi_4(\mathbf{x}_1) \right\rangle \\ &= \langle 3 | h | 3 \rangle + \langle 4 | h | 4 \rangle + \langle 34 | 34 \rangle - \langle 34 | 43 \rangle \end{aligned} \quad (2.3.10)$$

2.3.2 Notations for 1- and 2-Electron Integrals

2.3.3 General Rules for Matrix Elements

Ex 2.10

$$\langle K | \mathcal{H} | K \rangle = \sum_m^N [m|h|m] + \frac{1}{2} \sum_m^N \sum_n^N \langle mn || mn \rangle = \sum_m^N [m|h|m] + \frac{1}{2} \sum_m^N \sum_n^N ([mm|nn] - [mn|nm]) \quad (2.3.11)$$

When $m = n$,

$$[mm|mm] - [mm|mm] = 0 \quad (2.3.12)$$

thus

$$\langle K | \mathcal{H} | K \rangle = \sum_m^N [m|h|m] + \frac{1}{2} \sum_m^N \sum_{n \neq m}^N ([mm|nn] - [mn|nm]) = \sum_m^N [m|h|m] + \sum_m^N \sum_{n > m}^N ([mm|nn] - [mn|nm]) \quad (2.3.13)$$

Ex 2.11

$$\begin{aligned} \langle K | \mathcal{H} | K \rangle &= \langle K | \mathcal{O}_1 + \mathcal{O}_2 | K \rangle = \sum_m^N [m|h|m] + \sum_m^N \sum_{n > m}^N \langle mn || mn \rangle \\ &= \langle 1 | h | 1 \rangle + \langle 2 | h | 2 \rangle + \langle 3 | h | 3 \rangle + \langle 12 || 12 \rangle + \langle 13 || 13 \rangle + \langle 23 || 23 \rangle \end{aligned} \quad (2.3.14)$$

Ex 2.12

$$\begin{aligned} \langle \Psi_0 | \mathcal{H} | \Psi_0 \rangle &= \langle 1 | h | 1 \rangle + \langle 2 | h | 2 \rangle + \langle 12 || 12 \rangle \\ &= \langle 1 | h | 1 \rangle + \langle 2 | h | 2 \rangle + \langle 12 | 12 \rangle - \langle 12 | 21 \rangle \end{aligned} \quad (2.3.15)$$

$$\langle \Psi_0 | \mathcal{H} | \Psi_{12}^{34} \rangle = \langle 12 || 34 \rangle = \langle 12 | 34 \rangle - \langle 12 | 43 \rangle \quad (2.3.16)$$

$$\langle \Psi_{12}^{34} | \mathcal{H} | \Psi_0 \rangle = \langle 34 || 12 \rangle = \langle 34 | 12 \rangle - \langle 34 | 21 \rangle \quad (2.3.17)$$

$$\begin{aligned} \langle \Psi_{12}^{34} | \mathcal{H} | \Psi_{12}^{34} \rangle &= \langle 3 | h | 3 \rangle + \langle 4 | h | 4 \rangle + \langle 34 || 34 \rangle \\ &= \langle 3 | h | 3 \rangle + \langle 4 | h | 4 \rangle + \langle 34 | 34 \rangle - \langle 34 | 43 \rangle \end{aligned} \quad (2.3.18)$$

Which are exactly the same with Ex 2.9.

Ex 2.13 if $a = b, r = s$

$$\langle \Psi_a^r | \mathcal{O} | \Psi_b^s \rangle = \langle \Psi_a^r | \mathcal{O}_1 | \Psi_a^r \rangle = \sum_c^N \langle c | h | c \rangle - \langle a | h | a \rangle + \langle r | h | r \rangle \quad (2.3.19)$$

if $a = b, r \neq s$

$$\langle \Psi_a^r | \mathcal{O} | \Psi_b^s \rangle = \langle \Psi_a^r | \mathcal{O}_1 | \Psi_a^s \rangle = \langle r | h | s \rangle \quad (2.3.20)$$

if $a \neq b, r = s$

$$\langle \Psi_a^r | \mathcal{O} | \Psi_b^s \rangle = \langle \Psi_a^r | \mathcal{O}_1 | \Psi_b^r \rangle = \langle \Psi_a^r | \mathcal{O}_1 | -(\Psi_a^r)_b^a \rangle = -\langle b | h | a \rangle \quad (2.3.21)$$

if $a \neq b, r \neq s$

$$\langle \Psi_a^r | \mathcal{O} | \Psi_b^s \rangle = \langle \Psi_a^r | \mathcal{O}_1 | (\Psi_a^r)_{rb}^{as} \rangle = 0 \quad (2.3.22)$$

Ex 2.14

$${}^N E_0 = \sum_m^N \langle m | h | m \rangle + \sum_m^M \sum_{n>m}^M \langle mn || mn \rangle \quad (2.3.23)$$

$${}^{N-1} E_0 = \sum_{m \neq a}^N \langle m | h | m \rangle + \sum_{m \neq a}^M \sum_{n>m, n \neq a}^M \langle mn || mn \rangle \quad (2.3.24)$$

$${}^N E_0 - {}^{N-1} E_0 = \langle a | h | a \rangle + \sum_{b \neq a}^N \langle ab || ab \rangle \quad (2.3.25)$$

2.3.4 Derivation of the Rules for Matrix Elements

Ex 2.15

$$\begin{aligned} \langle \Psi | \mathcal{H} | \Psi \rangle &= \frac{1}{N!} \left\langle \sum_{n=1}^{N!} (-1)^{p_n} \mathcal{P}_n \{ \chi_i(1) \chi_j(2) \cdots \chi_k(N) \} \left| \sum_{c=1}^N h(c) \right| \sum_{m=1}^{N!} (-1)^{p_m} \mathcal{P}_m \{ \chi_i(1) \chi_j(2) \cdots \chi_k(N) \} \right\rangle \\ &= \frac{1}{N!} \sum_{n=1}^{N!} \sum_{m=1}^{N!} (-1)^{p_n+p_m} \sum_{c=1}^N \langle \mathcal{P}_n \{ \chi_i(1) \chi_j(2) \cdots \chi_k(N) \} | h(c) | \mathcal{P}_m \{ \chi_i(1) \chi_j(2) \cdots \chi_k(N) \} \rangle \end{aligned} \quad (2.3.26)$$

Since the integral inside equals 0 when $\mathcal{P}_n \neq \mathcal{P}_m$,

$$\langle \Psi | \mathcal{H} | \Psi \rangle = \frac{1}{N!} \sum_{n=1}^{N!} (-1)^{p_n+p_n} (\varepsilon_i + \varepsilon_j + \cdots + \varepsilon_k) = \varepsilon_i + \varepsilon_j + \cdots + \varepsilon_k \quad (2.3.27)$$

Ex 2.16 Suppose

$$c = \langle K^{HP} | \mathcal{H} | L \rangle = \left\langle K^{HP} \left| \mathcal{H} \right| \sum_{m=1}^{N!} (-1)^{p_m} \mathcal{P}_m L^{HP} \right\rangle \quad (2.3.28)$$

thus

$$\langle K | \mathcal{H} | L \rangle = \sum_{n=1}^{N!} (-1)^{p_n} \left\langle \mathcal{P}_n K^{HP} \left| \mathcal{H} \right| \sum_{m=1}^{N!} (-1)^{p_m} \mathcal{P}_m L^{HP} \right\rangle \quad (2.3.29)$$

2.3.5 Transition from Spin Orbitals to Spatial Orbitals

Ex 2.17

$$\begin{aligned} |1\rangle &= |\psi_1 \alpha\rangle & |2\rangle &= |\psi_1 \beta\rangle \\ |3\rangle &= |\psi_2 \alpha\rangle & |4\rangle &= |\psi_2 \beta\rangle \end{aligned} \quad (2.3.30)$$

thus

$$\begin{aligned} \mathbf{H} &= \begin{pmatrix} \langle 1 | h | 1 \rangle + \langle 2 | h | 2 \rangle + \langle 12 | 12 \rangle - \langle 12 | 21 \rangle & \langle 12 | 34 \rangle - \langle 12 | 43 \rangle \\ \langle 34 | 12 \rangle - \langle 34 | 21 \rangle & \langle 3 | h | 3 \rangle + \langle 4 | h | 4 \rangle + \langle 34 | 34 \rangle - \langle 34 | 43 \rangle \end{pmatrix} \\ &= \begin{pmatrix} 2(1|h|1) + (11|11) & (12|12) \\ (21|21) & 2(2|h|2) + (22|22) \end{pmatrix} \end{aligned} \quad (2.3.31)$$

Ex 2.18

$$\begin{aligned}
|\langle ab || rs \rangle|^2 &= (\langle ab | rs \rangle - \langle ab | sr \rangle)^* (\langle ab | rs \rangle - \langle ab | sr \rangle) \\
&= \langle rs | ab \rangle \langle ab | rs \rangle - \langle rs | ab \rangle \langle ab | sr \rangle - \langle sr | ab \rangle \langle ab | rs \rangle + \langle sr | ab \rangle \langle ab | sr \rangle \\
&= [ra|sb][ar|bs] - [ra|sb][as|br] - [sa|rb][ar|bs] + [sa|rb][as|br] \\
&= [ar|bs]^2 - 2[ar|bs][as|br] + [as|br]^2
\end{aligned} \tag{2.3.32}$$

Let's calculate $E_0^{(2)}$ term by term.

$$\begin{aligned}
(E_0^{(2)})_1 &= \frac{1}{4} \sum_{abrs} \frac{[ar|bs]^2}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= \frac{1}{4} \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{[ar|bs]^2 + [\bar{a}\bar{r}|bs]^2 + [ar|\bar{b}\bar{s}]^2 + [\bar{a}\bar{r}|\bar{b}\bar{s}]^2}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{[ar|bs]^2}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{\langle ab | rs \rangle \langle rs | ab \rangle}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s}
\end{aligned} \tag{2.3.33}$$

$$\begin{aligned}
(E_0^{(2)})_2 &= \frac{1}{4} \sum_{abrs} \frac{-2[ar|bs][as|br]}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= -\frac{1}{2} \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{[ar|bs][as|br] + [\bar{a}\bar{r}|\bar{b}\bar{s}][\bar{a}\bar{s}|\bar{b}\bar{r}]}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= -\sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{[ar|bs][as|br]}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \\
&= -\sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{\langle ab | rs \rangle \langle rs | ba \rangle}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s}
\end{aligned} \tag{2.3.34}$$

$$\begin{aligned}
(E_0^{(2)})_3 &= \frac{1}{4} \sum_{abrs} \frac{[as|br]^2}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} = \frac{1}{4} \sum_{abrs} \frac{[ar|bs]^2}{\varepsilon_a + \varepsilon_b - \varepsilon_s - \varepsilon_r} \\
&= \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{\langle ab | rs \rangle \langle rs | ab \rangle}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s}
\end{aligned} \tag{2.3.35}$$

thus,

$$E_0^{(2)} = \sum_{a,b}^{N/2} \sum_{r,s=N/2+1}^K \frac{\langle ab | rs \rangle (2 \langle rs | ab \rangle - \langle rs | ba \rangle)}{\varepsilon_a + \varepsilon_b - \varepsilon_r - \varepsilon_s} \tag{2.3.36}$$

2.3.6 Coulomb and Exchange Integrals

Ex 2.19

$$J_{ii} = (ii|ii) = K_{ii} \tag{2.3.37}$$

$$J_{ij}^* = \langle ij | ij \rangle^* = \langle ij | ij \rangle = J_{ij} \tag{2.3.38}$$

$$K_{ij}^* = \langle ij | ji \rangle^* = \langle ji | ij \rangle = \langle ij | ji \rangle = K_{ij} \tag{2.3.39}$$

$$J_{ij} = (ii|jj) = (jj|ii) = J_{ji} \tag{2.3.40}$$

$$K_{ij} = (ij|ji) = (ji|ij) = K_{ji} \tag{2.3.41}$$

Ex 2.20 For real spatial orbitals

$$K_{ij} = (ij|ji) = (ij|ij) = (ji|ji) \quad (2.3.42)$$

$$K_{ij} = \langle ij | ji \rangle = \langle ii | jj \rangle = \langle jj | ii \rangle \quad (2.3.43)$$

Ex 2.21

$$\mathbf{H} = \begin{pmatrix} 2(1|h|1) + (11|11) & (12|12) \\ (21|21) & 2(2|h|2) + (22|22) \end{pmatrix} = \begin{pmatrix} 2h_{11} + J_{11} & K_{12} \\ K_{12} & 2h_{22} + J_{22} \end{pmatrix} \quad (2.3.44)$$

Ex 2.22

$$E_{\uparrow\downarrow}^{HP} = \left\langle \Psi_{\uparrow\downarrow}^{HP} \left| h(1) + h(2) + \frac{1}{r_{12}} \right| \Psi_{\uparrow\downarrow}^{HP} \right\rangle = (1|h|1) + (2|h|2) + (11|22) = h_{11} + h_{22} + J_{12} \quad (2.3.45)$$

$$E_{\downarrow\downarrow}^{HP} = \left\langle \Psi_{\downarrow\downarrow}^{HP} \left| h(1) + h(2) + \frac{1}{r_{12}} \right| \Psi_{\downarrow\downarrow}^{HP} \right\rangle = (1|h|1) + (2|h|2) + (11|22) = h_{11} + h_{22} + J_{12} \quad (2.3.46)$$

2.3.7 Pseudo-Classical Interpretation of Determinantal Energies