

Modern Quantum Chemistry, Szabo & Ostlund

HW

王石嵘

February 5, 2020

Contents

3	The Hartree-Fock Approximation	3
3.1	The HF Equations	3
3.1.1	The Coulomb and Exchange Operators	3
3.1.2	The Fock Operator	3
	Ex 3.1	3
3.2	Derivation of the HF Equations	3
3.2.1	Functional Variation	3
3.2.2	Minimization of the Energy of a Single Determinant	3
	Ex 3.2	3
	Ex 3.3	3
3.2.3	The Canonical HF Equations	4
3.3	Interpretation of Solutions to the HF Equations	4
3.3.1	Orbital Energies and Koopmans' Theorem	4
	Ex 3.4	4
	Ex 3.5	4
	Ex 3.6	5
3.3.2	Brillouin's Theorem	5
3.3.3	The HF Hamiltonian	5
	Ex 3.7	5
	Ex 3.8	5
3.4	Restricted Closed-shell HF: The Roothaan Equations	5
3.4.1	Closed-shell HF: Restricted Spin Orbitals	5
	Ex 3.9	5
3.4.2	Introduction of a Basis: The Roothaan Equations	6
	Ex 3.10	6
3.4.3	The Charge Density	6
	Ex 3.11	6
	Ex 3.12	6
	Ex 3.13	7
3.4.4	Expression for the Fock Matrix	7
	Ex 3.14	7
3.4.5	Orthogonalization of the Basis	8
	Ex 3.15	8
	Ex 3.16	8
3.4.6	The SCF Procedure	9
3.4.7	Expectation Values and Population Analysis	9
	Ex 3.17	9
	Ex 3.18	9
3.5	Model Calculations on H_2 and HeH^+	10
3.5.1	The 1s Minimal STO-3G Basis Set	10
	Ex 3.19	10
	Ex 3.20	10
3.5.2	STO-3G H_2	11

	Ex 3.21	11
	Ex 3.22	11
	Ex 3.23	11
	Ex 3.24	11
	Ex 3.25	12
	Ex 3.26	12
	Ex 3.27	12
3.5.3	An SCF Calculation on STO-3G HeH^+	12
	Ex 3.28	12
	Ex 3.29	12
3.6	Polyatomic Basis Sets	12
3.6.1	Contracted Gaussian Functions	12
3.6.2	Minimal Basis Sets: STO-3G	12
3.6.3	Double Zeta Basis Sets: 4-31G	12
	Ex 3.30	12
3.6.4	Polarized Basis Sets: 6-31G* and 6-31G**	13
	Ex 3.31	13
3.7	Some Illustrative Closed-shell Calculations	13
3.7.1	Total Energies	13
	Ex 3.32	13
3.7.2	Ionization Potentials	13
3.7.3	Equilibrium Geometries	13
3.7.4	Population Analysis and Dipole Moments	13
3.8	Unrestricted Open-shell HF: The Pople-Nesbet Equations	13
3.8.1	Open-shell HF: Unrestricted Spin Orbitals	13
	Ex 3.33	13
	Ex 3.34	14
	Ex 3.35	14
3.8.2	Introduction of a Basis: The Pople-Nesbet Equations	14
3.8.3	Unrestricted Density Matrices	14
	Ex 3.36	14
	Ex 3.37	15
	Ex 3.38	15
	Ex 3.39	15
3.8.4	Expression for the Fock Matrices	16
3.8.5	Solution of the Unrestricted SCF Equations	16
	Ex 3.40	16

3 The Hartree-Fock Approximation

3.1 The HF Equations

3.1.1 The Coulomb and Exchange Operators

3.1.2 The Fock Operator

Ex 3.1

$$\begin{aligned}
 \langle \chi_i | \hat{f} | \chi_j \rangle &= \left\langle \chi_i(1) \left| h(1) + \sum_b [\mathcal{J}_b(1) - \mathcal{K}_b(1)] \right| \chi_j(1) \right\rangle \\
 &= [i|h|j] + \sum_{b \neq j} \left[\left\langle \chi_i(1) \chi_b(2) \left| \frac{1}{r_{12}} \right| \chi_b(2) \chi_j(1) \right\rangle - \left\langle \chi_i(1) \chi_b(2) \left| \frac{1}{r_{12}} \right| \chi_b(1) \chi_j(2) \right\rangle \right] \\
 &= [i|h|j] + \sum_{b \neq j} ([ij|bb] - [ib|bj])
 \end{aligned} \tag{3.1.1}$$

Since

$$[ij|jj] - [ij|jj] = 0 \tag{3.1.2}$$

we have

$$\begin{aligned}
 \langle \chi_i | \hat{f} | \chi_j \rangle &= \langle i | h | j \rangle + \sum_b (\langle ib | jb \rangle - \langle ib | bj \rangle) \\
 &= \langle i | h | j \rangle + \sum_b \langle ib || jb \rangle
 \end{aligned} \tag{3.1.3}$$

3.2 Derivation of the HF Equations

3.2.1 Functional Variation

3.2.2 Minimization of the Energy of a Single Determinant

Ex 3.2 Take the complex conjugate of

$$\mathcal{L}[\{\chi_\alpha\}] = E_0[\{\chi_\alpha\}] - \sum_a^N \sum_b^N \varepsilon_{ba}([a|b] - \delta_{ab}) \tag{3.2.1}$$

we have

$$\mathcal{L}[\{\chi_\alpha\}]^* = E_0[\{\chi_\alpha\}]^* - \sum_a^N \sum_b^N \varepsilon_{ba}^*([a|b]^* - \delta_{ab}^*) \tag{3.2.2}$$

i.e.

$$\mathcal{L}[\{\chi_\alpha\}] = E_0[\{\chi_\alpha\}] - \sum_a^N \sum_b^N \varepsilon_{ba}^*([b|a] - \delta_{ab}) \tag{3.2.3}$$

thus

$$\sum_a^N \sum_b^N \varepsilon_{ba}([a|b] - \delta_{ab}) = \sum_a^N \sum_b^N \varepsilon_{ba}^*([b|a] - \delta_{ab}) = \sum_b^N \sum_a^N \varepsilon_{ab}^*([a|b] - \delta_{ba}) \tag{3.2.4}$$

\therefore

$$\varepsilon_{ba} = \varepsilon_{ab}^* \tag{3.2.5}$$

Ex 3.3 \therefore

$$[\delta \chi_a | h | \chi_a] = [\chi_a | h | \delta \chi_a]^* \tag{3.2.6}$$

$$[\chi_a \delta \chi_a | \chi_b \chi_b] = [\delta \chi_a \chi_a | \chi_b \chi_b]^* \tag{3.2.7}$$

$$[\chi_a \chi_a | \chi_b \delta \chi_b] = [\chi_a \chi_a | \delta \chi_b \chi_b]^* \tag{3.2.8}$$

$$[\chi_a \chi_b | \chi_b \delta \chi_a] = [\chi_b \delta \chi_a | \chi_a \chi_b] = [\delta \chi_a \chi_b | \chi_b \chi_a]^* \tag{3.2.9}$$

$$[\chi_a \chi_b | \delta \chi_b \chi_a] = [\delta \chi_b \chi_a | \chi_a \chi_b] = [\chi_a \delta \chi_b | \chi_b \chi_a]^* \tag{3.2.10}$$

∴

$$\begin{aligned}\delta E_0 &= \sum_a^N [\delta \chi_a | h | \chi_a] + \frac{1}{2} \sum_a^N \sum_b^N ([\delta \chi_a \chi_a | \chi_b \chi_b] + [\chi_a \chi_a | \delta \chi_b \chi_b]) \\ &\quad - \frac{1}{2} \sum_a^N \sum_b^N ([\delta \chi_a \chi_b | \chi_b \chi_a] + [\chi_a \chi_b | \delta \chi_b \chi_a]) + \text{complex conjugates}\end{aligned}\quad (3.2.11)$$

while

$$\sum_a^N \sum_b^N [\chi_a \chi_a | \delta \chi_b \chi_b] = \sum_b^N \sum_a^N [\chi_b \chi_b | \delta \chi_a \chi_a] = \sum_a^N \sum_b^N [\delta \chi_a \chi_a | \chi_b \chi_b] \quad (3.2.12)$$

$$\sum_a^N \sum_b^N [\chi_a \chi_b | \delta \chi_b \chi_a] = \sum_b^N \sum_a^N [\chi_b \chi_a | \delta \chi_a \chi_b] = \sum_a^N \sum_b^N [\delta \chi_a \chi_b | \chi_b \chi_a] \quad (3.2.13)$$

thus

$$\delta E_0 = \sum_a^N [\delta \chi_a | h | \chi_a] + \sum_a^N \sum_b^N ([\delta \chi_a \chi_a | \chi_b \chi_b] - [\delta \chi_a \chi_b | \chi_b \chi_a]) + \text{complex conjugates} \quad (3.2.14)$$

3.2.3 The Canonical HF Equations

3.3 Interpretation of Solutions to the HF Equations

3.3.1 Orbital Energies and Koopmans' Theorem

Ex 3.4

$$f_{ij} = \langle \chi_i | f | \chi_j \rangle = \langle i | h | j \rangle + \sum_b \langle ib || jb \rangle \quad (3.3.1)$$

$$\begin{aligned}f_{ji}^* &= \langle \chi_j | f | \chi_i \rangle^* = \langle j | h | i \rangle^* + \sum_b \langle jb || ib \rangle^* \\ &= \langle i | h | j \rangle + \sum_b \langle ib || jb \rangle \\ &= f_{ij}\end{aligned}\quad (3.3.2)$$

thus the Fock operator is Hermitian.

Ex 3.5

$$\text{IP} = {}^{N-2} E - E_0$$

$$\begin{aligned}&= \sum_{a \neq c, d} \langle a | h | a \rangle + \frac{1}{2} \sum_{a \neq c, d} \sum_{b \neq c, d} \langle ab || ab \rangle - \left[\sum_a \langle a | h | a \rangle + \frac{1}{2} \sum_a \sum_b \langle ab || ab \rangle \right] \\ &= -\langle c | h | c \rangle - \langle d | h | d \rangle - \frac{1}{2} \sum_{a \neq c, d} \langle ac || ac \rangle - \frac{1}{2} \sum_{a \neq c, d} \langle ad || ad \rangle - \frac{1}{2} \sum_{b \neq c, d} \langle cb || cb \rangle - \frac{1}{2} \sum_{b \neq c, d} \langle db || db \rangle - \langle cd || cd \rangle \\ &= -\langle c | h | c \rangle - \langle d | h | d \rangle - \sum_{a \neq c, d} \langle ac || ac \rangle - \sum_{a \neq c, d} \langle ad || ad \rangle - \langle cd || cd \rangle \\ &= -\langle c | h | c \rangle - \langle d | h | d \rangle - \left(\sum_{a \neq c} \langle ac || ac \rangle - \langle dc || dc \rangle \right) - \left(\sum_{a \neq d} \langle ad || ad \rangle - \langle cd || cd \rangle \right) - \langle cd || cd \rangle \\ &= -\varepsilon_c - \varepsilon_d + \langle cd || cd \rangle - \langle cd || dc \rangle\end{aligned}\quad (3.3.3)$$

Ex 3.6

$$\begin{aligned}
{}^N E_0 - {}^{N+1} E^r &= \sum_a \langle a | h | a \rangle + \frac{1}{2} \sum_a \sum_b \langle ab || ab \rangle \\
&\quad - \left[\sum_a \langle a | h | a \rangle + \langle r | h | r \rangle + \frac{1}{2} \sum_a \sum_b \langle ab || ab \rangle + \frac{1}{2} \sum_b \langle rb || rb \rangle + \frac{1}{2} \sum_a \langle ar || ar \rangle \right] \\
&= - \langle r | h | r \rangle - \frac{1}{2} \sum_b \langle rb || rb \rangle - \frac{1}{2} \sum_b \langle br || br \rangle \\
&= - \langle r | h | r \rangle - \sum_b \langle rb || rb \rangle
\end{aligned} \tag{3.3.4}$$

3.3.2 Brillouin's Theorem

3.3.3 The HF Hamiltonian

Ex 3.7 Suppose \mathcal{H}_0 commutes with \mathcal{P}_n ,

$$\begin{aligned}
\mathcal{H}_0 |\Psi_0\rangle &= \mathcal{H}_0 \frac{1}{\sqrt{N!}} \sum_n (-1)^{p_n} \mathcal{P}_n \left\{ \sum_i^N f(i) \chi_j(1) \cdots \chi_k(N) \right\} \\
&= \frac{1}{\sqrt{N!}} \sum_n (-1)^{p_n} \mathcal{P}_n \{ (\varepsilon_j + \cdots + \varepsilon_k) \chi_j(1) \cdots \chi_k(N) \} \\
&= \sum_a \varepsilon_a
\end{aligned} \tag{3.3.5}$$

Now we show \mathcal{H}_0 commutes with \mathcal{P}_n , for example, \mathcal{P}_{ab}

$$\mathcal{P}_{ab} \mathcal{H}_0 = \mathcal{P}_{ab} (\cdots + f(a) + \cdots + f(b) + \cdots) = (\cdots + f(b) + \cdots + f(a) + \cdots) \mathcal{P}_{ab} = \mathcal{H}_0 \mathcal{P}_{ab} \tag{3.3.6}$$

Ex 3.8

$$\mathcal{V} = \sum_i^N \sum_{j>i}^N \mathcal{O}_2 - \sum_i^N \sum_b^N [\mathcal{G}_b(i) - \mathcal{K}_b(i)] \tag{3.3.7}$$

thus

$$\begin{aligned}
\langle \Psi_0 | \mathcal{V} | \Psi_0 \rangle &= \sum_i^N \sum_{j>i}^N \langle \Psi_0 | \mathcal{O}_2 | \Psi_0 \rangle - \sum_i^N \sum_b^N [\langle \Psi_0 | \mathcal{G}_b(i) - \mathcal{K}_b(i) | \Psi_0 \rangle] \\
&= \frac{1}{2} \sum_a^N \sum_b^N \langle ab || ab \rangle - \sum_i^N \sum_b^N [\langle ib | ib \rangle - \langle ib | bi \rangle] \\
&= -\frac{1}{2} \sum_a^N \sum_b^N \langle ab || ab \rangle
\end{aligned} \tag{3.3.8}$$

3.4 Restricted Closed-shell HF: The Roothaan Equations

3.4.1 Closed-shell HF: Restricted Spin Orbitals

Ex 3.9

$$\begin{aligned}
\varepsilon_i &= (i|h|i) + \sum_b^N (\langle ib | ib \rangle - \langle ib | bi \rangle) \\
&= (i|h|i) + \sum_c^{N/2} (\langle ic | ic \rangle - \langle ic | ci \rangle) + \sum_{\bar{c}}^{N/2} (\langle i\bar{c} | i\bar{c} \rangle - \langle i\bar{c} | \bar{c}i \rangle)
\end{aligned} \tag{3.4.1}$$

Assume χ_j has α spin, since assuming α or β is identical

$$\begin{aligned}
\varepsilon_i &= (i|h|i) + \sum_c^{N/2} [(ic|ic) \langle \alpha | \alpha \rangle \langle \alpha | \alpha \rangle - (ic|ci) \langle \alpha | \alpha \rangle \langle \alpha | \alpha \rangle] + \sum_c^{N/2} [(ic|ic) \langle \alpha | \alpha \rangle \langle \beta | \beta \rangle - (ic|ci) \langle \alpha | \beta \rangle \langle \beta | \alpha \rangle] \\
&= (i|h|i) + \sum_c^{N/2} [2(ic|ic) - (ic|ci)] \\
&= (i|h|i) + \sum_n^{N/2} (2J_{ib} - K_{ib})
\end{aligned} \tag{3.4.2}$$

3.4.2 Introduction of a Basis: The Roothaan Equations

Ex 3.10

$$\begin{aligned}
(\mathbf{C}^\dagger \mathbf{S} \mathbf{C})_{\mu\nu} &= \sum_i \sum_j C_{\mu i}^\dagger S_{ij} C_{j\nu} \\
&= \sum_i \sum_j C_{i\mu}^* \langle \phi_i | \phi_j \rangle C_{j\nu} \\
&= \langle \phi_\mu | \phi_\nu \rangle \\
&= \delta_{\mu\nu}
\end{aligned} \tag{3.4.3}$$

thus

$$\mathbf{C}^\dagger \mathbf{S} \mathbf{C} = \mathbf{1} \tag{3.4.4}$$

3.4.3 The Charge Density

Ex 3.11

$$\begin{aligned}
\rho(\mathbf{r}) &= \langle \Psi_0 | \hat{\rho}(\mathbf{r}) | \Psi_0 \rangle \\
&= \sum_i^N \frac{1}{N!} \sum_I^{N!} \sum_J^{N!} (-1)^{p_I} (-1)^{p_J} \int d\mathbf{x}_1 \cdots d\mathbf{x}_N \hat{\mathcal{P}}_I \{ \chi_1(1) \cdots \chi_N(N) \}^* \delta(\mathbf{r}_i - \mathbf{r}) \hat{\mathcal{P}}_J \{ \chi_1(1) \cdots \chi_N(N) \}
\end{aligned} \tag{3.4.5}$$

Since $\{\chi_m\}$ are orthogonal,

$$\begin{aligned}
\rho(\mathbf{r}) &= \sum_i^N \frac{1}{N!} \sum_I^{N!} \int d\mathbf{x}_1 \cdots d\mathbf{x}_N \hat{\mathcal{P}}_I \{ \chi_1(1) \cdots \chi_N(N) \}^* \delta(\mathbf{r}_i - \mathbf{r}) \hat{\mathcal{P}}_I \{ \chi_1(1) \cdots \chi_N(N) \} \\
&= \sum_i^N \frac{1}{N!} (N-1)! \sum_s^N \int d\mathbf{x}_i \chi_s^*(\mathbf{x}_i) \delta(\mathbf{r}_i - \mathbf{r}) \chi_s(\mathbf{x}_i) \\
&= \sum_i^N \frac{1}{N} \cdot 2 \sum_s^{N/2} \int d\mathbf{r}_i \phi_s(\mathbf{r}_i) \delta(\mathbf{r}_i - \mathbf{r}) \phi_s(\mathbf{r}_i) \\
&= \sum_i^N \frac{2}{N} \sum_s^{N/2} \phi_s(\mathbf{r}) \phi_s(\mathbf{r}) \\
&= N \frac{2}{N} \sum_s^{N/2} \phi_s(\mathbf{r}) \phi_s(\mathbf{r}) \\
&= 2 \sum_s^{N/2} \phi_s(\mathbf{r}) \phi_s(\mathbf{r})
\end{aligned} \tag{3.4.6}$$

Ex 3.12 From Ex 3.10, we have

$$\mathbf{C}^\dagger \mathbf{S} \mathbf{C} = \mathbf{1} \tag{3.4.7}$$

i.e.

$$\sum_i^K \sum_j^K C_{i\mu}^* S_{ij} C_{j\nu} = \delta_{\mu\nu} \quad (3.4.8)$$

thus

$$\begin{aligned} (\mathbf{PSP})_{\mu\sigma} &= \sum_{\nu}^K \sum_{\lambda}^K P_{\mu\nu} S_{\nu\lambda} P_{\lambda\sigma} \\ &= 4 \sum_{\nu}^K \sum_{\lambda}^K \sum_a^{N/2} C_{\mu a} C_{\nu a}^* S_{\nu\lambda} \sum_b^{N/2} C_{\lambda b} C_{\sigma b}^* \\ &= 4 \sum_a^{N/2} \sum_b^{N/2} C_{\mu a} \left(\sum_{\nu}^K \sum_{\lambda}^K C_{\nu a}^* S_{\nu\lambda} C_{\lambda b} \right) C_{\sigma b}^* \\ &= 4 \sum_a^{N/2} \sum_b^{N/2} C_{\mu a} \delta_{ab} C_{\sigma b}^* \\ &= 4 \sum_a^{N/2} C_{\mu a} C_{\sigma a}^* \\ &= 2P_{\mu\sigma} \end{aligned} \quad (3.4.9)$$

thus

$$\mathbf{PSP} = 2\mathbf{P} \quad (3.4.10)$$

Ex 3.13 Eq. 3.122 shows

$$f(\mathbf{r}_1) = h(\mathbf{r}_1) + \sum_a^{N/2} \int d\mathbf{r}_2 \psi_a^*(\mathbf{r}_2) (2 - \hat{\mathcal{P}}_{12}) r_{12}^{-1} \psi_a(\mathbf{r}_2) \quad (3.4.11)$$

thus

$$\begin{aligned} f(\mathbf{r}_1) &= h(\mathbf{r}_1) + \sum_a^{N/2} \int d\mathbf{r}_2 \sum_{\sigma} C_{\sigma a}^* \phi_{\sigma}^*(\mathbf{r}_2) (2 - \hat{\mathcal{P}}_{12}) r_{12}^{-1} \sum_{\lambda} C_{\lambda a} \phi_{\lambda}(\mathbf{r}_2) \\ &= h(\mathbf{r}_1) + \sum_{\sigma} \sum_{\lambda} \left(\sum_a^{N/2} C_{\sigma a}^* C_{\lambda a} \right) \int d\mathbf{r}_2 \phi_{\sigma}^*(\mathbf{r}_2) (2 - \hat{\mathcal{P}}_{12}) r_{12}^{-1} \phi_{\lambda}(\mathbf{r}_2) \\ &= h(\mathbf{r}_1) + \frac{1}{2} \sum_{\sigma, \lambda} P_{\lambda\sigma} \int d\mathbf{r}_2 \phi_{\sigma}^*(\mathbf{r}_2) (2 - \hat{\mathcal{P}}_{12}) r_{12}^{-1} \phi_{\lambda}(\mathbf{r}_2) \end{aligned} \quad (3.4.12)$$

3.4.4 Expression for the Fock Matrix

Ex 3.14 In expression $(\mu\nu|\lambda\sigma)$, there are three interchangeable pairs, i.e. $\mu \leftrightarrow \nu$, $\lambda \leftrightarrow \sigma$, and $\mu\nu \leftrightarrow \lambda\sigma$. Thus $(\mu\nu|\lambda\sigma)$ has an 8-fold symmetry. Similarly, $(\mu\mu|\lambda\sigma)$, $(\mu\nu|\mu\lambda)$, $(\mu\nu|\mu\nu)$, $(\mu\mu|\sigma\sigma)$ has 2-fold symmetry, and $(\mu\mu|\mu\nu)$, $(\mu\mu|\mu\mu)$ has 1-fold symmetry.

Therefore, the number of unique 2e integrals is

expression	number	$K = 100$
$(\mu\nu \lambda\sigma)$	$K(K-1)(K-2)(K-3)/8$	11763675
$(\mu\mu \lambda\sigma)$	$K(K-1)(K-2)/2$	485100
$(\mu\nu \mu\lambda)$	$K(K-1)(K-2)/2$	485100
$(\mu\nu \mu\nu)$	$K(K-1)/2$	4950
$(\mu\mu \sigma\sigma)$	$K(K-1)/2$	4950
$(\mu\mu \mu\nu)$	$K(K-1)$	9900
$(\mu\mu \mu\mu)$	K	100

thus the total number is 12 753 775.

3.4.5 Orthogonalization of the Basis

Ex 3.15 \therefore

$$\mathbf{U}^\dagger \mathbf{S} \mathbf{U} = \mathbf{s} \quad (3.4.13)$$

\therefore

$$\mathbf{S} \mathbf{U} = \mathbf{U} \mathbf{s} \quad (3.4.14)$$

i.e.

$$\sum_{\nu} S_{\mu\nu} U_{\nu i} = U_{\mu i} s_i \quad (3.4.15)$$

thus

$$\sum_{\mu} U_{\mu i}^* \sum_{\nu} S_{\mu\nu} U_{\nu i} = \sum_{\mu} U_{\mu i}^* U_{\mu i} s_i \quad (3.4.16)$$

$$\sum_{\mu} \sum_{\nu} U_{\mu i}^* \langle \phi_{\mu} | \phi_{\nu} \rangle U_{\nu i} = s_i \sum_{\mu} |U_{\mu i}|^2 \quad (3.4.17)$$

Suppose

$$\phi'_i = \sum_{\nu} U_{\nu i} \phi_{\nu} \quad (3.4.18)$$

thus

$$\langle \phi'_i | \phi'_i \rangle = s_i \sum_{\mu} |U_{\mu i}|^2 \quad (3.4.19)$$

\therefore

$$\langle \phi'_i | \phi'_i \rangle > 0 \quad |U_{\mu i}|^2 > 0 \quad (3.4.20)$$

\therefore

$$s_i > 0 \quad (3.4.21)$$

Ex 3.16

- (3.174)

Since (ϕ, ϕ', ψ) are row vectors

$$\psi = \phi \mathbf{C} \quad (3.4.22)$$

$$\psi = \phi' \mathbf{C}' = \phi \mathbf{X} \mathbf{C}' \quad (3.4.23)$$

we have

$$\mathbf{C} = \mathbf{X} \mathbf{C}' \quad (3.4.24)$$

i.e.

$$\mathbf{C}' = \mathbf{X}^{-1} \mathbf{C} \quad (3.4.25)$$

- (3.177)

$$\begin{aligned} F'_{\mu\nu} &= \langle \phi'_{\mu} | f | \phi'_{\nu} \rangle \\ &= \left\langle \sum_i \phi_i X_{i\mu} \left| f \right| \sum_j \phi_j X_{j\nu} \right\rangle \\ &= \sum_i \sum_j X_{i\mu}^* X_{j\nu} \langle \phi_i | f | \phi_j \rangle \\ &= \sum_i \sum_j X_{i\mu}^* F_{ij} X_{j\nu} \end{aligned} \quad (3.4.26)$$

i.e.

$$\mathbf{F}' = \mathbf{X}^\dagger \mathbf{F} \mathbf{X} \quad (3.4.27)$$

3.4.6 The SCF Procedure

3.4.7 Expectation Values and Population Analysis

Ex 3.17 From (3.148) in the textbook, we get

$$F_{\mu\nu} = H_{\mu\nu}^{\text{core}} + G_{\mu\nu} = H_{\mu\nu}^{\text{core}} + \sum_a^{N/2} [2(\mu\nu|aa) - (\mu a|a\nu)] \quad (3.4.28)$$

thus

$$\begin{aligned} E_0 &= \sum_a^{N/2} [2h_{aa} + \sum_b^{N/2} (2J_{ab} - K_{ab})] \\ &= 2 \sum_a^{N/2} (a|h|a) + \sum_a^{N/2} \sum_b^{N/2} [2(aa|bb) - (ab|ba)] \\ &= 2 \sum_a^{N/2} \sum_\mu \sum_\nu C_{\mu a}^* C_{\nu a} (\mu|h|\nu) + \sum_a^{N/2} \sum_b^{N/2} \left[2 \sum_\mu \sum_\nu C_{\mu a}^* C_{\nu a} (\mu\nu|bb) - \sum_\mu \sum_\nu C_{\mu a}^* C_{\nu a} (\mu b|b\nu) \right] \\ &= \sum_\mu \sum_\nu P_{\nu\mu} H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_b^{N/2} \sum_\mu \sum_\nu [2P_{\nu\mu} (\mu\nu|bb) - P_{\nu\mu} (\mu b|b\nu)] \\ &= \sum_\mu \sum_\nu P_{\nu\mu} [H_{\mu\nu}^{\text{core}} + \frac{1}{2} G_{\mu\nu}] \\ &= \frac{1}{2} \sum_\mu \sum_\nu P_{\nu\mu} [H_{\mu\nu}^{\text{core}} + F_{\mu\nu}] \end{aligned} \quad (3.4.29)$$

Ex 3.18 For symmetrically orthogonalized basis,

$$\mathbf{C}' = \mathbf{S}^{1/2} \mathbf{C} \quad (3.4.30)$$

thus

$$\begin{aligned} P'_{\mu\nu} &= 2 \sum_a^{N/2} C'_{\mu a} C'^*_{\nu a} \\ &= 2 \sum_a^{N/2} \sum_i S_{\mu i}^{1/2} C_{ia} \sum_j S_{\nu j}^{1/2*} C_{ja}^* \\ &= \sum_i \sum_j S_{\mu i}^{1/2} \left(2 \sum_a^{N/2} C_{ia} C_{ja}^* \right) S_{\nu j}^{1/2*} \\ &= \sum_i \sum_j S_{\mu i}^{1/2} P_{ij} S_{\nu j}^{1/2*} \\ &= \sum_i \sum_j S_{\mu i}^{1/2} P_{ij} S_{j\nu}^{1/2} \end{aligned} \quad (3.4.31)$$

i.e.

$$\mathbf{P}' = \mathbf{S}^{1/2} \mathbf{P} \mathbf{S}^{1/2} \quad (3.4.32)$$

thus

$$\sum_\mu (\mathbf{S}^{1/2} \mathbf{P} \mathbf{S}^{1/2})_{\mu\mu} = \sum_\mu \mathbf{P}'_{\mu\mu} \quad (3.4.33)$$

3.5 Model Calculations on H₂ and HeH⁺

3.5.1 The 1s Minimal STO-3G Basis Set

Ex 3.19

$$\begin{aligned}
\phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_A) \phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_B) &= \left(\frac{2\alpha}{\pi}\right)^{3/4} e^{-\alpha|\mathbf{r}-\mathbf{R}_A|^2} \left(\frac{2\beta}{\pi}\right)^{3/4} e^{-\beta|\mathbf{r}-\mathbf{R}_B|^2} \\
&= \left(\frac{2\alpha}{\pi}\right)^{3/4} \left(\frac{2\beta}{\pi}\right)^{3/4} e^{-\alpha|\mathbf{r}-\mathbf{R}_A|^2 - \beta|\mathbf{r}-\mathbf{R}_B|^2} \\
&= \left(\frac{2\alpha}{\pi}\right)^{3/4} \left(\frac{2\beta}{\pi}\right)^{3/4} \exp\left(-\left[(\alpha + \beta)|\mathbf{r}|^2 - 2\mathbf{r} \cdot (\alpha\mathbf{R}_A + \beta\mathbf{R}_B) + \alpha|\mathbf{R}_A|^2 + \beta|\mathbf{R}_B|^2\right]\right)
\end{aligned} \tag{3.5.1}$$

Let

$$p = \alpha + \beta \quad \mathbf{R}_P = \frac{\alpha\mathbf{R}_A + \beta\mathbf{R}_B}{\alpha + \beta} \tag{3.5.2}$$

we have

$$\begin{aligned}
\phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_A) \phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_B) &= \left(\frac{2\alpha}{\pi}\right)^{3/4} \left(\frac{2\beta}{\pi}\right)^{3/4} \exp\left(-\left[p|\mathbf{r}|^2 - 2\mathbf{r} \cdot (p\mathbf{R}_P) + \alpha|\mathbf{R}_A|^2 + \beta|\mathbf{R}_B|^2\right]\right) \\
&= \left(\frac{2\alpha}{\pi}\right)^{3/4} \left(\frac{2\beta}{\pi}\right)^{3/4} \exp\left(-\left[p|\mathbf{r} - \mathbf{R}_P|^2 - p|\mathbf{R}_P|^2 + \alpha|\mathbf{R}_A|^2 + \beta|\mathbf{R}_B|^2\right]\right) \\
&= \left(\frac{2\alpha\beta/p}{\pi}\right)^{3/4} \left(\frac{2p}{\pi}\right)^{3/4} e^{-p|\mathbf{r}-\mathbf{R}_P|^2} \exp\left(p|\mathbf{R}_P|^2 - \alpha|\mathbf{R}_A|^2 - \beta|\mathbf{R}_B|^2\right)
\end{aligned} \tag{3.5.3}$$

Let

$$\phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_A) \phi_{1s}^{\text{GF}}(\alpha, \mathbf{r} - \mathbf{R}_B) = K_{AB} \left(\frac{2p}{\pi}\right)^{3/4} e^{-p|\mathbf{r}-\mathbf{R}_P|^2} \tag{3.5.4}$$

thus

$$\begin{aligned}
K_{AB} &= \left(\frac{2\alpha\beta/p}{\pi}\right)^{3/4} \exp\left(p|\mathbf{R}_P|^2 - \alpha|\mathbf{R}_A|^2 - \beta|\mathbf{R}_B|^2\right) \\
&= \left(\frac{2\alpha\beta/p}{\pi}\right)^{3/4} \exp\left(\frac{1}{p}(\alpha^2|\mathbf{R}_A|^2 + \beta^2|\mathbf{R}_B|^2 + 2\alpha\beta\mathbf{R}_A \cdot \mathbf{R}_B) - \alpha|\mathbf{R}_A|^2 - \beta|\mathbf{R}_B|^2\right) \\
&= \left(\frac{2\alpha\beta/p}{\pi}\right)^{3/4} \exp\left(\frac{1}{p}(\alpha^2|\mathbf{R}_A|^2 + \beta^2|\mathbf{R}_B|^2 + 2\alpha\beta\mathbf{R}_A \cdot \mathbf{R}_B - p\alpha|\mathbf{R}_A|^2 - p\beta|\mathbf{R}_B|^2)\right) \\
&= \left(\frac{2\alpha\beta/p}{\pi}\right)^{3/4} \exp\left(\frac{1}{p}(-\alpha\beta|\mathbf{R}_A|^2 - \alpha\beta|\mathbf{R}_B|^2 + 2\alpha\beta\mathbf{R}_A \cdot \mathbf{R}_B)\right) \\
&= \left(\frac{2\alpha\beta}{p\pi}\right)^{3/4} \exp\left(-\frac{\alpha\beta}{p}|\mathbf{R}_A - \mathbf{R}_B|^2\right)
\end{aligned} \tag{3.5.5}$$

Ex 3.20 At $r = 0$,

$$\phi_{1s}^{\text{CGF}}(\zeta = 1.0, \text{STO-1G}) = 0.267\,656 \tag{3.5.6}$$

$$\phi_{1s}^{\text{CGF}}(\zeta = 1.0, \text{STO-2G}) = 0.389\,383 \tag{3.5.7}$$

$$\phi_{1s}^{\text{CGF}}(\zeta = 1.0, \text{STO-3G}) = 0.454\,986 \tag{3.5.8}$$

while

$$\phi_{1s}^{\text{SF}}(\zeta = 1.0) = \frac{1}{\sqrt{\pi}} = 0.564\,19 \tag{3.5.9}$$

3.5.2 STO-3G H₂

Ex 3.21

$$\phi_{1s}^{\text{CGF}}(\zeta = 1.0, \text{STO-1G}) = \phi_{1s}^{\text{GF}}(0.270\,950) \quad (3.5.10)$$

Since $\alpha = \alpha_{(\zeta=1.0)} \times \zeta^2$,

$$\phi_{1s}^{\text{CGF}}(\zeta = 1.24, \text{STO-1G}) = \phi_{1s}^{\text{GF}}(0.416\,613) \quad (3.5.11)$$

thus

$$\begin{aligned} S_{12} &= K_{AB} \left(\frac{2 \cdot 2\alpha}{\pi} \right)^{3/4} \int d\mathbf{r} e^{-2\alpha|\mathbf{r}-\mathbf{R}_P|^2} \\ &= \left(\frac{2\alpha}{2\pi} \right)^{3/4} e^{-\frac{\alpha}{2}R^2} \left(\frac{2 \cdot 2\alpha}{\pi} \right)^{3/4} \int d\mathbf{r} e^{-2\alpha|\mathbf{r}-\mathbf{R}_A|^2} \\ &= \left(\frac{2\alpha}{\pi} \right)^{3/2} e^{-\frac{\alpha}{2}R^2} 4\pi \int dr r^2 e^{-2\alpha r^2} \\ &= \left(\frac{2\alpha}{\pi} \right)^{3/2} e^{-\frac{\alpha}{2}R^2} 4\pi \frac{\sqrt{\pi}}{8\sqrt{2}\alpha^{3/2}} \\ &= e^{-\frac{\alpha}{2}R^2} \end{aligned} \quad (3.5.12)$$

At $R = 1.4, \alpha = 0.416\,613$,

$$S_{12} = 0.6648 \quad (3.5.13)$$

Ex 3.22 Let

$$\psi_1 = c_1(\phi_1 + \phi_2) \quad \psi_2 = c_2(\phi_1 - \phi_2) \quad (3.5.14)$$

$$\begin{aligned} 1 &= \langle \phi_1 | \psi_1 \rangle = c_1^2(S_{11} + S_{12} + S_{21} + S_{22}) \\ &= c_1^2(2 + 2S_{12}) \end{aligned} \quad (3.5.15)$$

\therefore

$$c_1 = [2(1 + S_{12})]^{-1/2} \quad (3.5.16)$$

$$\begin{aligned} 1 &= \langle \phi_2 | \psi_2 \rangle = c_2^2(S_{11} - S_{12} - S_{21} + S_{22}) \\ &= c_2^2(2 - 2S_{12}) \end{aligned} \quad (3.5.17)$$

\therefore

$$c_2 = [2(1 - S_{12})]^{-1/2} \quad (3.5.18)$$

Ex 3.23 Suppose

$$\psi_1 = c_1(\phi_1 + \phi_2) \quad \psi_2 = c_2(\phi_1 - \phi_2) \quad (3.5.19)$$

thus

$$\mathbf{H}^{\text{core}} \mathbf{C} = \mathbf{S} \mathbf{C} \boldsymbol{\varepsilon} \quad (3.5.20)$$

$$\begin{pmatrix} H_{11}^{\text{core}} & H_{12}^{\text{core}} \\ H_{21}^{\text{core}} & H_{22}^{\text{core}} \end{pmatrix} \begin{pmatrix} c_1 & c_2 \\ c_1 & -c_2 \end{pmatrix} = \begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix} \begin{pmatrix} c_1 & c_2 \\ c_1 & -c_2 \end{pmatrix} \begin{pmatrix} \varepsilon_1 & 0 \\ 0 & \varepsilon_2 \end{pmatrix} \quad (3.5.21)$$

$$\begin{pmatrix} (H_{11}^{\text{core}} + H_{12}^{\text{core}})c_1 & (H_{11}^{\text{core}} - H_{12}^{\text{core}})c_2 \\ (H_{21}^{\text{core}} + H_{22}^{\text{core}})c_1 & (H_{21}^{\text{core}} - H_{22}^{\text{core}})c_2 \end{pmatrix} = \begin{pmatrix} (S_{11} + S_{12})c_1\varepsilon_1 & (S_{11} - S_{12})c_2\varepsilon_2 \\ (S_{21} + S_{22})c_1\varepsilon_1 & (S_{21} - S_{22})c_2\varepsilon_2 \end{pmatrix} \quad (3.5.22)$$

\therefore

$$\begin{cases} \varepsilon_1 = (H_{11}^{\text{core}} + H_{12}^{\text{core}})/(1 + S_{12}) \\ \varepsilon_2 = (H_{11}^{\text{core}} - H_{12}^{\text{core}})/(1 - S_{12}) \end{cases} \quad (3.5.23)$$

$$\varepsilon_1 = (-1.1204 - 0.9584)/(1 + 0.6593) = -1.2528 \quad (3.5.24)$$

$$\varepsilon_2 = (-1.1204 + 0.9584)/(1 - 0.6593) = -0.4755 \quad (3.5.25)$$

Ex 3.24

Ex 3.25

Ex 3.26

Ex 3.27

3.5.3 An SCF Calculation on STO-3G HeH⁺

Ex 3.28

$$\begin{aligned}
\mathbf{X}_{\text{Schmidt}}^\dagger \mathbf{S} \mathbf{X}_{\text{Schmidt}} &= \begin{pmatrix} 1 & 0 \\ -S_{12}/\sqrt{1-S_{12}^2} & 1/\sqrt{1-S_{12}^2} \end{pmatrix} \begin{pmatrix} 1 & S_{12} \\ S_{12} & 1 \end{pmatrix} \begin{pmatrix} 1 & -S_{12}/\sqrt{1-S_{12}^2} \\ 0 & 1/\sqrt{1-S_{12}^2} \end{pmatrix} \\
&= \begin{pmatrix} 1 & S_{12} \\ 0 & \sqrt{1-S_{12}^2} \end{pmatrix} \begin{pmatrix} 1 & -S_{12}/\sqrt{1-S_{12}^2} \\ 0 & 1/\sqrt{1-S_{12}^2} \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}
\end{aligned} \tag{3.5.26}$$

thus the Schmidt transformation produces orthonormal basis.

Ex 3.29

$$E_0(R \rightarrow \infty) = \frac{1}{2} \sum_{\mu} \sum_{\nu} P_{\nu\mu}(R \rightarrow \infty) [2H_{\mu\nu}^{\text{core}} + G_{\mu\nu}] \tag{3.5.27}$$

where

$$P_{\nu\mu}(R \rightarrow \infty) = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} \tag{3.5.28}$$

$$\begin{aligned}
G_{\mu\nu} &= \sum_{\lambda} \sum_{\sigma} P_{\lambda\sigma}(R \rightarrow \infty) \left[(\mu\nu|\sigma\lambda) - \frac{1}{2}(\mu\lambda|\sigma\nu) \right] \\
&= 2 \left[(\mu\nu|\phi_1\phi_1) - \frac{1}{2}(\mu\phi_1|\phi_1\nu) \right]
\end{aligned} \tag{3.5.29}$$

thus

$$\begin{aligned}
E_0(R \rightarrow \infty) &= \frac{1}{2} \sum_{\mu} \sum_{\nu} P_{\nu\mu}(R \rightarrow \infty) [2H_{\mu\nu}^{\text{core}} + G_{\mu\nu}] \\
&= \frac{1}{2} \times 2 [2H_{11}^{\text{core}} + G_{11}] \\
&= 2(T_{11} + V_{11}^1) + 2 \left[(\phi_1\phi_1|\phi_1\phi_1) - \frac{1}{2}(\phi_1\phi_1|\phi_1\phi_1) \right] \\
&= 2T_{11} + 2V_{11}^1 + (\phi_1\phi_1|\phi_1\phi_1)
\end{aligned} \tag{3.5.30}$$

3.6 Polyatomic Basis Sets

3.6.1 Contracted Gaussian Functions

3.6.2 Minimal Basis Sets: STO-3G

3.6.3 Double Zeta Basis Sets: 4-31G

Ex 3.30 The outer basis function

$$\phi_{1s}''(\mathbf{r}) = g_{1s}(0.298073, \mathbf{r}) \tag{3.6.1}$$

The inner basis function

$$\phi_{1s}'(\mathbf{r}) = N[0.46954g_{1s}(1.242567, \mathbf{r}) + 0.15457g_{1s}(5.782948, \mathbf{r}) + 0.02373g_{1s}(38.47497, \mathbf{r})] \tag{3.6.2}$$

Renormalize it, we get

$$N = 1.689 \tag{3.6.3}$$

thus

$$\phi_{1s}'(\mathbf{r}) = 0.79330g_{1s}(1.242567, \mathbf{r}) + 0.26115g_{1s}(5.782948, \mathbf{r}) + 0.04009g_{1s}(38.47497, \mathbf{r}) \tag{3.6.4}$$

3.6.4 Polarized Basis Sets: 6-31G* and 6-31G**

Ex 3.31

	C	H	total
STO-3G	5	1	36
4-31G	9	2	66
6-31G* (Cartesian)	15	2	102
6-31G** (Cartesian)	15	5	120

3.7 Some Illustrative Closed-shell Calculations

3.7.1 Total Energies

Ex 3.32

Reaction I

basis	$\Delta E/$ a.u.	$\Delta E/(\text{kcal/mol})$	
STO-3G	-0.061	-38.28	exoergic
4-31G	-0.069	-43.30	
6-31G*	-0.045	-28.24	
6-31G**	-0.055	-34.51	
HF-limit	-0.051	-32.00	

Reaction II

basis	$\Delta E/$ a.u.	$\Delta E/(\text{kcal/mol})$	
STO-3G	0.186	116.72	endoergic
4-31G	-0.114	-71.54	exoergic
6-31G*	-0.088	-55.22	
6-31G**	-0.095	-59.61	
HF-limit	-0.097	-60.87	

The contribution of zero-point vibrations to the energy change of reaction I would be -0.37 kcal/mol, to the energy change of reaction II would be 17.78 kcal/mol. Thus the effect of zero-point vibrations should not be ignored.

3.7.2 Ionization Potentials

3.7.3 Equilibrium Geometries

3.7.4 Population Analysis and Dipole Moments

3.8 Unrestricted Open-shell HF: The Pople-Nesbet Equations

3.8.1 Open-shell HF: Unrestricted Spin Orbitals

Ex 3.33

$$\begin{aligned}
 f^\alpha(1) &= \int d\omega_1 \alpha^*(\omega_1) \left[h(1) + \sum_a \int d\mathbf{x}_2 \chi_a^*(2) r_{12}^{-1} (1 - \hat{\mathcal{P}}_{12}) \chi_a(2) \right] \alpha(\omega_1) \\
 &= h(1) + \sum_a^{N_\alpha} \left[\int d\omega_1 \alpha^*(\omega_1) \int d\mathbf{x}_2 \chi_a^*(2) r_{12}^{-1} \chi_a(2) \alpha(\omega_1) - \int d\omega_1 \alpha^*(\omega_1) \int d\mathbf{x}_2 \chi_a^*(2) r_{12}^{-1} \chi_a(1) \alpha(\omega_2) \right] \\
 &\quad + \sum_a^{N_\beta} \left[\int d\omega_1 \alpha^*(\omega_1) \int d\mathbf{x}_2 \chi_a^*(2) r_{12}^{-1} \chi_a(2) \alpha(\omega_1) - \int d\omega_1 \alpha^*(\omega_1) \int d\mathbf{x}_2 \chi_a^*(2) r_{12}^{-1} \chi_a(1) \alpha(\omega_2) \right] \\
 &= h(1) + \sum_a^{N_\alpha} \left[\int d\mathbf{r}_2 \psi_a^{\alpha*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\alpha(\mathbf{r}_2) - \int d\mathbf{r}_2 \int d\omega_2 \int d\omega_1 \alpha^*(\omega_1) \alpha^*(\omega_2) \psi_a^{\alpha*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\alpha(\mathbf{r}_1) \alpha(\omega_1) \alpha(\omega_2) \right]
 \end{aligned}$$

$$\begin{aligned}
& + \sum_a^{N_\beta} \left[\int d\mathbf{r}_2 \psi_a^{\beta*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\beta(\mathbf{r}_2) - \int d\mathbf{r}_2 \int d\omega_2 \int d\omega_1 \alpha^*(\omega_1) \beta^*(\omega_2) \psi_a^{\beta*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\beta(\mathbf{r}_1) \beta(\omega_1) \alpha(\omega_2) \right] \\
& = h(1) + \sum_a^{N_\alpha} \left[\int d\mathbf{r}_2 \psi_a^{\alpha*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\alpha(\mathbf{r}_2) - \int d\mathbf{r}_2 \psi_a^{\alpha*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\alpha(\mathbf{r}_1) \right] + \sum_a^{N_\beta} \left[\int d\mathbf{r}_2 \psi_a^{\beta*}(\mathbf{r}_2) r_{12}^{-1} \psi_a^\beta(\mathbf{r}_2) - 0 \right] \\
& = h(1) + \sum_a^{N_\alpha} [J_a^\alpha(1) - K_a^\alpha(1)] + \sum_a^{N_\beta} J_a^\beta(1)
\end{aligned} \tag{3.8.1}$$

Ex 3.34

$$\begin{aligned}
E_0 & = \sum_a h_{aa} + \frac{1}{2} \sum_a^{N_\alpha} \sum_b^{N_\alpha} (J_{ab}^{\alpha\alpha} - K_{ab}^{\alpha\alpha}) + \sum_a^{N_\alpha} \sum_b^{N_\beta} J_{ab}^{\alpha\beta} \\
& = h_{11}^\alpha + h_{22}^\alpha + h_{11}^\beta + J_{12}^{\alpha\alpha} - K_{12}^{\alpha\alpha} + J_{11}^{\alpha\beta} + J_{21}^{\alpha\beta}
\end{aligned} \tag{3.8.2}$$

Ex 3.35

$$\begin{aligned}
\varepsilon_i^\alpha & = (\psi_i^\alpha(1) | h(1) + \sum_a^{N_\alpha} [J_a^\alpha(1) - K_a^\alpha(1)] + \sum_a^{N_\beta} J_a^\beta(1) | \psi_i^\alpha(1)) \\
& = h_{ii}^\alpha + \sum_a^{N_\alpha} [J_{ia}^{\alpha\alpha} - K_{ia}^{\alpha\alpha}] + \sum_a^{N_\beta} J_{ia}^{\alpha\beta}
\end{aligned} \tag{3.8.3}$$

$$\begin{aligned}
\varepsilon_i^\beta & = (\psi_i^\beta(1) | h(1) + \sum_a^{N_\alpha} [J_a^\beta(1) - K_a^\beta(1)] + \sum_a^{N_\beta} J_a^\alpha(1) | \psi_i^\beta(1)) \\
& = h_{ii}^\beta + \sum_a^{N_\alpha} [J_{ia}^{\beta\alpha} - K_{ia}^{\beta\alpha}] + \sum_a^{N_\beta} J_{ia}^{\beta\beta}
\end{aligned} \tag{3.8.4}$$

Since

$$E_0 = \sum_a h_{aa} + \frac{1}{2} \sum_a^{N_\alpha} \sum_b^{N_\alpha} (J_{ab}^{\alpha\alpha} - K_{ab}^{\alpha\alpha}) + \frac{1}{2} \sum_a^{N_\beta} \sum_b^{N_\beta} (J_{ab}^{\beta\beta} - K_{ab}^{\beta\beta}) + \sum_a^{N_\alpha} \sum_b^{N_\beta} J_{ab}^{\alpha\beta} \tag{3.8.5}$$

we have

$$E_0 = \sum_i^{N_\alpha} \varepsilon_i^\alpha + \sum_i^{N_\beta} \varepsilon_i^\beta - \frac{1}{2} \sum_i^{N_\alpha} \sum_a^{N_\alpha} (J_{ia}^{\alpha\alpha} - K_{ia}^{\alpha\alpha}) - \frac{1}{2} \sum_i^{N_\beta} \sum_a^{N_\beta} (J_{ia}^{\beta\beta} - K_{ia}^{\beta\beta}) - \sum_i^{N_\beta} \sum_a^{N_\alpha} J_{ia}^{\beta\alpha} \tag{3.8.6}$$

3.8.2 Introduction of a Basis: The Pople-Nesbet Equations

3.8.3 Unrestricted Density Matrices

Ex 3.36

$$\int d\mathbf{r} \rho^S(\mathbf{r}) = \int d\mathbf{r} [\rho^\alpha(\mathbf{r}) - \rho^\beta(\mathbf{r})] \tag{3.8.7}$$

$$= N_\alpha - N_\beta \tag{3.8.8}$$

Since

$$\langle \hat{\mathcal{J}}_z \rangle = \frac{1}{2} (N_\alpha - N_\beta) \tag{3.8.9}$$

we get

$$\int d\mathbf{r} \rho^S(\mathbf{r}) = 2 \langle \hat{\mathcal{J}}_z \rangle \tag{3.8.10}$$

Ex 3.37

$$\begin{aligned}
\rho^\alpha(\mathbf{r}) &= \sum_a^{N_\alpha} \psi_a^{\alpha*}(\mathbf{r}) \psi_a^\alpha(\mathbf{r}) \\
&= \sum_a^{N_\alpha} \sum_\nu C_{\nu a}^{\alpha*} \phi_\nu^*(\mathbf{r}) \sum_\mu C_{\mu a}^\alpha \phi_\mu(\mathbf{r}) \\
&= \sum_\nu \sum_\mu \left[\sum_a^{N_\alpha} C_{\nu a}^{\alpha*} C_{\mu a}^\alpha \right] \phi_\nu^*(\mathbf{r}) \phi_\mu(\mathbf{r})
\end{aligned} \tag{3.8.11}$$

Let

$$P_{\mu\nu}^\alpha = \sum_a^{N_\alpha} C_{\nu a}^{\alpha*} C_{\mu a}^\alpha \tag{3.8.12}$$

thus

$$\rho^\alpha(\mathbf{r}) = \sum_\nu \sum_\mu P_{\mu\nu}^\alpha \phi_\mu(\mathbf{r}) \phi_\nu^*(\mathbf{r}) \tag{3.8.13}$$

The formulation for β spin is similar.

Ex 3.38

$$\begin{aligned}
\langle \mathcal{O}_1 \rangle &= \sum_i^N \langle \chi_i | h | \chi_i \rangle \\
&= \sum_i^{N_\alpha} (\psi_i^\alpha | h | \psi_i^\alpha) + \sum_i^{N_\beta} (\psi_i^\beta | h | \psi_i^\beta) \\
&= \sum_i^{N_\alpha} \sum_\nu \sum_\mu C_{\nu a}^{\alpha*} (\phi_\nu | h | \phi_\mu) C_{\mu a}^\alpha + \sum_i^{N_\beta} \sum_\nu \sum_\mu C_{\nu a}^{\beta*} (\phi_\nu | h | \phi_\mu) C_{\mu a}^\beta \\
&= \sum_\nu \sum_\mu P_{\mu\nu}^\alpha (\phi_\nu | h | \phi_\mu) + \sum_\nu \sum_\mu P_{\mu\nu}^\beta (\phi_\nu | h | \phi_\mu) \\
&= \sum_\nu \sum_\mu P_{\mu\nu}^T (\phi_\nu | h | \phi_\mu)
\end{aligned} \tag{3.8.14}$$

Ex 3.39

$$\begin{aligned}
\langle \hat{\rho}^S \rangle &= \langle \Psi_0 | \hat{\rho}^S | \Psi_0 \rangle \\
&= \frac{1}{N!} \sum_{i,j}^{N!} (-1)^{p_i} (-1)^{p_j} \int d\mathbf{x}_1 \cdots d\mathbf{x}_N \hat{\mathcal{P}}_i \{ \chi_1(1) \cdots \chi_N(N) \} \sum_m^N 2\delta(\mathbf{r}_m - \mathbf{R}) s_z(m) \hat{\mathcal{P}}_j \{ \chi_1(1) \cdots \chi_N(N) \} \\
&= \frac{2}{N!} \sum_i^{N!} \sum_m^N \int d\mathbf{x}_1 \cdots d\mathbf{x}_N \hat{\mathcal{P}}_i \{ \chi_1(1) \cdots \chi_N(N) \} \delta(\mathbf{r}_m - \mathbf{R}) s_z(m) \hat{\mathcal{P}}_i \{ \chi_1(1) \cdots \chi_N(N) \} \\
&= \frac{2}{N} \sum_s^N \sum_m^N \int d\mathbf{x}_m \chi_s^*(m) \delta(\mathbf{r}_m - \mathbf{R}) s_z(m) \chi_s(m) \\
&= \frac{2}{N} \sum_m^N \left[\sum_s^{N_\alpha} \int d\mathbf{r}_m \psi_s^{\alpha*}(m) \delta(\mathbf{r}_m - \mathbf{R}) s_z(m) \psi_s^\alpha(m) + \sum_s^{N_\beta} \int d\mathbf{r}_m \psi_s^{\beta*}(m) \delta(\mathbf{r}_m - \mathbf{R}) s_z(m) \psi_s^\beta(m) \right] \\
&= \frac{2}{N} \sum_m^N \left[\frac{1}{2} \sum_s^{N_\alpha} \int d\mathbf{r}_m \psi_s^{\alpha*}(m) \delta(\mathbf{r}_m - \mathbf{R}) \psi_s^\alpha(m) - \frac{1}{2} \sum_s^{N_\beta} \int d\mathbf{r}_m \psi_s^{\beta*}(m) \delta(\mathbf{r}_m - \mathbf{R}) \psi_s^\beta(m) \right] \\
&= \frac{2}{N} N \left[\frac{1}{2} \sum_s^{N_\alpha} \psi_s^{\alpha*}(\mathbf{R}) \psi_s^\alpha(\mathbf{R}) - \frac{1}{2} \sum_s^{N_\beta} \psi_s^{\beta*}(\mathbf{R}) \psi_s^\beta(\mathbf{R}) \right]
\end{aligned}$$

$$\begin{aligned}
&= 2 \left[\frac{1}{2} \rho^\alpha(\mathbf{R}) - \frac{1}{2} \rho^\beta(\mathbf{R}) \right] \\
&= \rho^S(\mathbf{R})
\end{aligned} \tag{3.8.15}$$

where

$$\begin{aligned}
\rho^S(\mathbf{R}) &= \sum_\nu \sum_\mu P_{\nu\mu}^S \phi_\mu^*(\mathbf{R}) \phi_\nu(\mathbf{R}) \\
&= \sum_\nu (\mathbf{P}^S \mathbf{A})_{\nu\nu} \\
&= \text{tr}(\mathbf{P}^S \mathbf{A})
\end{aligned} \tag{3.8.16}$$

3.8.4 Expression for the Fock Matrices

3.8.5 Solution of the Unrestricted SCF Equations

Ex 3.40

$$\begin{aligned}
E_0 &= \sum_a^{N_\alpha} h_{aa}^\alpha + \sum_a^{N_\beta} h_{aa}^\beta + \frac{1}{2} \sum_a^{N_\alpha} \sum_b^{N_\alpha} (J_{ab}^{\alpha\alpha} - K_{ab}^{\alpha\alpha}) + \frac{1}{2} \sum_a^{N_\beta} \sum_b^{N_\beta} (J_{ab}^{\beta\beta} - K_{ab}^{\beta\beta}) + \sum_a^{N_\alpha} \sum_b^{N_\beta} J_{ab}^{\alpha\beta} \\
&= \sum_\mu \sum_\nu P_{\nu\mu}^\alpha H_{\mu\nu}^{\text{core}} + \sum_\mu \sum_\nu P_{\nu\mu}^\beta H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_\mu \sum_\nu \sum_\lambda \sum_\sigma P_{\nu\mu}^\alpha P_{\sigma\lambda}^\alpha [(\mu\nu|\lambda\sigma) - (\mu\sigma|\lambda\nu)] \\
&\quad + \frac{1}{2} \sum_\mu \sum_\nu \sum_\lambda \sum_\sigma P_{\nu\mu}^\beta P_{\sigma\lambda}^\beta [(\mu\nu|\lambda\sigma) - (\mu\sigma|\lambda\nu)] + \sum_\mu \sum_\nu \sum_\lambda \sum_\sigma P_{\nu\mu}^\alpha P_{\sigma\lambda}^\beta (\mu\nu|\lambda\sigma) \\
&= \sum_\mu \sum_\nu P_{\nu\mu}^\alpha \left\{ H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_\lambda \sum_\sigma P_{\sigma\lambda}^\alpha [(\mu\nu|\lambda\sigma) - (\mu\sigma|\lambda\nu)] + \frac{1}{2} \sum_\lambda \sum_\sigma P_{\sigma\lambda}^\beta (\mu\nu|\lambda\sigma) \right\} \\
&\quad + \sum_\mu \sum_\nu P_{\nu\mu}^\beta \left\{ H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_\lambda \sum_\sigma P_{\sigma\lambda}^\beta [(\mu\nu|\lambda\sigma) - (\mu\sigma|\lambda\nu)] + \frac{1}{2} \sum_\lambda \sum_\sigma P_{\sigma\lambda}^\alpha (\mu\nu|\lambda\sigma) \right\} \\
&= \sum_\mu \sum_\nu P_{\nu\mu}^\alpha \left\{ H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_\lambda \sum_\sigma [P_{\sigma\lambda}^T (\mu\nu|\lambda\sigma) - P_{\sigma\lambda}^\alpha (\mu\sigma|\lambda\nu)] \right\} \\
&\quad + \sum_\mu \sum_\nu P_{\nu\mu}^\beta \left\{ H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_\lambda \sum_\sigma [P_{\sigma\lambda}^T (\mu\nu|\lambda\sigma) - P_{\sigma\lambda}^\beta (\mu\sigma|\lambda\nu)] \right\}
\end{aligned} \tag{3.8.17}$$

while

$$F_{\mu\nu}^\alpha = H_{\mu\nu}^{\text{core}} + \sum_\lambda \sum_\sigma [P_{\lambda\sigma}^T (\mu\nu|\sigma\lambda) - P_{\lambda\sigma}^\alpha (\mu\lambda|\sigma\nu)] \tag{3.8.18}$$

$$F_{\mu\nu}^\beta = H_{\mu\nu}^{\text{core}} + \sum_\lambda \sum_\sigma [P_{\lambda\sigma}^T (\mu\nu|\sigma\lambda) - P_{\lambda\sigma}^\beta (\mu\lambda|\sigma\nu)] \tag{3.8.19}$$

thus

$$\begin{aligned}
E_0 &= \sum_\mu \sum_\nu \left\{ P_{\nu\mu}^\alpha \left[\frac{1}{2} H_{\mu\nu}^{\text{core}} + \frac{1}{2} F_{\mu\nu}^\alpha \right] + P_{\nu\mu}^\beta \left[\frac{1}{2} H_{\mu\nu}^{\text{core}} + \frac{1}{2} F_{\mu\nu}^\beta \right] \right\} \\
&= \frac{1}{2} \sum_\mu \sum_\nu [P_{\nu\mu}^T H_{\mu\nu}^{\text{core}} + P_{\nu\mu}^\alpha F_{\mu\nu}^\alpha + P_{\nu\mu}^\beta F_{\mu\nu}^\beta]
\end{aligned} \tag{3.8.20}$$